AUTHOR INDEX VOLUMES 132–135

A

Abe, H., Obinata, T., Minamide, L.S., and Bamburg, J.R. *Xenopus laevis* actin-depolymerizing factor/cofilin: A phosphorylation-regulated protein essential for development. 132:871.
Adam, E.J.H. *See* Chi, Adam, Visser, and Adam.
Adam, S.A. *See* Chi, Adam, Visser, and Adam.
Adams, C.L., Nelson, W.J., and Smith, S.J. Quantitative analysis of cadherin-catenin-actin reorganization during development of cell–cell adhesion. 135:1899.
— *See* Näthke, Adams, Polakis, Sellin, and Nelson.
Adamson, E.D. *See* Huang, Wu, Fan, and Adamson.
Adelstein, R.S. *See* Kelley, Sellers, Gard, Bui, Adelstein, and Baines.
Adesnik, M. *See* Simon, Ivanov, Adesnik, and Sabatini.
Aebi, U. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Aggujaro, D. *See* Borgese, Aggujaro, Carrera, Pietrini, and Bassetti.
Aikawa, M. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Aïssi, F. *See* Vainio, Dunon, Aïssi, Dangy, McNagny, and Imhof.
Aizawa, H., Sutoh, K., and Yahara, I. Overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and cell movement in *Dictyostelium*. 132:335.
Aizawa, S. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Akada, R. *See* Matsui, Matsui, Akada, and Toh-e.
Alabouvette, J. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Al-Awar, O. *See* Field, Al-Awar, Rosenblatt, Wong, Alberts, and Mitchison.
Albers, K.M., Perrone, T.N., Goodness, T.P., Jones, M.E., Green, M.A., and Davis, B.M. Cutaneous overexpression of NT-3 increases sensory and sympathetic neuron number and enhances touch dome and hair follicle innervation. 134:487.
Alberts, B. *See* Field, Al-Awar, Rosenblatt, Wong, Alberts, and Mitchison.
Albright, C.F. *See* Song, Mach, Chen, Reynolds, and Albright.
Alexander, C.M., Howard, E.W., Bissell, M.J., and Werb, Z. Rescue of mammary epithelial cell apoptosis and entactin degradation by a tissue inhibitor of metalloproteinase-1 transgene. 135:1669.
— *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Allan, V.J. *See* Niclas, Allan, and Vale.
Allen, E., Yu, Q.-C., and Fuchs, E. Mice expressing a mutant desmosomal cadherin exhibit abnormalities in desmosomes, proliferation, and epidermal differentiation. 133:1367.
Allen, T.StC. *See* Myers, Goh, Allen, Bucher, and Bogaert.
Allet, B., Hochmann, A., Martinou, I., Berger, A., Missotten, M., Antonsson, B., Sadoul, R., Martinou, J.-C., and Bernasconi, L. Dissecting processing and apoptotic activity of a cysteine protease by mutant analysis. 135:479.
Allis, C.D. *See* Sweet, Jones, and Allis.
Allsopp, R. *See* Henderson, Allsopp, Spector, Wang, and Harley.
Almendro, N. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Alon, R., Fuhlbrigge, R.C., Finger, E.B., and Springer, T.A. Interactions through L-selectin between leukocytes and adherent leukocytes nucleate rolling adhesions on selectins and VCAM-1 in shear flow. 135:849.
— *See* Clark, Alon, and Springer.
— *See* Fuhlbrigge, Alon, Puri, Lowe, and Springer.
— *See* Weber, Alon, Moser, and Springer.
Aloyz, R. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Altruda, F. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Amalric, F. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Amann, K.J. *See* Rybakova, Amann, and Ervasti.
America, T. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Amherdt, M. *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
— *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Anderson, B.L. *See* Goodson, Anderson, Warrick, Pon, and Spudich.
Anderson, K.I., Wang, Y.-L., and Small, J.V. Coordination of protusion and translocation of the keratocyte involves rolling of the cell body. 134:1209.
Anderson, M. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Anderson, P. *See* Maryon, Coronado, and Anderson.
Anderson, R.G.W. *See* Smart, Mineo, and Anderson.
Ando-Akatsuka, Y., Saitou, M., Hirase, T., Kishi, M., Sakakibara, A., Itoh, M., Yonemura, S., Furuse, M., and Tsukita, S. Interspecies diversity of the occludin sequence: cDNA cloning of human, mouse, dog, and rat–kangaroo homologues. 133:43.
Andreassen, P.R. *See* Trielli, Andreassen, Lacroix, and Margolis.
Andrews, N.W. *See* Rodríguez, Samoff, Rioult, Chung, and Andrews.
André, C. *See* Roussel, André, Comai, and Hernandez-Verdun.
Andrés, V., and Walsh, K. Myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis. 132:657.
Angres, B., Barth, A., and Nelson, W.J. Mechanism for transition from initial to stable cell–cell adhesion: Kinetic analysis of E-cadherin–mediated adhesion using a quantitative adhesion assay. 134:549.
Aniento, F., Gu, F., Parton, R.G., and Gruenberg, J. An endosomal βCOP is involved in the pH-dependent formation of transport vesicles destined for late endosomes. 133:29.
Antonsson, B. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
— *See* Di Paolo, Pellier, Catsicas, Antonsson, Catsicas, and Grenningloh.
Arai, K. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Araki, N., Johnson, M.T., and Swanson, J.A. A role for phosphoinositide 3-kinase in the completion of macropinocytosis and phagocytosis by macrophages. 135:1249.
Ardito, T. *See* Devarajan, Stabach, Mann, Ardito, Kashgarian, and Morrow.
Arenas, E. *See* Neveu and Arenas.
Aridor, M. *See* Rowe, Aridor, McCaffery, Plutner, Nuoffer, and Balch.
— *See* Traub, Bannykh, Rodel, Aridor, Balch, and Kornfeld.
Arnoult, C., Zeng, Y., and Florman, H.M. ZP3-dependent activation of sperm cation channels regulates acrosomal secretion during mammalian fertilization. 134:637.
Arredondo, J. *See* Maroto, Arredondo, Goulding, Marco, Bullard, and Cervera.
Arrigo, A.-P. *See* Kretz-Remy, Mehlen, Mirault, and Arrigo.
Artuc, M. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Aruffo, A. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Arvan, P. *See* Kim, Kwon, and Arvan.
Assoian, R.K. *See* Zhu, Ohtsubo, Böhmer, Roberts, and Assoian.
Atsumi, T. *See* Shukunami, Shigeno, Atsumi, Ishizeki, Suzuki, and Hiraki.
Atwal, J. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Aubert, S., Gout, E., Bligny, R., Marty-Mazars, D., Barrieu, F., Alabouvette, J., Marty, F., and Douce, R. Ultrastructural and biochemical characterization of autophagy in higher plant cells subjected to carbon deprivation: Control by the supply of mitochondria with respiratory substrates. 133:1251.
Aubry, J.-M. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Audigier, S. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Auerbach, S. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
Aunis, D. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Austen, K.F. *See* Friend, Ghildyal, Austen, Gurish, Matsumoto, and Stevens.
Austin, C.D., and Shields, D. Formation of nascent secretory vesicles from the *trans*-Golgi network of endocrine cells is inhibited by tyrosine kinase and phosphatase inhibitors. 135:1471.
Auvinen, P. *See* Peränen, Auvinen, Virta, Wepf, and Simons.
Avery, L. *See* Starich, Lee, Panzarella, Avery, and Shaw.

B

Baas, P.W. *See* Yu, Schwei, and Baas.
Babinet, C. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Bacarra, A.E. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Bachant, J.B., and Klymkowsky, M.W. A nontetrameric species is the major soluble form of keratin in *Xenopus* oocytes and rabbit reticulocyte lysates. 132:153.

Bacon, K.B. *See* Campbell, Qin, Bacon, Mackay, and Butcher.
Bader, M.-F. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Baeuerle, P.A. *See* Grimm, Bauer, Baeuerle, and Schulze-Osthoff.
— *See* Pahl, Sester, Burgert, and Baeuerle.
Bagai, S., and Lamb, R.A. Truncation of the COOH-terminal region of the paramyxovirus SV5 fusion protein leads to hemifusion but not complete fusion. 135:73.
Bahrami, S. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Baillat, G. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Baines, I.C. *See* Kelley, Sellers, Gard, Bui, Adelstein, and Baines.
Baird, B. *See* Pierini, Holowka, and Baird.
Bajer, A.S. *See* Khodjakov, Cole, Bajer, and Rieder.
Balch, W.E. *See* Bannykh, Rowe, and Balch.
— *See* Rowe, Aridor, McCaffery, Plutner, Nuoffer, and Balch.
— *See* Traub, Bannykh, Rodel, Aridor, Balch, and Kornfeld.
Balda, M.S., Whitney, J.A., Flores, C., González, S., Cereijido, M., and Matter, K. Functional dissociation of paracellular permeability and transepithelial electrical resistance and disruption of the apical–basolateral intramembrane diffusion barrier by expression of a mutant tight junction membrane protein. 134:1031.
Baldini, G. *See* Scherer, Lederkremer, Williams, Fogliano, Baldini, and Lodish.
Ballotti, R. *See* Bertolotto, Bille, Ortonne, and Ballotti.
Balsamo, J., Leung, T., Ernst, H., Zanin, M.K.B., Hoffman, S., and Lilien, J. Regulated binding of a PTP1B-like phosphatase to N-cadherin: Control of cadherin-mediated adhesion by dephosphorylation of β-catenin. 134:801.
Balzac, F. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Bamburg, J.R. *See* Abe, Obinata, Minamide, and Bamburg.
Bannykh, S.I., Rowe, T., and Balch, W.E. The organization of endoplasmic reticulum export complexes. 135:19.
— *See* Traub, Bannykh, Rodel, Aridor, Balch, and Kornfeld.
Barkalow, K., Witke, W., Kwiatkowski, D.J., and Hartwig, J.H. Coordinated regulation of platelet actin filament barbed ends by gelsolin and capping protein. 134:389.
Barnier, J.V. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Barondes, S.H. *See* Cleves, Cooper, Barondes, and Kelly.
Barrieu, F. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Barth, A. *See* Angres, Barth, and Nelson.
Bartolazzi, A., Nocks, A., Aruffo, A., Spring, F., and Stamenkovic, I. Glycosylation of CD44 is implicated in CD44-mediated cell adhesion to hyaluronan. 132:1199.
Bartoli, M. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Bassetti, M. *See* Borgese, Aggujaro, Carrera, Pietrini, and Bassetti.
Bastiaens, P.I.H. *See* Majoul, Bastiaens, and Söling.
Bastos, R., Lin, A., Enarson, M., and Burke, B. Targeting and function in mRNA export of nuclear pore complex protein Nup153. 134:1141.
Bauer, K.D. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Bauer, M.K.A. *See* Grimm, Bauer, Baeuerle, and Schulze-Osthoff.
Baurén, G., Jiang, W.-Q., Bernholm, K., Gu, F., and Wieslander, L. Demonstration of a dynamic, transcription-dependent organization of pre-mRNA splicing factors in polytene nuclei. 133:929.
Bayard, F. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Bazett-Jones, D.P. *See* Rattner, Hendzel, Furbee, Muller, and Bazett-Jones.
Becher, M.W. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
Becker, E. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Becker, J.C. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Becker, K. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Beckerle, M.C. *See* Stronach, Siegrist, and Beckerle.
Beggah, A. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Behrend, M. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Belkin, A.M., Zhidkova, N.I., Balzac, F., Altruda, F., Tomatis, D., Maier, A., Tarone, G., Koteliansky, V.E., and Burridge, K. β1D integrin displaces the β1A isoform in striated muscles: Localization at junctional structures and signaling potential in nonmuscle cells. 132:211.
Belliveau, D.J. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Bellon, P.L. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Belmont, A.S. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Benbow, R.M. *See* Fang and Benbow.
Bendayan, M. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Bender, A. *See* Bender, Lo, Lee, Kokojan, Peterson, and Bender.
Bender, L., Lo, H.S., Lee, H., Kokojan, V., Peterson, J., and Bender, A. Associations among PH and SH3 domain–containing proteins and Rho-type GTPases in yeast. 133:879.
Benedetti, E.L. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
Benian, G.M., Tinley, T.L., Tang, X., and Borodovsky, M. The *Caenorhabditis elegans* gene *unc-89*, required for muscle M-line assembly, encodes a giant modular protein composed of Ig and signal transduction domains. 132:835.
Bénichou, J.-C. *See* Imaizumi-Scherrer, Faust, Bénichou, Hellio, and Weiss.
Benner, A. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Bennett, V. *See* Davis, Lambert, and Bennett.
Ben-Ze'ev, A. *See* Simcha, Geiger, Yehuda-Levenberg, Salomon, and Ben-Ze'ev.
Berger, A. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Berger, P. *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
Berglund, E.B. *See* Engert, Berglund, and Rosenthal.
Bernabéu, C. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Bernard, A. *See* Deckert, Ticchioni, and Bernard.
Bernasconi, L. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Bernholm, K. *See* Baurén, Jiang, Bernholm, Gu, and Wieslander.
Bertin, E. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Bertolotto, C., Bille, K., Ortonne, J.-P., and Ballotti, R. Regulation of tyrosinase gene expression by cAMP in B16 melanoma cells involves two CATGTG motifs surrounding the TATA box: Implication of the microphthalmia gene product. 134:747.
Berton, G. *See* Lowell, Fumagalli, and Berton.
Beug, H. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Beyer, A. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Biggins, S., Ivanovska, I., and Rose, M.D. Yeast ubiquitin-like genes are involved in duplication of the microtubule organizing center. 133:1331.
Bille, K. *See* Bertolotto, Bille, Ortonne, and Ballotti.
Bingham, J.B. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Birchmeier, C. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Birchmeier, W. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
— *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Bird, C. *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Birk, D.E. *See* Marchant, Hahn, Linsenmayer, and Birk.
Bishop, J.D. *See* Van Haastert, Bishop, and Gomer.
Bissell, M.J. *See* Alexander, Howard, Bissell, and Werb.
— *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell. and Werb.
Bjorkman, J. *See* Gould, Kalish, Morrell, Bjorkman, Urquhart, and Crane.
Black, C. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Blandino, G. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Blank, P.S. *See* Vogel, Blank, and Zimmerberg.
Blau, H.M. *See* Yang, Rando, Mohler, Rayburn, Blau, and Hynes.
Blessing, M., Schirmacher, P., and Kaiser, S. Overexpression of bone morphogenetic protein-6 (BMP-6) in the epidermis of transgenic mice: Inhibition or stimulation of proliferation depending on the pattern of transgene expression and formation of psoriatic lesions. 135:227.
Bligny, R. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Blobel, C.P. *See* Weskamp, Krätzschmar, Reid, and Blobel.
Blobel, G. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.

— *See* Erdmann and Blobel.
— *See* Matunis, Coutavas, and Blobel.
— *See* Nehrbass, Rout, Maguire, Blobel, and Wozniak.
Block, G.D., Locker, J., Bowen, W.C., Petersen, B.E., Katyal, S., Strom, S.C., Riley, T., Howard, T.A., and Michalopoulos, G.K. Population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF and TGFα in a chemically defined (HGM) medium. 132: 1133.
Bloemendal, H. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
Blumenthal, R., Sarkar, D.P., Durell, S., Howard, D.E., and Morris, S.J. Dilation of the influenza hemagglutinin fusion pore revealed by the kinetics of individual cell–cell fusion events. 135:63.
Bogaert, T. *See* Myers, Goh, Allen, Bucher, and Bogaert.
Böhmer, R.M. *See* Zhu, Ohtsubo, Böhmer, Roberts, and Assoian.
Bonaldo, P. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
Bonifacino, J.S. *See* Marks, Woodruff, Ohno, and Bonifacino.
Borgese, N., Aggujaro, D., Carrera, P., Pietrini, G., and Bassetti, M. A role for N-myristoylation in protein targeting: NADH-cytochrome b_5 reductase requires myristic acid for association with outer mitochondrial but not ER membranes. 135:1501.
Borisy, G.G. *See* Svitkina, Verkhovsky, and Borisy.
— *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
— *See* Zhai, Kronebusch, Simon, and Borisy.
Bornslaeger, E.A., Corcoran, C.M., Stappenbeck, T.S., and Green, K.J. Breaking the connection: Displacement of the desmosomal plaque protein desmoplakin from cell–cell interfaces disrupts anchorage of intermediate filament bundles and alters intercellular junction assembly. 134:985.
Borodovsky, M. *See* Benian, Tinley, Tang, and Borodovsky.
Borst, P. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
Bos-Mikich, A. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.
Bossi, G. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Boudier, J.-A. *See* Paillart, Boudier, Boudier, Rochat, Couraud, and Dargent.
Boudier, J.-L. *See* Paillart, Boudier, Boudier, Rochat, Couraud, and Dargent.
Bouillet, P. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Boukamel, R. *See* De Arcangelis, Neuville, Boukamel, Lefebvre, Kedinger, and Simon-Assmann.
Bourgeois, J.-P. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Bowen, W.C. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Bradl, J. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Bradley, A. *See* Milner, Weitzer, Tran, Bradley, and Capetanaki.
Brady, B. *See* Wein, Foss, Brady, and Cande.
Braghetta, P., Fabbro, C., Piccolo, S., Marvulli, D., Bonaldo, P., Volpin, D., and Bressan, G.M. Distinct regions control transcriptional activation of the α1(VI) collagen promoter in different tissues of transgenic mice. 135:1163.
Brandenberger, R., Kammerer, R.A., Engel, J., and Chiquet, M. Native chick laminin-4 containing the β2 chain (s-laminin) promotes motor axon growth. 135:1583.
Brandin, E. *See* Lue, Brandin, Chan, and Branton.
Branton, D. *See* Lue, Brandin, Chan, and Branton.
Bravo, N.S. *See* Paladini, Takahashi, Bravo, and Coulombe.
Bremser, M. *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Brenner, J. *See* Prestle, Pfizenmaier, Brenner, and Johannes.
Bressan, G.M. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
Bretaudiere, J.-P. *See* Schroeter, Bretaudiere, Sass, and Goldstein.
Bridge, E., Riedel, K.-U., Johansson, B.-M., and Pettersson, U. Spliced exons of adenovirus late RNAs colocalize with snRNP in a specific nuclear domain. 135:303.
Brieher, W.M., Yap, A.S., and Gumbiner, B.M. Lateral dimerization is required for the homophilic binding activity of C-cadherin. 135:487.
Bright, N.A. *See* Simpson, Bright, West, Newman, Darnell, and Robinson.
Brinkmann, V. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
— *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Brizzio, V., Gammie, A.E., Nijbroek, G., Michaelis, S., and Rose, M.D. Cell fusion during yeast mating requires high levels of a-factor mating pheromone. 135:1727.
Brock, J., Midwinter, K., Lewis, J., and Martin, P. Healing of incisional wounds in the embryonic chick wing bud: Characterization of the actin purse-string and demonstration of a requirement for Rho activation. 135:1097.
Brodsky, F.M. *See* Wilde and Brodsky.
Brodsky, J.L. *See* McCracken and Brodsky.
Bron, R. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Bronner, S. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Brown, E.J. *See* Gao, Lindberg, Dimitry, Brown, and Frazier.
— *See* Lindberg, Gresham, Reinhold, and Brown.
Brown, H.A. *See* Ktistakis, Brown, Waters, Sternweis, and Roth.
Brown, J.D. *See* Ng, Brown, and Walter.
Brown, M.C., Perrotta, J.A., and Turner, C.E. Identification of LIM3 as the principal determinant of paxillin focal adhesion localization and characterization of a novel motif on paxillin directing vinculin and focal adhesion kinase binding. 135:1109.
Brown, N.H. *See* Martin-Bermudo and Brown.
Brown, S.D. *See* Zhang, Lee, Chen, Brown, and Hsieh.
Bruckner, P. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Brunton, V.G. *See* Fincham, Unlu, Brunton, Pitts, Wyke, and Frame.
Bryan, J. *See* Tao, Edwards, Tubb, Wang, Bryan, and McCrea.
Bryant, P.J. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
— *See* Woods, Hough, Peel, Callaini, and Bryant.
Bucher, E.A. *See* Myers, Goh, Allen, Bucher, and Bogaert.
Buchstaller, A., Kunz, S., Berger, P., Kunz, B., Ziegler, U., Rader, C., and Sonderegger, P. Cell adhesion molecules NgCAM and axonin-1 form heterodimers in the neuronal membrane and cooperate in neurite outgrowth promotion. 135:1593.
Bui, D. *See* Kelley, Sellers, Gard, Bui, Adelstein, and Baines.
Bulinski, J.C. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
Bullard, B. *See* Maroto, Arredondo, Goulding, Marco, Bullard, and Cervera.
Burger, K.N.J., van der Bijl, P., and van Meer, G. Topology of sphingolipid galactosyltransferases in ER and Golgi: Transbilayer movement of monohexosyl sphingolipids is required for higher glycosphingolipid biosynthesis. 133:15.
Burgert, H.-G. *See* Pahl, Sester, Burgert, and Baeuerle.
Burgeson, R.E. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Burke, B. *See* Bastos, Lin, Enarson, and Burke.
Burkhardt, H. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Burridge, K. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
— *See* Chrzanowska-Wodnicka and Burridge.
Busetto, S. *See* Menegazzi, Busetto, Dri, Cramer, and Patriarca.
Bush, G.L., and Meyer, D.I. The refolding activity of the yeast heat shock proteins Ssa1 and Ssa2 defines their role in protein translocation. 135:1229.
Busslinger, M. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Butcher, E.C. *See* Campbell, Qin, Bacon, Mackay, and Butcher.
Byers, B. *See* Sundberg, Goetsch, Byers, and Davis.

C

Cabral, F. *See* Gonzalez-Garay and Cabral.
Cabrera, N., Díaz-Rodríguez, E., Becker, E., Martín-Zanca, D., and Pandiella, A. TrkA receptor ectodomain cleavage generates a tyrosine-phosphorylated cell-associated fragment. 132:427.
Caffarelli, A.D. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Calafat, J. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Calfon, M. *See* Constam, Calfon, and Robertson.
Callaini, G. *See* Woods, Hough, Peel, Callaini, and Bryant.
Calothy, G. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Cameron, M. *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Campbell, J.J., Qin, S., Bacon, K.B., Mackay, C.R., and Butcher, E.C. Biology of chemokine and classical chemoattractant receptors: Differential requirements for adhesion-triggering versus chemotactic responses in lymphoid cells. 134:255.
Campbell, K.P. *See* Rafael, Cox, Corrado, Jung, Campbell, and Chamberlain.
Campenot, R.B., Lund, K., and Senger, D.L. Delivery of newly synthesized tubulin to rapidly growing distal axons of rat sympathetic neurons in compartmented cultures. 135:701.

Cande, W.Z. *See* Wein, Foss, Brady, and Cande.
Capetanaki, Y. *See* Milner, Weitzer, Tran, Bradley, and Capetanaki.
Cardone, M.H., Smith, B.L., Mennitt, P.A., Mochly-Rosen, D., Silver, R.B., and Mostov, K.E. Signal transduction by the polymeric immunoglobulin receptor suggests a role in regulation of receptor transcytosis. 133:997.
Carey, K.L., Richards, S.A., Lounsbury, K.M., and Macara, I.G. Evidence using a green fluorescent protein-glucocorticoid receptor chimera that the RAN/TC4 GTPase mediates an essential function independent of nuclear protein import. 133:985. Correction. 135:1193.
—*See* Richards, Lounsbury, Carey, and Macara.
Carey, T.E. *See* Islam, Carey, Wolf, Wheelock, and Johnson.
Carmo-Fonseca, M. *See* Jordan, Mannervik, Tora, and Carmo-Fonseca.
Carnes, D.K. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Carrera, P. *See* Borgese, Aggujaro, Carrera, Pietrini, and Bassetti.
Carroll, J. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.
Carter, W.G. *See* Xia, Gil, and Carter.
Cass, D. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Castets, F., Bartoli, M., Barnier, J.V., Baillat, G., Salin, P., Moqrich, A., Bourgeois, J.-P., Denizot, F., Rougon, G., Calothy, G., and Monneron, A. A novel calmodulin-binding protein, belonging to the WD-repeat family, is localized in dendrites of a subset of CNS neurons. 134:1051.
Catlett, N.L. *See* Hill, Catlett, and Weisman.
Catsicas, M. *See* Di Paolo, Pellier, Catsicas, Antonsson, Catsicas, and Grenningloh.
Catsicas, S. *See* Di Paolo, Pellier, Catsicas, Antonsson, Catsicas, and Grenningloh.
Catterall, W.A. *See* Sakurai, Westenbroek, Rettig, Hell, and Catterall.
Cereijido, M. *See* Balda, Whitney, Flores, González, Cereijido, and Matter.
Cervera, M. *See* Maroto, Arredondo, Goulding, Marco, Bullard, and Cervera.
Chalfie, M. *See* Lai, Hong, Kinnell, Chalfie, and Driscoll.
Chamberlain, J.S. *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
— *See* Rafael, Cox, Corrado, Jung, Campbell, and Chamberlain.
Chambon, P. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Champliaud, M.-F., Lunstrum, G.P., Rousselle, P., Nishiyama, T., Keene, D.R., and Burgeson, R.E. Human amnion contains a novel laminin variant, laminin 7, which like laminin 6, covalently associates with laminin 5 to promote stable epithelial–stromal attachment. 132:1189.
Chan, E.P. *See* Lue, Brandin, Chan, and Branton.
Chan, P.-Y. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Chang, A. *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Chang, L.Y. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Chang, S. *See* DeBernardo and Chang.
Chan-Hui, P.-Y. *See* Frisch, Vuori, Ruoslahti, and Chan-Hui.
Charness, M.E. *See* Ramanathan, Wilkemeyer, Mittal, Perides, and Charness.
Chasserot-Golaz, S., Vitale, N., Sagot, I., Delouche, B., Dirrig, S., Pradel, L.-A., Henry, J.-P., Aunis, D., and Bader, M.-F. Annexin II in exocytosis: Catecholamine secretion requires the translocation of p36 to the subplasmalemmal region in chromaffin cells. 133:1217.
Chavez, R.A., Miller, S.G., and Moore, H.-P.H. A biosynthetic regulated secretory pathway in constitutive secretory cells. 133:1177.
Chen, A.D. *See* Zhang, Lee, Chen, Brown, and Hsieh.
Chen, C.-Y. *See* Song, Mach, Chen, Reynolds, and Albright.
Chen, P., Murphy-Ullrich, J.E., and Wells, A. A role for gelsolin in actuating epidermal growth factor receptor–mediated cell motility. 134:689.
Cheresh, D.A. *See* Lewis, Cheresh, and Schwartz.
— *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Chi, N.C., Adam, E.J.H., Visser, G.D., and Adam, S.A. RanBP1 stabilizes the interaction of Ran with p97 in nuclear protein import. 135:559.
Chilcoat, N.D., Melia, S.M., Haddad, A., and Turkewitz, A.P. Granule lattice protein 1 (Grl1p), an acidic, calcium-binding protein in *Tetrahymena thermophila* dense-core secretory granules, influences granule size, shape, content organization, and release but not protein sorting or condensation. 135:1775.
Chin, J.R. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Chin, L.-S. *See* Ryan, Li, Chin, Greengard, and Smith.
Chiquet, M. *See* Brandenberger, Kammerer, Engel, and Chiquet.
— *See* Hagios, Koch, Spring, Chiquet, and Chiquet-Ehrismann.
Chiquet-Ehrismann, R. *See* Hagios, Koch, Spring, Chiquet, and Chiquet-Ehrismann.
Chishti, A.H. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
Chitaev, N.A., Leube, R.E., Troyanovsky, R.B., Eshkind, L.G., Franke, W.W., and Troyanovsky, S.M. The binding of plakoglobin to desmosomal cadherins: Patterns of binding sites and topogenic potential. 133:359.
Chong, L. *See* Ludérus, van Steensel, Chong, Sibon, Cremers, and de Lange.
Chou, Y.H. *See* Goldman, Khuon, Chou, Opal, and Steinert.
Chrzanowska-Wodnicka, M., and Burridge, K. Rho-stimulated contractility drives the formation of stress fibers and focal adhesions. 133:1403.
Chu, G.C. *See* Tansey, Chu, and Merlie.
Chuang, J.S., and Schekman, R.W. Differential trafficking and timed localization of two chitin synthase proteins, Chs2p and Chs3p. 135:597. Correction. 135:1925.
Chun, J. *See* Hecht, Weiner, Post, and Chun.
Chung, A. *See* Rodríguez, Samoff, Rioult, Chung, and Andrews.
Cimino, L. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Clark, R.A., Alon, R., and Springer, T.A. CD44 and hyaluronan-dependent rolling interactions of lymphocytes on tonsillar stroma. 134:1075.
Clark, R.A.F. *See* Xu and Clark.
— *See* Xu, Zutter, Santoro, and Clark.
Clase, K.L. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
Clausen, H. *See* Sönnichsen, Watson, Clausen, Misteli, and Warren.
Clayton, C. *See* Häusler, Stierhof, Wirtz, and Clayton.
Clayton, D.A. *See* Davis and Clayton.
Clement, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Clemson, C.M., McNeil, J.A., Willard, H.F., and Lawrence, J.B. XIST RNA paints the inactive X chromosome at interphase: Evidence for a novel RNA involved in nuclear/chromosome structure. 132:259.
Cleveland, D.W. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
— *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Cleves, A.E., Cooper, D.N.W., Barondes, S.H., and Kelly, R.B. A new pathway for protein export in *Saccharomyces cerevisiae.* 133:1017.
Clift, S.M. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Coen, S. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Cole, D.G. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Cole, N.M. *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
Cole, R.W. *See* Khodjakov, Cole, Bajer, and Rieder.
Coles, H.S.R. *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
Colgan, S.P. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Collart, M.A. *See* Irminger-Finger, Hurt, Roebuck, Collart, and Edelstein.
Collas, P., Courvalin, J.-C., and Poccia, D. Targeting of membranes to sea urchin sperm chromatin is mediated by a lamin B receptor-like integral membrane protein. 135:1715.
Colucci-Guyon, E. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Comai, L. *See* Roussel, André, Comai, and Hernandez-Verdun.
Compton, D.A. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Compton, M.T. *See* Hungerford, Compton, Matter, Hoffstrom, and Otey.
Condeelis, J. *See* Liu, Tang, Edmonds, Murray, Levin, and Condeelis.
— *See* Murray, Edmonds, Liu, and Condeelis.
Condon, P.J. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Connelly, P. *See* Tilney, Connelly, Smith, and Guild.
Constam, D.B., Calfon, M., and Robertson, E.J. SPC4, SPC6, and the novel protease SPC7 are coexpressed with bone morphogenetic proteins at distinct sites during embryogenesis. 134:181.
Coonrod, A. *See* Li, Coonrod, and Horwitz.
Cooper, A.A., and Stevens, T.H. Vps10p cycles between the late-Golgi and prevacuolar compartments in its function as the sorting receptor for multiple yeast vacuolar hydrolases. 133:529.
Cooper, D.N.W. *See* Cleves, Cooper, Barondes, and Kelly.
Cooper, E. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.

Cooper, J.A., and Kiehart, D.P. Septins may form a ubiquitous family of cytoskeletal filaments. 134:1345.
— *See* Hopmann, Cooper, and Miller.
— *See* Schafer, Jennings, and Cooper.
— *See* Waddle, Karpova, Waterston, and Cooper.
Cooper, J.B. *See* Prekeris, Mayhew, Cooper, and Terrian.
Corada, M. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
Corbett, A.H. *See* Koepp, Wong, Corbett, and Silver.
Corcoran, C.M. *See* Bornslaeger, Corcoran, Stappenbeck, and Green.
Cordey, A.-S. *See* Spertini, Cordey, Monai, Giuffrè, and Schapira.
Coronado, R. *See* Maryon, Coronado, and Anderson.
Corrado, K., Rafael, J.A., Mills, P.L., Cole, N.M., Faulkner, J.A., Wang, K., and Chamberlain, J.S. Transgenic *mdx* mice expressing dystrophin with a deletion in the actin-binding domain display a "mild becker" phenotype. 134:873.
— *See* Rafael, Cox, Corrado, Jung, Campbell, and Chamberlain.
Corry, D. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Corvera, S. *See* Shpetner, Joly, Hartley, and Corvera.
Coulombe, P.A. *See* Paladini, Takahashi, Bravo, and Coulombe.
Couraud, F. *See* Paillart, Boudier, Boudier, Rochat, Couraud, and Dargent.
Courvalin, J.-C. *See* Collas, Courvalin, and Poccia.
Coutavas, E. *See* Matunis, Coutavas, and Blobel.
Cowan, N.J. *See* Lewis, Tian, Vainberg, and Cowan.
Cox, G.A. *See* Rafael, Cox, Corrado, Jung, Campbell, and Chamberlain.
Cox, R.T., Kirkpatrick, C., and Peifer, M. Armadillo is required for adherens junction assembly, cell polarity, and morphogenesis during *Drosophila* embryogenesis. 134:133.
Craig, E. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Craig, R. *See* Xu, Harder, Uman, and Craig.
Craig, R.A. *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Cramer, R. *See* Menegazzi, Busetto, Dri, Cramer, and Patriarca.
Crane, D.I. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
— *See* Gould, Kalish, Morrell, Bjorkman, Urquhart, and Crane.
Crawford, T.O. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
— *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Cregg, J.M. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Cremer, C. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Cremer, T. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
— *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
— *See* Scherthan, Weich, Schwegler, Härle, Heyting, and Cremer.
Cremers, F.F.M. *See* Ludérus, van Steensel, Chong, Sibon, Cremers, and de Lange.
Crescenzi, M. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Creutz, C.E., Snyder, S.L., Daigle, S.N., and Redick, J. Identification, localization, and functional implications of an abundant nematode annexin. 132:1079.
Cumberledge, S. *See* Reichsman and Cumberledge.
Curriden, S.A. *See* Deng, Curriden, Wang, Rosenberg, and Loskutoff.
Cutler, D.F. *See* Norcott, Solari, and Cutler.

D

Daigle, S.N. *See* Creutz, Snyder, Daigle, and Redick.
Daily, D.R. *See* Debec, Kalpin, Daily, McCallum, Rothwell, and Sullivan.
Dallai, R. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Daneholt, B. *See* Visa, Izaurralde, Ferreira, Daneholt, and Mattaj.
Dangy, J.-P. *See* Vainio, Dunon, Aïssi, Dangy, McNagny, and Imhof.
Danieli, T., Pelletier, S.L., Henis, Y.I., and White, J.M. Membrane fusion mediated by the influenza virus hemagglutinin requires the concerted action of at least three hemagglutinin trimers. 133:559.
Daniell, L. *See* Takei, Mundigl, Daniell, and De Camilli.
Danpure, C.J. *See* Leiper, Oatey, and Danpure.
Daou, M.-C. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Dargent, B. *See* Paillart, Boudier, Boudier, Rochat, Couraud, and Dargent.
Darnell, R.B. *See* Simpson, Bright, West, Newman, Darnell, and Robinson.
David, G. *See* Mertens, van der Schueren, van den Berghe, and David.
— *See* Steinfeld, Van Den Berghe, and David.
Davidson, A.R. *See* Hong, Davidson, and Kaiser.
Davies, P. *See* Vincent, Rosado, and Davies.
Davies, T.J. *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
Davis, A.F., and Clayton, D.A. In situ localization of mitochondrial DNA replication in intact mammalian cells. 135:883.
Davis, B.M. *See* Albers, Perrone, Goodness, Jones, Green, and Davis.
Davis, J. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Davis, J.Q., Lambert, S., and Bennett, V. Molecular composition of the node of Ranvier: Identification of ankyrin-binding cell adhesion molecules neurofascin (mucin+/third FNIII domain−) and NrCAM at nodal axon segments. 135:1355.
Davis, N.G. *See* Roth and Davis.
Davis, T.N. *See* Friedman, Sundberg, Huang, and Davis.
— *See* Sundberg, Goetsch, Byers, and Davis.
De Arcangelis, A., Neuville, P., Boukamel, R., Lefebvre, O., Kedinger, M., and Simon-Assmann, P. Inhibition of laminin α1-chain expression leads to alteration of basement membrane assembly and cell differentiation. 133:417.
Debec, A., Kalpin, R.F., Daily, D.R., McCallum, P.D., Rothwell, W.F., and Sullivan, W. Live analysis of free centrosomes in normal and aphidicolin-treated *Drosophila* embryos. 134:103.
DeBernardo, A.P., and Chang, S. Heterophilic interactions of DM-GRASP: GRASP-NgCAM interactions involved in neurite extension. 133:657.
De Bie, I., Marcinkiewicz, M., Malide, D., Lazure, C., Nakayama, K., Bendayan, M., and Seidah, N.G. The isoforms of proprotein convertase PC5 are sorted to different subcellular compartments. 135:1261.
De Camilli, P. *See* Takei, Mundigl, Daniell, and De Camilli.
Décimo, D. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Deckert, M., Ticchioni, M., and Bernard, A. Endocytosis of GPI-anchored proteins in human lymphocytes: Role of glycolipid-based domains, actin cytoskeleton, and protein kinases. 133:791.
Deckwerth, T.L. *See* Deshmukh, Vasilakos, Deckwerth, Lampe, Shivers, and Johnson.
de Crombrugghe, B. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Deiner, M. *See* Ferns, Deiner, and Hall.
Dejana, E. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
de Lange, T. *See* Ludérus, van Steensel, Chong, Sibon, Cremers, and de Lange.
Del Maschio, A., Zanetti, A., Corada, M., Rival, Y., Ruco, L., Lampugnani, M.G., and Dejana, E. Polymorphonuclear leukocyte adhesion triggers the disorganization of endothelial cell-to-cell adherens junctions. 135:497.
Delouche, B. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
De Lozanne, A. *See* Larochelle, Vithalani, and De Lozanne.
DeMarais, A.A. *See* Torres, Yang-Snyder, Purcell, DeMarais, McGrew, and Moon.
Demaurex, N., Downey, G.P., Waddell, T.K., and Grinstein, S. Intracellular pH regulation during spreading of human neutrophils. 133:1391.
Dembinsky, A., Rubin, H., and Ravid, S. Chemoattractant-mediated increases in cGMP induce changes in *Dictyostelium* myosin II heavy chain–specific protein kinase C activities. 134:911.
Deng, G., Curriden, S.A., Wang, S., Rosenberg, S., and Loskutoff, D.J. Is plasminogen activator inhibitor-1 the molecular switch that governs urokinase receptor-mediated cell adhesion and release? 134:1571.
Denizot, F. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Denker, S.P., McCaffery, J.M., Palade, G.E., Insel, P.A., and Farquhar, M.G. Differential distribution of α subunits and βγ subunits of heterotrimeric G proteins on Golgi membranes of the exocrine pancreas. 133:1027.
Derynck, R. *See* Erlebacher and Derynck.
Deshmukh, M., Vasilakos, J., Deckwerth, T.L., Lampe, P.A., Shivers, B.D., and Johnson, E.M., Jr. Genetic and metabolic status of NGF-deprived sympathetic neurons saved by an inhibitor of ICE family proteases. 135:1341.
DeSimone, D.W. *See* Ramos and DeSimone.

Devarajan, P., Stabach, P.R., Mann, A.S., Ardito, T., Kashgarian, M., and Morrow, J.S. Identification of a small cytoplasmic ankyrin (Ank_{G119}) in the kidney and muscle that binds βΙΣ* spectrin and associates with the Golgi apparatus. 133:819.
de Vree, W.J.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Díaz-Rodríguez, E. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Dice, J.F. *See* Hayes and Dice.
Didenko, V.V., and Hornsby, P.J. Presence of double-strand breaks with single-base 3′ overhangs in cells undergoing apoptosis but not necrosis. 135:1369.
Dietrich, J., Neisig, A., Hou, X., Wegener, A.-M.K., Gajhede, M., and Geisler, C. Role of CD3γ in T cell receptor assembly. 132:299.
Dietzel, S. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Dimitry, J.M. *See* Gao, Lindberg, Dimitry, Brown, and Frazier.
Dingwall, C. *See* Palacios, Weis, Klebe, Mattaj, and Dingwall.
Di Paolo, G., Pellier, V., Catsicas, M., Antonsson, B., Catsicas, S., and Grenningloh, G. The phosphoprotein stathmin is essential for nerve growth factor–stimulated differentiation. 133:1383.
Dirrig, S. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Dissanayake, S. *See* Dubreuil, MacVicar, Dissanayake, Liu, Homer, and Hortsch.
Distel, B., Erdmann, R., Gould, S.J., Blobel, G., Crane, D.I., Cregg, J.M., Dodt, G., Fujiki, Y., Goodman, J.M., Just, W.W., Kiel, J.A.K.W., Kunau, W.-H., Lazarow, P.B., Mannaerts, G.P., Moser, H.W., Osumi, T., Rachubinski, R.A., Roscher, A., Subramani, S., Tabak, H.F., Tsukamoto, T., Valle, D., van der Klei, I., van Veldhoven, P.P., and Veenhuis, M. A unified nomenclature for peroxisome biogenesis factors. 135:1.
— *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Dittié, A.S., Hajibagheri, N., and Tooze, S.A. The AP-1 adaptor complex binds to immature secretory granules from PC12 cells, and is regulated by ADP-ribosylation factor. 132:523.
Dittmer, F. *See* Kasper, Dittmer, von Figura, and Pohlmann.
Dodt, G., and Gould, S.J. Multiple PEX genes are required for proper subcellular distribution and stability of Pex5p, the PTS1 receptor: Evidence that PTS1 protein import is mediated by a cycling receptor. 135:1763.
— *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Dogterom, M., Félix, M.-A., Guet, C.C., and Leibler, S. Influence of M-phase chromatin on the anisotropy of microtubule asters. 133:125.
Dollé, P. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Domingo, D. *See* Odorizzi, Pearse, Domingo, Trowbridge, and Hopkins.
Donaldson, A.D., and Kilmartin, J.V. Spc42p: A phosphorylated component of the *S. cerevisiae* spindle pole body (SPB) with an essential function during SPB duplication. 132:887.
Donaldson, J.G. *See* Radhakrishna, Klausner, and Donaldson.
Dooley, T.P. *See* Ruhrberg, Hajibagheri, Simon, Dooley, and Watt.
Douce, R. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Dowling, J., Yu, Q.-C., and Fuchs, E. β4 integrin is required for hemidesmosome formation, cell adhesion and cell survival. 134:559.
Downey, G.P. *See* Demaurex, Downey, Waddell, and Grinstein.
— *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Doye, V. *See* Zabel, Doye, Tekotte, Wepf, Grandi, and Hurt.
Dreyfuss, G. *See* Nakielny and Dreyfuss.
Dri, P. *See* Menegazzi, Busetto, Dri, Cramer, and Patriarca.
Driscoll, M. *See* Lai, Hong, Kinnell, Chalfie, and Driscoll.
Dubreuil, R.R., MacVicar, G., Dissanayake, S., Liu, C., Homer, D., and Hortsch, M. Neuroglian-mediated cell adhesion induces assembly of the membrane skeleton at cell contact sites. 133:647.
Duffus, W.A. *See* Kielian, Klimjack, Ghosh, and Duffus.
Dunia, I., Smit, J.J.M., van der Valk, M.A., Bloemendal, H., Borst, P., and Benedetti, E.L. Human multidrug resistance 3–P-glycoprotein expression in transgenic mice induces lens membrane alterations leading to cataract. 132:701.
Dunn, K.W. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
Dunon, D. *See* Vainio, Dunon, Aïssi, Dangy, McNagny, and Imhof.
Dupouey, P. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Durell, S. *See* Blumenthal, Sarkar, Durell, Howard, and Morris.
Dusseljee, S. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Dustin, M.L., Ferguson, L.M., Chan, P.-Y., Springer, T.A., and Golan, D.E. Visualization of CD2 interaction with LFA-3 and determination of the two-dimensional dissociation constant for adhesion receptors in a contact area. 132:465.
Dvorak, J.A. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Dyer, J.M., McNew, J.A., and Goodman, J.M. The sorting sequence of the peroxisomal integral membrane protein PMP47 is contained within a short hydrophilic loop. 133:269.
— *See* Marshall, Dyer, Quick, and Goodman.

E

Eady, R.A.J. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Eaton, S., Wepf, R., and Simons, K. Roles for Rac1 and Cdc42 in planar polarization and hair outgrowth in the wing of *Drosophila.* 135:1277.
Eberspaecher, H. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Echeverri, C.J., Paschal, B.M., Vaughan, K.T., and Vallee, R.B. Molecular characterization of the 50-kD subunit of dynactin reveals function for the complex in chromosome alignment and spindle organization during mitosis. 132:617.
Edelstein, S.J. *See* Irminger-Finger, Hurt, Roebuck, Collart, and Edelstein.
Edidin, M. *See* Hannan and Edidin.
Edmonds, B.T. *See* Liu, Tang, Edmonds, Murray, Levin, and Condeelis.
— *See* Murray, Edmonds, Liu, and Condeelis.
Edwards, R.A. *See* Tao, Edwards, Tubb, Wang, Bryan, and McCrea.
Egelhoff, T.T. *See* Kolman, Futey, and Egelhoff.
Eils, R., Dietzel, S., Bertin, E., Schröck, E., Speicher, M.R., Ried, T., Robert-Nicoud, M., Cremer, C., and Cremer, T. Three-dimensional reconstruction of painted human interphase chromosomes: Active and inactive X chromosome territories have similar volumes but differ in shape and surface structure. 135:1427.
Ekblom, P. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Elgersma, Y., Kwast, L., Klein, A., Voorn-Brouwer, T., van den Berg, M., Metzig, B., America, T., Tabak, H.F., and Distel, B. The SH3 domain of the *Saccharomyces cerevisiae* peroxisomal membrane protein Pex13p functions as a docking site for Pex5p, a mobile receptor for the import of PTS1-containing proteins. 135:97.
Elia, L., and Marsh, L. Role of the ABC transporter Ste6 in cell fusion during yeast conjugation. 135:741.
Ellis-Davies, G. *See* Wilding, Wright, Patel, Ellis-Davies, and Whitaker.
Emr, S.D. *See* Wendland, McCaffery, Xiao, and Emr.
Enarson, M. *See* Bastos, Lin, Enarson, and Burke.
Engel, J. *See* Brandenberger, Kammerer, Engel, and Chiquet.
Engert, J.C., Berglund, E.B., and Rosenthal, N. Proliferation precedes differentiation in IGF-I–stimulated myogenesis. 135:431.
Engvall, E. *See* Vachon, Loechel, Xu, Wewer, and Engvall.
— *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Ensergueix, D. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Erdmann, R., and Blobel, G. Identification of Pex13p, a peroxisomal membrane receptor for the PTS1 recognition factor. 135:111.
— *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Erdmann, S., Müller, W., Bahrami, S., Vornehm, S.I., Mayer, H., Bruckner, P., von der Mark, K., and Burkhardt, H. Differential effects of parathyroid hormone fragments on collagen gene expression in chondrocytes. 135:1179.
Erickson, H.P., and Stoffler, D. Protofilaments and rings, two conformations of the tubulin family conserved from bacterial FtsZ to α/β and γ tubulin. 135:5.
Erle, D.J. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Erlebacher, A., and Derynck, R. Increased expression of TGF-β2 in osteoblasts results in an osteoporosis-like phenotype. 132:195.
Ernst, H. *See* Balsamo, Leung, Ernst, Zanin, Hoffman, and Lilien.
Ernst, J.D. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Ervasti, J.M. *See* Rybakova, Amann, and Ervasti.
Eshkind, L.G. *See* Chitaev, Leube, Troyanovsky, Eshkind, Franke, and Troyanovsky.

F

Fabbro, C. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
Fabra, A. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Fagotto, F., Funayama, N., Glück, U., and Gumbiner, B.M. Binding to cadherins antagonizes the signaling activity of β-catenin during axis formation in *Xenopus.* 132:1105.
Faissner, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Fan, Y. *See* Huang, Wu, Fan, and Adamson.
Fang, J., and Benbow, R.M. Nuclear proteins of quiescent *Xenopus laevis* cells inhibit DNA replication in intact and permeabilized nuclei. 133:955.
Fardeau, M. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Fares, H., Goetsch, L., and Pringle, J.R. Identification of a developmentally regulated septin and involvement of the septins in spore formation in *Saccharomyces cerevisiae.* 132:399.
Farese, R.V., Jr. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Farquhar, M.G. *See* Denker, McCaffery, Palade, Insel, and Farquhar.
Fässler, R. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Faulkner, J.A. *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
Faulstich, D., Auerbach, S., Orci, L., Ravazzola, M., Wegehingel, S., Lottspeich, F., Stenbeck, G., Harter, C., Wieland, F.T., and Tschochner, H. Architecture of coatomer: Molecular characterization of δ–COP and protein interactions within the complex. 135:53.
Faust, D.M. *See* Imaizumi-Scherrer, Faust, Bénichou, Hellio, and Weiss.
Fazeli, S., Wells, D.J., Hobbs, C., and Walsh, F.S. Altered secondary myogenesis in transgenic animals expressing the neural cell adhesion molecule under the control of a skeletal muscle α-actin promoter. 135:241.
Fechheimer, M. *See* Rivero, Furukawa, Noegel, and Fechheimer.
Fehon, R.G. *See* McCartney and Fehon.
Félix, M.-A. *See* Dogterom, Félix, Guet, and Leibler.
Ferguson, L.M. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Fernandez-Borja, M. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Ferns, M., Deiner, M., and Hall, Z. Agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine phosphorylation. 132:937.
Ferreira, J. *See* Visa, Izaurralde, Ferreira, Daneholt, and Mattaj.
Fialka, I., Schwarz, H., Reichmann, E., Oft, M., Busslinger, M., and Beug, H. The estrogen-dependent c-JunER protein causes a reversible loss of mammary epithelial cell polarity involving a destabilization of adherens junctions. 132:1115.
Fichtner, I. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Fiedler, K. *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Field, C.M., Al-Awar, O., Rosenblatt, J., Wong, M.L., Alberts, B., and Mitchison, T.J. A purified *Drosophila* septin complex forms filaments and exhibits GTPase activity. 133:605.
Filbin, M.T. *See* Wong and Filbin.
Fincham, V.J., Unlu, M., Brunton, V.G., Pitts, J.D., Wyke, J.A., and Frame, M.C. Translocation of Src kinase to the cell periphery is mediated by the actin cytoskeleton under the control of the Rho family of small G proteins. 135:1551.
Finger, E.B. *See* Alon, Fuhlbrigge, Finger, and Springer.
Fire, E. *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Fischer, E.R. *See* Scidmore, Fischer, and Hackstadt.
Flores, C. *See* Balda, Whitney, Flores, González, Cereijido, and Matter.
Florman, H.M. *See* Arnoult, Zeng, and Florman.
Flucher, B.E. *See* Powell, Petherbridge, and Flucher.
Fogliano, M. *See* Scherer, Lederkremer, Williams, Fogliano, Baldini, and Lodish.
Folmer, J. *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Forbes, D.J. *See* Macaulay and Forbes.
Formenton, A. *See* Gotta, Laroche, Formenton, Maillet, Scherthan, and Gasser.
Forster, J. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Foss, M. *See* Wein, Foss, Brady, and Cande.
Frame, M.C. *See* Fincham, Unlu, Brunton, Pitts, Wyke, and Frame.
Francis, R. *See* Iwasaki, McCarter, Francis, and Schedl.
Frank, J. *See* Verschoor, Srivastava, Grassucci, and Frank.
Frank, R. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Franke, W.W. *See* Chitaev, Leube, Troyanovsky, Eshkind, Franke, and Troyanovsky.
— *See* Keon, Schäfer, Kuhn, Grund, and Franke.
— *See* Mertens, Kuhn, and Franke.
— *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Frazier, W.A. *See* Gao, Lindberg, Dimitry, Brown, and Frazier.
French, L.E., Hahne, M., Viard, I., Radlgruber, G., Zanone, R., Becker, K., Müller, C., and Tschopp, J. Fas and Fas ligand in embryos and adult mice: Ligand expression in several immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover. 133:335.
Friedman, D.B., Sundberg, H.A., Huang, E.Y., and Davis, T.N. The 110-kD spindle pole body component of *Saccharomyces cerevisiae* is a phosphoprotein that is modified in a cell cycle–dependent manner. 132:903.
Friend, D.S., Ghildyal, N., Austen, K.F., Gurish, M.F., Matsumoto, R., and Stevens, R.L. Mast cells that reside at different locations in the jejunum of mice infected with *Trichinella spiralis* exhibit sequential changes in their granule ultrastructure and chymase phenotype. 135:279.
Frisch, S.M., Vuori, K., Kelaita, D., and Sicks, S. A role for Jun-N-terminal kinase in anoikis; suppression by bcl-2 and crmA. 135:1377.
—, **Vuori, K., Ruoslahti, E., and Chan-Hui, P.-Y.** Control of adhesion-dependent cell survival by focal adhesion kinase. 134:793.
Fuchs, E. *See* Allen, Yu, and Fuchs.
— *See* Dowling, Yu, and Fuchs.
Fuhlbrigge, R.C., Alon, R., Puri, K.D., Lowe, J.B., and Springer, T.A. Sialylated, fucosylated ligands for L-selectin expressed on leukocytes mediate tethering and rolling adhesions in physiologic flow conditions. 135:837.
— *See* Alon, Fuhlbrigge, Finger, and Springer.
Fujiki, Y. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Fujioka, H. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Fujisawa-Sehara, A. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Fukushima, T., Waddell, T.K., Grinstein, S., Goss, G.G., Orlowski, J., and Downey, G.P. Na^+/H^+ exchange activity during phagocytosis in human neutrophils: Role of Fcγ receptors and tyrosine kinases. 132:1037.
Fumagalli, L. *See* Lowell, Fumagalli, and Berton.
Funakoshi, T., Takeda, S., and Hirokawa, N. Active transport of photoactivated tubulin molecules in growing axons revealed by a new electron microscopic analysis. 133:1347.
— *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Funayama, N. *See* Fagotto, Funayama, Glück, and Gumbiner.
Furbee, C.S. *See* Rattner, Hendzel, Furbee, Muller, and Bazett-Jones.
Fürst. D.O. *See* Obermann, Gautel, Steiner, van der Ven, Weber, and Fürst.
Furukawa, R. *See* Rivero, Furukawa, Noegel, and Fechheimer.
Furuse, M. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Furuya, W. *See* Kim, Lingwood, Williams, Furuya, Manolson, and Grinstein.
Futey, L.M. *See* Kolman, Futey, and Egelhoff.
Futter, C.E., Pearse, A., Hewlett, L.J., and Hopkins, C.R. Multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and then fuse directly with lysosomes. 132:1011.

G

Gaglio, T., Saredi, A., Bingham, J.B., Hasbani, M.J., Gill, S.R., Schroer, T.A., and Compton, D.A. Opposing motor activities are required for the organization of the mammalian mitotic spindle pole. 135:399.
Gajhede, M. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Galou, M., Colucci-Guyon, E., Ensergueix, D., Ridet, J.-L., Giminez y Ribotta, M., Privat, A., Babinet, C., and Dupouey, P. Disrupted glial fibrillary acidic protein network in astrocytes from vimentin knockout mice. 133:853.
Galy, B. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Gammie, A.E. *See* Brizzio, Gammie, Nijbroek, Michaelis, and Rose.
Gansmuller, A. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.

Gao, A.-G., Lindberg, F.P., Dimitry, J.M., Brown, E.J., and Frazier, W.A. Thrombospondin modulates $\alpha_v\beta_3$ function through integrin-associated protein. 135:533.
Gard, D.L. *See* Kelley, Sellers, Gard, Bui, Adelstein, and Baines.
Gardner, L.C. *See* Rupp, O'Toole, Gardner, Mitchell, and Porter.
Gardner, R.L. *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
Garfinkel, S., Hu, X., Prudovsky, I.A., McMahon, G.A., Kapnik, E.M., McDowell, S.D., and Maciag, T. FGF-1–dependent proliferative and migratory responses are impaired in senescent human umbilical vein endothelial cells and correlate with the inability to signal tyrosine phosphorylation of fibroblast growth factor receptor-1 substrates. 134:783.
Garippa, R. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
Garoff, H. *See* Suomalainen, Hultenby, and Garoff.
Garrett, L.A. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Gasser, S.M. *See* Gotta, Laroche, Formenton, Maillet, Scherthan, and Gasser.
Gatti, M. *See* Williams, Gatti, and Goldberg.
Gautel, M. *See* Obermann, Gautel, Steiner, van der Ven, Weber, and Fürst.
Gebbink, M.F.B.G. *See* Zondag, Moolenaar, and Gebbink.
Geering, K., Beggah, A., Good, P., Girardet, S., Roy, S., Schaer, D., and Jaunin, P. Oligomerization and maturation of Na,K-ATPase: Functional interaction of the cytoplasmic NH_2 terminus of the β subunit with the α subunit. 133:1193.
Geiger, B. *See* Simcha, Geiger, Yehuda-Levenberg, Salomon, and Ben-Ze'ev.
Geisler, C. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Geissler, S. *See* Spang, Geissler, Grein, and Schiebel.
Geller, H.M. *See* Morris and Geller.
Gensac, M.-C. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Georgatos, S.D. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Gerace, L. *See* Hu, Guan, and Gerace.
— *See* Sweet and Gerace.
Geuze, H.J. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
— *See* Stoorvogel, Oorschot, and Geuze.
Ghildyal, N. *See* Friend, Ghildyal, Austen, Gurish, Matsumoto, and Stevens.
Ghosh, R.N. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
Ghosh, S. *See* Kielian, Klimjack, Ghosh, and Duffus.
Giancotti, F.G. *See* Mainiero, Pepe, Yeon, Ren, and Giancotti.
Gil, S.G. *See* Xia, Gil, and Carter.
Gilbert, D.M. *See* Lawlis, Keezer, Wu, and Gilbert.
Gill, S.R. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Gimbrone, M.A., Jr. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Giminez y Ribotta, M. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Ginsberg, M.H. *See* Huttenlocher, Ginsberg, and Horwitz.
Girardet, S. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Giuffrè, L. *See* Spertini, Cordey, Monai, Giuffrè, and Schapira.
Glabe, C.G. *See* Moy, Mendoza, Schulz, Swanson, Glabe, and Vacquier.
Glickman, J.N., Morton, P.A., Slot, J.W., Kornfeld, S., and Geuze, H.J. The biogenesis of the MHC class II compartment in human I-Cell disease B lymphoblasts. 132:769.
Glover, D.M., Ohkura, H., and Tavares, ŽA. Polo kinase: The choreographer of the mitotic stage? 135:1681.
Glück, U. *See* Fagotto, Funayama, Glück, and Gumbiner.
Godek, C.P. *See* Muresan, Godek, Reese, and Schnapp.
Godowski, P.J. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Goetsch, L. *See* Fares, Goetsch, and Pringle.
— *See* Sundberg, Goetsch, Byers, and Davis.
Gogos, J.A., Thompson, R., Lowry, W., Sloane, B.F., Weintraub, H., and Horwitz, M. Gene trapping in differentiating cell lines: Regulation of the lysosomal protease cathepsin B in skeletal myoblast growth and fusion. 134:837.
Goh, P.-Y. *See* Myers, Goh, Allen, Bucher, and Bogaert.
Golan, D.E. *See* Dustin, Ferguson, Chan, Springer, and Golan.
Goldberg, G.S. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Goldberg, M.L. *See* Williams, Gatti, and Goldberg.
Goldfarb, D.S. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Goldman, R.D., Khuon, S., Chou, Y.H., Opal, P., and Steinert, P.M. The function of intermediate filaments in cell shape and cytoskeletal integrity. 134:971.
Goldstein, L.S.B. *See* Matthies, McDonald, Goldstein, and Theurkauf.
Goldstein, M.A. *See* Schroeter, Bretaudiere, Sass, and Goldstein.
Gomer, R.H. *See* Van Haastert, Bishop, and Gomer.
González, S. *See* Balda, Whitney, Flores, González, Cereijido, and Matter.
Gonzalez-Garay, M.L., and Cabral, F. α-Tubulin limits its own synthesis: Evidence for a mechanism involving translational repression. 135:1525.
Good, P. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Goodman, J.M. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
— *See* Dyer, McNew, and Goodman.
— *See* Marshall, Dyer, Quick, and Goodman.
Goodness, T.P. *See* Albers, Perrone, Goodness, Jones, Green, and Davis.
Goodson, H.V., Anderson, B.L., Warrick, H.M., Pon, L.A., and Spudich, J.A. Synthetic lethality screen identifies a novel yeast myosin I gene (*MYO5*): Myosin I proteins are required for polarization of the actin cytoskeleton. 133:1277.
Goping, I.S. *See* McBride, Goping, and Shore.
Gorza, L. *See* Rossetto, Gorza, Schiavo, Scheller, and Montecucco.
Goss, G.G. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Gotta, M., Laroche, T., Formenton, A., Maillet, L., Scherthan, H., and Gasser, S.M. The clustering of telomeres and colocalization with Rap1, Sir3, and Sir4 proteins in wild-type *Saccharomyces cerevisiae*. 134:1349.
Götz, B., Scholze, A., Clement, A., Joester, A., Schütte, K., Wigger, F., Frank, R., Spiess, E., Ekblom, P., and Faissner, A. Tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons. 132:681.
Gould, G.W. *See* Martin, Tellam, Livingstone, Slot, Gould, and James.
Gould, S.J., Kalish, J.E., Morrell, J.C., Bjorkman, J., Urquhart, A.J., and Crane, D.I. Pex13p is an SH3 protein of the peroxisome membrane and a docking factor for the predominantly cytoplasmic PTS1 receptor. 135:85.
— *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
— *See* Dodt and Gould.
Goulding, D. *See* Maroto, Arredondo, Goulding, Marco, Bullard, and Cervera.
Goulet, M. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Goulielmos, G., Gounari, F., Remington, S., Müller, S., Häner, M., Aebi, U., and Georgatos, S.D. Filensin and phakinin form a novel type of beaded intermediate filaments and coassemble de novo in cultured cells. 132:643.
Gounari, F. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Gout, E. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Grandi, P. *See* Zabel, Doye, Tekotte, Wepf, Grandi, and Hurt.
Grassucci, R. *See* Verschoor, Srivastava, Grassucci, and Frank.
Green, K.J. *See* Bornslaeger, Corcoran, Stappenbeck, and Green.
Green, M.A. *See* Albers, Perrone, Goodness, Jones, Green, and Davis.
Green, M.R. *See* Zhang, Zapp, Yan, and Green.
Greengard, P. *See* Ryan, Li, Chin, Greengard, and Smith.
Grein, K. *See* Spang, Geissler, Grein, and Schiebel.
Grenningloh, G. *See* Di Paolo, Pellier, Catsicas, Antonsson, Catsicas, and Grenningloh.
Gresham, H.D. *See* Lindberg, Gresham, Reinhold, and Brown.
Griffin, J.W. *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Grimm, S., Bauer, M.K.A., Baeuerle, P.A., and Schulze-Osthoff, K. Bcl-2 down-regulates the activity of transcription factor NF-κB induced upon apoptosis. 134:13.
Grinstein, S. *See* Demaurex, Downey, Waddell, and Grinstein.
— *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
— *See* Kim, Lingwood, Williams, Furuya, Manolson, and Grinstein.
Grissom, P.M. *See* Vaisberg, Grissom, and McIntosh.
Grote, E., and Kelly, R.B. Endocytosis of VAMP is facilitated by a synaptic vesicle targeting signal. 132:537.
Gruenberg, J. *See* Aniento, Gu, Parton, and Gruenberg.
Grund, C. *See* Keon, Schäfer, Kuhn, Grund, and Franke.
— *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.

Gu, F. *See* Aniento, Gu, Parton, and Gruenberg.
— *See* Baurén, Jiang, Bernholm, Gu, and Wieslander.
Gu, Y. *See* Jin, Gu, and Morgan.
Gu, Z. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Guacci, V. *See* Yamamoto, Guacci, and Koshland.
Guan, T. *See* Hu, Guan, and Gerace.
Guet, C.C. *See* Dogterom, Félix, Guet, and Leibler.
Guild, G.M. *See* Tilney, Connelly, Smith, and Guild.
— *See* Tilney, Tilney, and Guild.
Guinet, F., Dvorak, J.A., Fujioka, H., Keister, D.B., Muratova, O., Kaslow, D.C., Aikawa, M., Vaidya, A.B., and Wellems, T.E. A developmental defect in *Plasmodium falciparum* male gametogenesis. 135:269.
Gullberg, D. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Gumbiner, B.M. *See* Brieher, Yap, and Gumbiner.
— *See* Fagotto, Funayama, Glück, and Gumbiner.
Günthert, A.R., Sträter, J., von Reyher, U., Henne, C., Joos, S., Koretz, K., Moldenhauer, G., Krammer, P.H., and Möller, P. Early detachment of colon carcinoma cells during CD95(APO-1/Fas)-mediated apoptosis I. De-adhesion from hyaluronate by shedding of CD44. 134:1089.
Günthert, U. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Gurish, M.F. *See* Friend, Ghildyal, Austen, Gurish, Matsumoto, and Stevens.
Gustavsson, M. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Gutkind, J.S. *See* Miyamoto, Teramoto, Gutkind, and Yamada.

H

Haass, N.K., Kartenbeck, J., and Leube, R.E. Pantophysin is a ubiquitously expressed synaptophysin homologue and defines constitutive transport vesicles. 134:731.
Hackstadt, T. *See* Scidmore, Fischer, and Hackstadt.
Haddad, A. *See* Chilcoat, Melia, Haddad, and Turkewitz.
Haggarty, B.S. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Haghighi, A. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Hagios, C., Koch, M., Spring, J., Chiquet, M., and Chiquet-Ehrismann, R. Tenascin-Y: A protein of novel domain structure is secreted by differentiated fibroblasts of muscle connective tissue. 134:1499.
Hahn, K.M. *See* Shelby, Hahn, and Sullivan.
Hahn, R.A. *See* Marchant, Hahn, Linsenmayer, and Birk.
Hahne, M. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Hajibagheri, M.A.N. *See* Ruhrberg, Hajibagheri, Simon, Dooley, and Watt.
Hajibagheri, N. *See* Dittié, Hajibagheri, and Tooze.
Hall, Z. *See* Ferns, Deiner, and Hall.
Hall, Z.W. *See* Wang, Hardy, and Hall.
Hamada, H. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Hamamoto, S. *See* Lupashin, Hamamoto, and Schekman.
Hammer, J.A., III *See* Jung, Wu, and Hammer.
Hanemaaijer, R. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Häner, M. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Hannan, L.A., and Edidin, M. Traffic, polarity, and detergent solubility of a glycosylphosphotidylinositol–anchored protein after LDL-deprivation of MDCK cells. 133:1265.
Hannon, K., Kudla, A.J., McAvoy, M.J., Clase, K.L., and Olwin, B.B. Differentially expressed fibroblast growth factors regulate skeletal muscle development through autocrine and paracrine mechanisms. 132:1151.
Hansen, P.A. *See* Wang, Hansen, Marshall, Holloszy, and Mueckler.
Hanzawa, N. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Harder, B.A. *See* Xu, Harder, Uman, and Craig.
Hardy, S.F. *See* Wang, Hardy, and Hall.
Härle, M. *See* Scherthan, Weich, Schwegler, Härle, Heyting, and Cremer.
Harley, C. *See* Henderson, Allsopp, Spector, Wang, and Harley.
Harris, S.L., and Waters, M.G. Localization of a yeast early Golgi mannosyltransferase, Och1p, involves retrograde transport. 132:985.
Hart, T.K. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Harter, C. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
Hartley, D. *See* Shpetner, Joly, Hartley, and Corvera.
Hartmann, E. *See* Voigt, Jungnickel, Hartmann, and Rapoport.
Hartwig, J.H. *See* Barkalow, Witke, Kwiatkowski, and Hartwig.
Hasbani, M.J. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Häusler, T., Stierhof, Y.-D., Wirtz, E., and Clayton, C. Import of a DHFR hybrid protein into glycosomes in vivo is not inhibited by the folate-analogue aminopterin. 132:311.
Hayashi, Y. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Hayes, S.A., and Dice, J.F. Roles of molecular chaperones in protein degradation. 132:255.
Hechi, J.H., Weiner, J.A., Post, S.R., and Chun, J. *Ventricular zone gene-1* (*vzg-1*) encodes a lysophosphatidic acid receptor expressed in neurogenic regions of the developing cerebral cortex. 135:1071.
Hefner-Gravink, A. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Hell, J. *See* Sakurai, Westenbroek, Rettig, Hell, and Catterall.
Hellio, R. *See* Imaizumi-Scherrer, Faust, Bénichou, Hellio, and Weiss.
Helms, J.B. *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Henderson, S., Allsopp, R., Spector, D., Wang, S.-S., and Harley, C. *In situ* analysis of changes in telomere size during replicative aging and cell transformation. 134:1.
— *See* Piperno, Mead, and Henderson.
Hendzel, M.J. *See* Rattner, Hendzel, Furbee, Muller, and Bazett-Jones.
Henis, Y.I. *See* Danieli, Pelletier, Henis, and White.
— *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Henley, J.R., and McNiven, M.A. Association of a dynamin-like protein with the Golgi apparatus in mammalian cells. 133:761.
Henne, C. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Henry, J.-P. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Henz, B.M. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Hernandez-Verdun, D. *See* Roussel, André, Comai, and Hernandez-Verdun.
Herrlich, P. *See* Sleeman, Rudy, Hofmann, Moll, Herrlich, and Ponta.
Herskowitz, I. *See* Sanders and Herskowitz.
— *See* Valtz and Herskowitz.
Heuser, J.E. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Hewlett, L.J. *See* Futter, Pearse, Hewlett, and Hopkins.
Heyberger, S. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Heyting, C. *See* Scherthan, Weich, Schwegler, Härle, Heyting, and Cremer.
Hill, K.L., Catlett, N.L., and Weisman, L.S. Actin and myosin function in directed vacuole movement during cell division in *Saccharomyces cerevisiae.* 135:1535.
Hillan, K.J. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Hiraki, Y. *See* Shukunami, Shigeno, Atsumi, Ishizeki, Suzuki, and Hiraki.
Hirao, M., Sato, N., Kondo, T., Yonemura, S., Monden, M., Sasaki, T., Takai, Y., Tsukita, S., and Tsukita, S. Regulation mechanism of ERM (ezrin/radixin/moesin) protein/plasma membrane association: Possible involvement of phosphatidylinositol turnover and Rho-dependent signaling pathway. 135:37.
Hirase, T. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Hirata, A. *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.
Hirayoshi, K. *See* Satoh, Hirayoshi, Yokota, Hosokawa, and Nagata.
Hirokawa, N., Funakoshi, T., Sato-Harada, R., and Kanai, Y. Selective stabilization of tau in axons and microtubule-associated protein 2C in cell bodies and dendrites contributes to polarized localization of cytoskeletal proteins in mature neurons. 132:667.
— *See* Funakoshi, Takeda, and Hirokawa.
Hisanaga, S.-I. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Hobbs, C. *See* Fazeli, Wells, Hobbs, and Walsh.
Hochmann, A. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Hocking, D.C., Smith, R.K., and McKeown-Longo, P.J. A novel role for the integrin-binding III-10 module in fibronectin matrix assembly. 133:431.

Hoffman, P.N. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
Hoffman, S. *See* Balsamo, Leung, Ernst, Zanin, Hoffman, and Lilien.
Hoffstrom, B.G. *See* Hungerford, Compton, Matter, Hoffstrom, and Otey.
Hofmann, M. *See* Sleeman, Rudy, Hofmann, Moll, Herrlich, and Ponta.
Hojo, M. *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Holash, J.A. *See* Soans, Holash, Pavlova, and Pasquale.
Holleran, E.A., Tokito, M.K., Karki, S., and Holzbaur, E.L.F. Centractin (ARP1) associates with spectrin revealing a potential mechanism to link dynactin to intracellular organelles. 135:1815.
Holloszy, J.O. *See* Wang, Hansen, Marshall, Holloszy, and Mueckler.
Holowka, D. *See* Pierini, Holowka, and Baird.
Holstein, S.E.H., Ungewickell, H., and Ungewickell, E. Mechanism of clathrin basket dissociation: Separate functions of protein domains of the DnaJ homologue auxilin. 135:925.
Holt, J.T. *See* Jotte and Holt.
Holzbaur, E.L.F. *See* Holleran, Tokito, Karki, and Holzbaur.
Homer, D. *See* Dubreuil, MacVicar, Dissanayake, Liu, Homer, and Hortsch.
Hong, E., Davidson, A.R., and Kaiser, C.A. A pathway for targeting soluble misfolded proteins to the yeast vacuole. 135:623.
Hong, K. *See* Lai, Hong, Kinnell, Chalfie, and Driscoll.
Honjo, Y. *See* Uchida, Honjo, Johnson, Wheelock, and Takeichi.
Hopkins, C.R. *See* Futter, Pearse, Hewlett, and Hopkins.
— *See* Odorizzi, Pearse, Domingo, Trowbridge, and Hopkins.
Hopmann, R., Cooper, J.A., and Miller, K.G. Actin organization, bristle morphology, and viability are affected by actin-capping protein mutations in *Drosophila*. 133:1293.
Hoppe, P.E., and Waterston, R.H. Hydrophobicity variations along the surface of the coiled-coil rod may mediate striated muscle myosin assembly in *Caenorhabditis elegans*. 135:371.
Hornsby, P.J. *See* Didenko and Hornsby.
Hortsch, M. *See* Dubreuil, MacVicar, Dissanayake, Liu, Homer, and Hortsch.
Horwitz, A.F. *See* Huttenlocher, Ginsberg, and Horwitz.
— *See* Sastry, Lakonishok, Thomas, Muschler, and Horwitz.
Horwitz, M. *See* Li, Coonrod, and Horwitz.
— *See* Gogos, Thompson, Lowry, Sloane, Weintraub, and Horwitz.
Hosokawa, N. *See* Satoh, Hirayoshi, Yokota, Hosokawa, and Nagata.
Hou, X. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Hough, C. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
— *See* Woods, Hough, Peel, Callaini, and Bryant.
Howard, D.E. *See* Blumenthal, Sarkar, Durell, Howard, and Morris.
Howard, E.W. *See* Alexander, Howard, Bissell, and Werb.
Howard, J.P. *See* Tan, Howard, and Payne.
Howard, T.A. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Hoxie, J.A. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Hsieh, S.-T. *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Hsieh, T.-S. *See* Zhang, Lee, Chen, Brown, and Hsieh.
Hu, T., Guan, T., and Gerace, L. Molecular and functional characterization of the p62 complex, an assembly of nuclear pore complex glycoproteins. 134:589.
Hu, X. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
Huang, E.Y. *See* Friedman, Sundberg, Huang, and Davis.
Huang, R.-P., Wu, J.-X., Fan, Y., and Adamson, E.D. UV activates growth factor receptors via reactive oxygen intermediates. 133:211.
Huang, S., and Spector, D.L. Intron-dependent recruitment of pre-mRNA splicing factors to sites of transcription. 133:719.
Huang, X.-Z., Wu, J.F., Cass, D., Erle, D.J., Corry, D., Young, S.G., Farese, R.V., Jr., and Sheppard, D. Inactivation of the integrin β6 subunit gene reveals a role of epithelial integrins in regulating inflammation in the lungs and skin. 133:921.
Huima, T. *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Hultenby, K. *See* Suomalainen, Hultenby, and Garoff.
Hungerford, J.E., Compton, M.T., Matter, M.L., Hoffstrom, B.G., and Otey, C.A. Inhibition of pp125^{FAK} in cultured fibroblasts results in apoptosis. 135:1383.
Hurt, E. *See* Irminger-Finger, Hurt, Roebuck, Collart, and Edelstein.
Hurt, E.C. *See* Zabel, Doye, Tekotte, Wepf, Grandi, and Hurt.
Huttenlocher, A., Ginsberg, M.H., and Horwitz, A.F. Modulation of cell migration by integrin-mediated cytoskeletal linkages and ligand-binding affinity. 134:1551.
Hynes, J.B. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Hynes, R.O. *See* Yang, Rando, Mohler, Rayburn, Blau, and Hynes.

I

Ide, C. *See* Igarashi, Kozaki, Terakawa, Kawano, Ide, and Komiya.
Igarashi, M., Kozaki, S., Terakawa, S., Kawano, S., Ide, C., and Komiya, Y. Growth cone collapse and inhibition of neurite growth by *botulinum* neurotoxin C1: A t-SNARE is involved in axonal growth. 134:205.
Ikebuchi, Y. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Ikeda, K. *See* Koenig and Ikeda.
Imafuku, I. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Imaizumi-Scherrer, T., Faust, D.M., Bénichou, J.-C., Hellio, R., and Weiss, M.C. Accumulation in fetal muscle and localization to the neuromuscular junction of cAMP-dependent protein kinase A regulatory and catalytic subunits RIα and Cα. 134:1241.
Imajoh-Ohmi, S. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Imhof, B.A. *See* Vainio, Dunon, Aïssi, Dangy, McNagny, and Imhof.
Inagaki, M. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
— *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Inagaki, N. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Ingber, D.E. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Inoue, T. *See* Muto, Kume, Inoue, Okano, and Mikoshiba.
Insel, P.A. *See* Denker, McCaffery, Palade, Insel, and Farquhar.
Irminger-Finger, I., Hurt, E., Roebuck, A., Collart, M.A., and Edelstein, S.J. *MHP1*, an essential gene in *Saccharomyces cerevisiae* required for microtubule function. 135:1323.
Ishizeki, K. *See* Shukunami, Shigeno, Atsumi, Ishizeki, Suzuki, and Hiraki.
Ishov, A.M., and Maul, G.G. The periphery of nuclear domain 10 (ND10) as site of DNA virus deposition. 134:815.
Islam, S., Carey, T.E., Wolf, G.T., Wheelock, M.J., and Johnson, K.R. Expression of N-cadherin by human squamous carcinoma cells induces a scattered fibroblastic phenotype with disrupted cell–cell adhesion. 135:1643.
Itoh, M. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Ivanov, I.E. *See* Simon, Ivanov, Adesnik, and Sabatini.
Ivanovska, I. *See* Biggins, Ivanovska, and Rose.
Iwasaki, K., McCarter, J., Francis, R., and Schedl, T. *emo-1*, a *Caenorhabditis elegans* Sec61p γ homologue, is required for oocyte development and ovulation. 134:699.
Izaurralde, E. *See* Visa, Izaurralde, Ferreira, Daneholt, and Mattaj.

J

Jacks, T. *See* Novitch, Mulligan, Jacks, and Lassar.
Jacobson, M.D., Weil, M., and Raff, M.C. Role of Ced-3/ICE-family proteases in staurosporine-induced programmed cell death. 133: 1041.
— *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
James, D.E. *See* Martin, Tellam, Livingstone, Slot, Gould, and James.
James, P. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
James, P.L., Stewart, C.E.H., and Rotwein, P. Insulin-like growth factor binding protein-5 modulates muscle differentiation through an insulin-like growth factor–dependent mechanism. 133:683.
Jansen, G. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Janssen, H. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Jass, G.K. *See* Opas, Szewczenko-Pawlikowski, Jass, Mesaeli, and Michalak.
Jaunin, P. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Jay, D.G. *See* Sydor, Su, Wang, Xu, and Jay.
Jeng, R.L. *See* Marschall, Jeng, Mulholland, and Stearns.
Jennings, P.B. *See* Schafer, Jennings, and Cooper.
Jiang, W.-Q. *See* Baurén, Jiang, Bernholm, Gu, and Wieslander.
Jin, P., Gu, Y., and Morgan, D.O. Role of inhibitory CDC2 phosphorylation in radiation-induced G2 arrest in human cells. 134:963.
Joester, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Johannes, F.-J. *See* Prestle, Pfizenmaier, Brenner, and Johannes.
Johansson, B.-M. *See* Bridge, Riedel, Johansson, and Pettersson.
Johansson, S. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.

Johnson, A.O., Ghosh, R.N., Dunn, K.W., Garippa, R., Park, J., Mayor, S., Maxfield, F.R., and McGraw, T.E. Transferrin receptor containing the SDYQRL motif of TGN38 causes a reorganization of the recycling compartment but is not targeted to the TGN. 135:1749.
Johnson, E.M., Jr. *See* Deshmukh, Vasilakos, Deckwerth, Lampe, Shivers, and Johnson.
Johnson, J. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Johnson, K.R. *See* Islam, Carey, Wolf, Wheelock, and Johnson.
— *See* Uchida, Honjo, Johnson, Wheelock, and Takeichi.
Johnson, M.T. *See* Araki, Johnson, and Swanson.
Johnson, R.G. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Joly, M. *See* Shpetner, Joly, Hartley, and Corvera.
Jones, K. *See* Sweet, Jones, and Allis.
Jones, K.T. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.
Jones, M.E. *See* Albers, Perrone, Goodness, Jones, Green, and Davis.
Joos, S. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Jordan, P., Mannervik, M., Tora, L., and Carmo-Fonseca, M. In vivo evidence that TATA-binding protein/SL1 colocalizes with UBF and RNA polymerase I when rRNA synthesis is either active or inactive. 133:225.
Joshi, H.C. *See* Miller and Joshi.
Jotte, R.M., and Holt, J.T. Myristylation of FBR v-fos dictates the differentiation pathways in malignant osteosarcoma. 135:457.
Jung, D. *See* Rafael, Cox, Corrado, Jung, Campbell, and Chamberlain.
Jung, G., Wu, X., and Hammer, J.A., III Dictyostelium mutants lacking multiple classic myosin I isoforms reveal combinations of shared and distinct functions. 133:305.
Jungnickel, B. *See* Voigt, Jungnickel, Hartmann, and Rapoport.
Jurgovsky, K. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Just, W.W. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.

K

Kahn, D.W. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Kaiser, C.A. *See* Hong, Davidson, and Kaiser.
Kaiser, S. *See* Blessing, Schirmacher, and Kaiser.
Kalish, J.E. *See* Gould, Kalish, Morrell, Bjorkman, Urquhart, and Crane.
Kalpin, R.F. *See* Debec, Kalpin, Daily, McCallum, Rothwell, and Sullivan.
Kamei, M. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Kammerer, R.A. *See* Brandenberger, Kammerer, Engel, and Chiquet.
Kanai, Y., Kanai-Azuma, M., Noce, T., Saido, T.C., Shiroishi, T., Hayashi, Y., and Yazaki, K. Identification of two Sox17 messenger RNA isoforms, with and without the high mobility group box region, and their differential expression in mouse spermatogenesis. 133:667.
— *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Kanai-Azuma, M. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Kanazawa, I. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Kanwar, Y.S. Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Kaplan, D.R. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Kapnik, E.M. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
Karki, S. *See* Holleran, Tokito, Karki, and Holzbaur.
Karpova, T.S. *See* Waddle, Karpova, Waterston, and Cooper.
Kartenbeck, J. *See* Haass, Kartenbeck, and Leube.
Kashgarian, M. *See* Devarajan, Stabach, Mann, Ardito, Kashgarian, and Morrow.
Kaslow, D.C. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Kasper, D., Dittmer, F., von Figura, K., and Pohlmann, R. Neither type of mannose 6-phosphate receptor is sufficient for targeting of lysosomal enzymes along intracellular routes. 134:615.
Kato, N. *See* Sakai, Saiganji, Yurimoto, Takabe, Saiki, and Kato.
Katyal, S. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Kawano, S. *See* Igarashi, Kozaki, Terakawa, Kawano, Ide, and Komiya.
Kedinger, M. *See* De Arcangelis, Neuville, Boukamel, Lefebvre, Kedinger, and Simon-Assmann.
Keen, J.H. *See* Santini and Keen.
Keene, D.R. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Keezer, S.M. *See* Lawlis, Keezer, Wu, and Gilbert.
Keister, D.B. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Kelaita, D. *See* Frisch, Vuori, Kelaita, and Sicks.
Keller, P. *See* Yoshimori, Keller, Roth, and Simons.
Kelley, C.A., Sellers, J.R., Gard, D.L., Bui, D., Adelstein, R.S., and Baines, I.C. Xenopus nonmuscle myosin heavy chain isoforms have different subcellular localizations and enzymatic activities. 134:675.
Kelly, R.B. *See* Cleves, Cooper, Barondes, and Kelly.
— *See* Grote and Kelly.
Keon, B.H., Schäfer, S., Kuhn, C., Grund, C., and Franke, W.W. Symplekin, a novel type of tight junction plaque protein. 134:1003.
Kern, M.A. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Khodjakov, A., Cole, R.W., Bajer, A.S., and Rieder, C.L. The force for poleward chromosome motion in *Haemanthus* cells acts along the length of the chromosome during metaphase but only at the kinetochore during anaphase. 132:1093.
—, **and Rieder, C.L.** Kinetochores moving away from their associated pole do not exert a significant pushing force on the chromosome. 135:315.
Khuon, S. *See* Goldman, Khuon, Chou, Opal, and Steinert.
Kidambi, U.S. *See* Kinose, Wang, Kidambi, Moncman, and Winkelmann.
Kiehart, D.P. *See* Cooper and Kiehart.
Kiel, J.A.K.W. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Kielian, M., Klimjack, M.R., Ghosh, S., and Duffus, W.A. Mechanisms of mutations inhibiting fusion and infection by Semliki forest virus. 134:863.
Kilmartin, J.V. *See* Donaldson and Kilmartin.
Kim, J.H., Lingwood, C.A., Williams, D.B., Furuya, W., Manolson, M.F., and Grinstein, S. Dynamic measurement of the pH of the Golgi complex in living cells using retrograde transport of the verotoxin receptor. 134:1387.
Kim, P.S., Kwon, O-Y., and Arvan, P. An endoplasmic reticulum storage disease causing congenital goiter with hypothyroidism. 133:517.
Kimura, T. *See* Tsumaki, Kimura, Matsui, Nakata, and Ochi.
Kincade, P.W. *See* Oritani and Kincade.
Kinnell, M. *See* Lai, Hong, Kinnell, Chalfie, and Driscoll.
Kinose, F., Wang, S.X., Kidambi, U.S., Moncman, C.L., and Winkelmann, D.A. Glycine 699 is pivotal for the motor activity of skeletal muscle myosin. 134:895.
Kinoshita, I. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Kirkpatrick, C. *See* Cox, Kirkpatrick, and Peifer.
Kishi, M. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Kishimoto, T. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Kiss, J.Z. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Kitsis, R.N. *See* Liu and Kitsis.
Klausner, R.D. *See* Radhakrishna, Klausner, and Donaldson.
Klebe, C. *See* Palacios, Weis, Klebe, Mattaj, and Dingwall.
Klein, A. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Klimjack, M.R. *See* Kielian, Klimjack, Ghosh, and Duffus.
Klionsky, D.J. *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Klymkowsky, M.W. *See* Bachant and Klymkowsky.
Knibbs, R.N., Craig, R.A., Natsuka, S., Chang, A., Cameron, M., Lowe, J.B., and Stoolman, L.M. The fucosyltransferase FucT-VII regulates E-selectin ligand synthesis in human T-cells. 133:911.
Koch, M. *See* Hagios, Koch, Spring, Chiquet, and Chiquet-Ehrismann.
Koenig, J.H., and Ikeda, K. Synaptic vesicles have two distinct recycling pathways. 135:797.
Koepp, D.M., Wong, D.H., Corbett, A.H., and Silver, P.A. Dynamic localization of the nuclear import receptor and its interactions with transport factors. 133:1163.
Kokojan, V. *See* Bender, Lo, Lee, Kokojan, Peterson, and Bender.
Kolde, G. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.

Kolman, M.F., Futey, L.M., and Egelhoff, T.T. *Dictyostelium* myosin heavy chain kinase A regulates myosin localization during growth and development. 132:101.
Komiya, Y. *See* Igarashi, Kozaki, Terakawa, Kawano, Ide, and Komiya.
Kondo, T. *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
Kono, T., Jones, K.T., Bos-Mikich, A., Whittingham, D.G., and Carroll, J. A cell cycle–associated change in Ca^{2+} releasing activity leads to the generation of Ca^{2+} transients in mouse embryos during the first mitotic division. 132:915.
Koolwijk, P., van Erck, M.G.M., de Vree, W.J.A., Vermeer, M.A., Weich, H.A., Hanemaaijer, R., and van Hinsbergh, V.W.M. Cooperative effect of TNFα, bFGF, and VEGF on the formation of tubular structures of human microvascular endothelial cells in a fibrin matrix. Role of urokinase activity. 132:1177.
Koretz, K. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Kornfeld, S. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
—*See* Rohrer, Schweizer, Russell, and Kornfeld.
— *See* Schweizer, Kornfeld, and Rohrer.
— *See* Traub, Bannykh, Rodel, Aridor, Balch, and Kornfeld.
Koshland, D. *See* Yamamoto, Guacci, and Koshland.
Koteliansky, V.E. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Kouranov, A. *See* Ma, Kouranov, LaSala, and Schnell.
Kozaki, S. *See* Igarashi, Kozaki, Terakawa, Kawano, Ide, and Komiya.
Krämer, H., and Phistry, M. Mutations in the *Drosophila hook* gene inhibit endocytosis of the boss transmembrane ligand into multivesicular bodies. 133:1205.
Krammer, P.H. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Krätzschmar, J. *See* Weskamp, Krätzschmar, Reid, and Blobel.
Kretz-Remy, C., Mehlen, P., Mirault, M.-E., and Arrigo, A.-P. Inhibition of IκB-α phosphorylation and degradation and subsequent NF-κB activation by glutathione peroxidase overexpression. 133:1083.
Kreuz, A.J., Simcox, A., and Maughan, D. Alterations in flight muscle ultrastructure and function in *Drosophila* tropomyosin mutants. 135:673.
Kronebusch, P.J. *See* Zhai, Kronebusch, Simon, and Borisy.
Ktistakis, N.T., Brown, H.A., Waters, M.G., Sternweis, P.C., and Roth, M.G. Evidence that phospholipase D mediates ADP ribosylation factor–dependent formation of Golgi coated vesicles. 134:295.
Kudla, A.J. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
Kuehn, M.J., Schekman, R., and Ljungdahl, P.O. Amino acid permeases require COPII components and the ER resident membrane protein Shr3p for packaging into transport vesicles in vitro. 135:585.
Kuhn, C. *See* Keon, Schäfer, Kuhn, Grund, and Franke.
—*See* Mertens, Kuhn, and Franke.
Kumar, A. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Kume, S. *See* Muto, Kume, Inoue, Okano, and Mikoshiba.
Kunau, W.-H. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Kunz, B. *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
— *See* Kunz, Ziegler, Kunz, and Sonderegger.
Kunz, S., Ziegler, U., Kunz, B., and Sonderegger, P. Intracellular signaling is changed after clustering of the neural cell adhesion molecules axonin-1 and NgCAM during neurite fasciculation. 135:253.
— *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
Kurz, A., Lampel, S., Nickolenko, J.E., Bradl, J., Benner, A., Zirbel, R.M., Cremer, T., and Lichter, P. Active and inactive genes localize preferentially in the periphery of chromosome territories. 135:1195.
Kwast, L. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Kwiatkowski, D.J. *See* Barkalow, Witke, Kwiatkowski, and Hartwig.
Kwon, O-Y. *See* Kim, Kwon, and Arvan.
Kypta, R.M., Su, H., and Reichardt, L.F. Association between a transmembrane protein tyrosine phosphatase and the cadherin–catenin complex. 134:1519.

L

LaBranche, C.C. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Lach, B. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Lachyankar, M.B. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Lacomis, L. *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
Lacroix, F.B. *See* Trielli, Andreassen, Lacroix, and Margolis.
Lai, C.-C., Hong, K., Kinnell, M., Chalfie, M., and Driscoll, M. Sequence and transmembrane topology of MEC-4, and ion channel subunit required for mechanotransduction in *Caenorhabditis elegans*. 133:1071.
Lakonishok, M. *See* Sastry, Lakonishok, Thomas, Muschler, and Horwitz.
Laloraya, S. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Lamb, R.A. *See* Bagai and Lamb.
— *See* Sakaguchi, Leser, and Lamb.
Lambert, S. *See* Davis, Lambert, and Bennett.
Lampe, P.A. *See* Deshmukh, Vasilakos, Deckwerth, Lampe, Shivers, and Johnson.
Lampe, P.D. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Lampel, S. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Lampugnani, M.G. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
Landmesser, L. *See* Rafuse and Landmesser.
Lane, H.A., and Nigg, E.A. Antibody microinjection reveals an essential role for human polo-like kinase 1 (Plk1) in the functional maturation of mitotic centrosomes. 135:1701.
Langa, C. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Langer, T. *See* Schmitt, Neupert, and Langer.
Lanzavecchia, S. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Laroche, T. *See* Gotta, Laroche, Formenton, Maillet, Scherthan, and Gasser.
Larochelle, D.A., Vithalani, K., and De Lozanne, A. A novel member of the *rho* family of small GTP-binding proteins is specifically required for cytokinesis. 133:1321.
LaSala, S.E. *See* Ma, Kouranov, LaSala, and Schnell.
Lassar, A.B. *See* Novitch, Mulligan, Jacks, and Lassar.
Lastres, P., Letamendía, A., Zhang, H., Rius, C., Almendro, N., Raab, U., López, L.A., Langa, C., Fabra, A., Letarte, M., and Bernabéu, C. Endoglin modulates cellular responses to TGF-β1. 133:1109.
Lawlis, S.J., Keezer, S.M., Wu, J.-R., and Gilbert, D.M. Chromosome architecture can dictate site-specific initiation of DNA replication in *Xenopus* egg extracts. 135:1207.
Lawrence, J.B. *See* Clemson, McNeil, Willard, and Lawrence.
Lazarovits, J., Naim, H.Y., Rodriguez, A.C., Wang, R.-H., Fire, E., Bird, C., Henis, Y.I., and Roth, M.G. Endocytosis of chimeric influenza virus hemagglutinin proteins that lack a cytoplasmic recognition feature for coated pits. 134:339.
Lazarow, P.B. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
— *See* Purdue and Lazarow.
— *See* Zhang and Lazarow.
Lazrak, A. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Lazure, C. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Lebakken, C.S., and Rapraeger, A.C. Syndecan-1 mediates cell spreading in transfected human lymphoblastoid (Raji) cells. 132:1209.
Lederkremer, G.Z. *See* Scherer, Lederkremer, Williams, Fogliano, Baldini, and Lodish.
Ledermann, B. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Lee, H. *See* Bender, Lo, Lee, Kokojan, Peterson, and Bender.
Lee, M.K. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
— *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
Lee, M.P. *See* Zhang, Lee, Chen, Brown, and Hsieh.
Lee, R.Y.N. *See* Starich, Lee, Panzarella, Avery, and Shaw.
Lee, W.M.F. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Lefebvre, O. *See* De Arcangelis, Neuville, Boukamel, Lefebvre, Kedinger, and Simon-Assmann.
Lefebvre, P.A. *See* Smith and Lefebvre.
Leibler, S. *See* Dogterom, Félix, Guet, and Leibler.

Leiper, J.M., Oatey, P.B., and Danpure, C.J. Inhibition of alanine: Glyoxylate aminotransferase 1 dimerization is a prerequisite for its peroxisome-to-mitochondrion mistargeting in primary hyperoxaluria type 1. 135:939.
Leitgeb, S. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Lemmon, V.P. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Leser, G.P. *See* Sakaguchi, Leser, and Lamb.
Letamendía, A. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Letarte, M. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Leube, R.E. *See* Chitaev, Leube, Troyanovsky, Eshkind, Franke, and Troyanovsky.
— *See* Haass, Kartenbeck, and Leube.
Leung, T. *See* Balsamo, Leung, Ernst, Zanin, Hoffman, and Lilien.
Leuschner, R. *See* Volkmer, Leuschner, Zacharias, and Rathjen.
Leventis, R. *See* Schroeder, Leventis, Shahinian, Walton, and Silvius.
Levin, S. *See* Liu, Tang, Edmonds, Murray, Levin, and Condeelis.
Lewis, J. *See* Brock, Midwinter, Lewis, and Martin.
Lewis, J.M., Cheresh, D.A., and Schwartz, M.A. Protein kinase C regulates αvβ5-dependent cytoskeletal associations and focal adhesion kinase phosphorylation. 134:1323.
Lewis, S.A., Tian, G., Vainberg, I.E., and Cowan, N.J. Chaperonin-mediated folding of actin and tubulin. 132:1.
Li, F.-Q., Coonrod, A., and Horwitz, M. Preferential MyoD homodimer formation demonstrated by a general method of dominant negative mutation employing fusion with a lysosomal protease. 135:1043.
Li, G. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Li, H., Liu, T.-F., Lazrak, A., Peracchia, C., Goldberg, G.S., Lampe, P.D., and Johnson, R.G. Properties and regulation of gap junctional hemichannels in the plasma membranes of cultured cells. 134:1019.
Li, L., Miano, J.M., Mercer, B., and Olson, E.N. Expression of the *SM22α* promoter in transgenic mice provides evidence for distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells. 132:849.
— *See* Ryan, Li, Chin, Greengard, and Smith.
Liang, T. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Liao, J., and Omary, M.B. 14-3-3 proteins associate with phosphorylated simple epithelial keratins during cell cycle progression and act as a solubility cofactor. 133:345.
Lichter, P. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Liesi, P., and Wright, J.M. Weaver granule neurons are rescued by calcium channel antagonists and antibodies against a neurite outgrowth domain of the B2 chain of laminin. 134:477.
Lilien, J. *See* Balsamo, Leung, Ernst, Zanin, Hoffman, and Lilien.
Lin, A. *See* Bastos, Lin, Enarson, and Burke.
Lin, L. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
Lindberg, F.P., Gresham, H.D., Reinhold, M.I., and Brown, E.J. Integrin-associated protein immunoglobulin domain is necessary for efficient vitronectin bead binding. 134:1313.
— *See* Gao, Lindberg, Dimitry, Brown, and Frazier.
Linder, M.E. *See* Timson Gauen, Linder, and Shaw.
Lingwood, C.A. *See* Kim, Lingwood, Williams, Furuya, Manolson, and Grinstein.
Linsenmayer, T.F. *See* Marchant, Hahn, Linsenmayer, and Birk.
Liu, C. *See* Dubreuil, MacVicar, Dissanayake, Liu, Homer, and Hortsch.
Liu, G., Tang, J., Edmonds, B.T., Murray, J., Levin, S., and Condeelis, J. F-actin sequesters elongation factor 1α from interaction with aminoacyl-tRNA in a pH-dependent reaction. 135:953.
— *See* Murray, Edmonds, Liu, and Condeelis.
Liu, T.-F. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Liu, Y., and Kitsis, R.N. Induction of DNA synthesis and apoptosis in cardiac myocytes by E1A oncoprotein. 133:325.
Liu, Z. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Livingstone, C. *See* Martin, Tellam, Livingstone, Slot, Gould, and James.
Ljungdahl, P.O. *See* Kuehn, Schekman, and Ljungdahl.
Lo, H.S. *See* Bender, Lo, Lee, Kokojan, Peterson, and Bender.
Lochmüller, H. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Locker, J. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Lodish, H.F. *See* Scherer, Lederkremer, Williams, Fogliano, Baldini, and Lodish.
Loechel, F. *See* Vachon, Loechel, Xu, Wewer, and Engvall.
Lohikangas, L. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Long, E.O. *See* Warmerdam, Long, and Roche.
Lopes-Cardozo, M. *See* van der Bijl, Lopes-Cardozo, and van Meer.
López, L.A. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Lottspeich, F. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
— *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Lounsbury, K.M. *See* Carey, Richards, Lounsbury, and Macara.
— *See* Richards, Lounsbury, Carey, and Macara.
Lowe, J.B. *See* Fuhlbrigge, Alon, Puri, Lowe, and Springer.
— *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Lowell, C.A., Fumagalli, L., and Berton, G. Deficiency of Src family kinases p59/61^{hck} and p58$^{c\text{-}fgr}$ results in defective adhesion-dependent neutrophil functions. 133:895.
Lowry, W. *See* Gogos, Thompson, Lowry, Sloane, Weintraub, and Horwitz.
Lu, D., Yang, H., and Raizada, M.K. Angiotensin II regulation of neuromodulation: Downstream signaling mechanism from activation of mitogen-activated protein kinase. 135:1609.
Ludérus, M.E.E., van Steensel, B., Chong, L., Sibon, O.C.M., Cremers, F.F.M., and de Lange, T. Structure, subnuclear distribution, and nuclear matrix association of the mammalian telomeric complex. 135:867.
Lue, R.A., Brandin, E., Chan, E.P., and Branton, D. Two independent domains of hDlg are sufficient for subcellular targeting: The PDZ1-2 conformational unit and an alternatively spliced domain. 135:1125.
Lund, K. *See* Campenot, Lund, and Senger.
Lunstrum, G.P. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Lupashin, V.V., Hamamoto, S., and Schekman, R.W. Biochemical requirements for the targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes. 132:277.
— *See* Sapperstein, Lupashin, Schmitt, and Waters.
Lupetti, P., Heuser, J.E., Manetti, R., Massari, P., Lanzavecchia, S., Bellon, P.L., Dallai, R., Rappuoli, R., and Telford, J.L. Oligomeric and subunit structure of the *Helicobacter pylori* vacuolating cytotoxin. 133:801.
Luskutoff, D.J. *See* Deng, Curriden, Wang, Rosenberg, and Loskutoff.
Lutz, Y. *See* Oulad-Abdelghani, Bouillet, Décimo, Gansmuller, Heyberger, Dollé, Bronner, Lutz, and Chambon.
Lyon, M.K. *See* Marr, Mastronarde, and Lyon.

M

Ma, Y., Kouranov, A., LaSala, S.E., and Schnell, D.J. Two components of the chloroplast protein import apparatus, IAP86 and IAP75, interact with the transit sequence during the recognition and translocation of precursor proteins at the outer envelope. 134:315.
Macara, I.G. *See* Carey, Richards, Lounsbury, and Macara.
— *See* Richards, Lounsbury, Carey, and Macara.
Macaulay, C., and Forbes, D.J. Assembly of the nuclear pore: Biochemically distinct steps revealed with NEM, GTPγS, and BAPTA. 132:5.
Mach, K.E. *See* Song, Mach, Chen, Reynolds, and Albright.
Maciag, T. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
Mackay, C.R. *See* Campbell, Qin, Bacon, Mackay, and Butcher.
Mac Nulty, E. *See* Nikolic, Mac Nulty, Mir, and Wiche.
MacVicar, G. *See* Dubreuil, MacVicar, Dissanayake, Liu, Homer, and Hortsch.
Madania, A. *See* Moreau, Madania, Martin, and Winsor.
Madara, J.L. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.
Magin, T.M. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Maguire, S. *See* Nehrbass, Rout, Maguire, Blobel, and Wozniak.
Maher, P.A. Nuclear translocation of fibroblast growth factor (FGF) receptors in response to FGF-2. 134:529.
Maier, A. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Maillet, L. *See* Gotta, Laroche, Formenton, Maillet, Scherthan, and Gasser.
Mainiero, F., Pepe, A., Yeon, M., Ren, Y., and Giancotti, F.G. The intracellular functions of $\alpha_6\beta_4$ integrin are regulated by EGF. 134:241.
Majeed, M. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Majoul, I.V., Bastiaens, P.I.H., and Söling, H.-D. Transport of an external Lys-Asp-Glu-Leu (KDEL) protein from the plasma membrane to the endoplasmic reticulum: Studies with cholera toxin in vero cells. 133:777.

Makki, A. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Malide, D. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Mammoto, A. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Manetti, R. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Mann, A.S. *See* Devarajan, Stabach, Mann, Ardito, Kashgarian, and Morrow.
Mannaerts, G.P. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Mannervik, M. *See* Jordan, Mannervik, Tora, and Carmo-Fonseca.
Manolson, M.F. *See* Kim, Lingwood, Williams, Furuya, Manolson, and Grinstein.
Marchant, J.K., Hahn, R.A., Linsenmayer, T.F., and Birk, D.E. Reduction of type V collagen using a dominant-negative strategy alters the regulation of fibrillogenesis and results in the loss of corneal-specific fibril morphology. 135:1415.
Marchetti, A. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Marcinkiewicz, M. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Marco, R. *See* Maroto, Arredondo, Goulding, Marco, Bullard, and Cervera.
Marfatia, S.M., Morais Cabral, J.H., Lin, L., Hough, C., Bryant, P.J., Stolz, L., and Chishti, A.H. Modular organization of the PDZ domains in the human discs-large protein suggests a mechanism for coupling PDZ domain-binding proteins to ATP and the membrane cytoskeleton. 135:753.
Margolis, R.L. *See* Trielli, Andreassen, Lacroix, and Margolis.
Markova, N.G. *See* Morasso, Markova, and Sargent.
Marks, M.S., Woodruff, L., Ohno, H., and Bonifacino, J.S. Protein targeting by tyrosine- and Di-leucine–based signals: Evidence for distinct saturable components. 135:341.
Maroto, M., Arredondo, J., Goulding, D., Marco, R., Bullard, B., and Cervera, M. *Drosophila* paramyosin/miniparamyosin gene products show a large diversity in quantity, localization, and isoform pattern: A possible role in muscle maturation and function. 134:81.
Marr, K.M., Mastronarde, D.N., and Lyon, M.K. Two-dimensional crystals of photosystem II: Biochemical characterization, cryo-electron microscopy and localization of the D1 and cytochrome b559 polypeptides. 132:823.
Marschall, L.G., Jeng, R.L., Mulholland, J., and Stearns, T. Analysis of Tub4p, a yeast γ-tubulin-like protein: Implications for microtubule-organizing center function. 134:443.
Marsh, L. *See* Elia and Marsh.
Marsh, M. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Marshall, B.A. *See* Wang, Hansen, Marshall, Holloszy, and Mueckler.
Marshall, P.A., Dyer, J.M., Quick, M.E., and Goodman, J.M. Redox-sensitive homodimerization of Pex11p: A proposed mechanism to regulate peroxisomal division. 135:123.
Marszalek, J.R. *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.
—, **Williamson, T.L., Lee, M.K., Xu, Z., Hoffman, P.N., Becher, M.W., Crawford, T.O., and Cleveland, D.W.** Neurofilament subunit NF-H modulates axonal diameter by selectively slowing neurofilament transport. 135:711.
Martin, P. *See* Brock, Midwinter, Lewis, and Martin.
Martin, R.P. *See* Moreau, Madania, Martin, and Winsor.
Martin, S., Tellam, J., Livingstone, C., Slot, J.W., Gould, G.W., and James, D.E. The glucose transporter (GLUT-4) and vesicle-associated membrane protein-2 (VAMP-2) are segregated from recycling endosomes in insulin-sensitive cells. 134:625.
Martin-Bermudo, M.D., and Brown, N.H. Intracellular signals direct integrin localization to sites of function in embryonic muscles. 134:217.
Martinou, I. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Martinou, J.-C. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Martín-Zanca, D. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Marty, F. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Marty-Mazars, D. *See* Aubert, Gout, Bligny, Marty-Mazars, Barrieu, Alabouvette, Marty, and Douce.
Marvaldi, J. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Marvulli, D. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
Maryon, E.B., Coronado, R., and Anderson, P. *unc-68* encodes a ryanodine receptor involved in regulating *C. elegans* body-wall muscle contraction. 134:885.
Massari, P. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Mastronarde, D.N. *See* Marr, Mastronarde, and Lyon.
Masumoto, N., Sasaki, T., Tahara, M., Mammoto, A., Ikebuchi, Y., Tasaka, K., Tokunaga, M., Takai, Y., and Miyake, A. Involvement of Rabphilin-3A in cortical granule exocytosis in mouse eggs. 135:1741.
Matlin, K.S. *See* Zinkl, Zuk, van der Bijl, van Meer, and Matlin.
Matsui, R. *See* Matsui, Matsui, Akada, and Toh-e.
Matsui, Y., Matsui, R., Akada, R., and Toh-e, A. Yeast *src* homology region 3 domain-binding proteins involved in bud formation. 133:865.
— *See* Tsumaki, Kimura, Matsui, Nakata, and Ochi.
Matsumoto, R. *See* Friend, Ghildyal, Austen, Gurish, Matsumoto, and Stevens.
Mattaj, I.W. *See* Palacios, Weis, Klebe, Mattaj, and Dingwall.
— *See* Visa, Izaurralde, Ferreira, Daneholt, and Mattaj.
Matter, K. *See* Balda, Whitney, Flores, González, Cereijido, and Matter.
Matter, M.L. *See* Hungerford, Compton, Matter, Hoffstrom, and Otey.
Matthies, H.J.G., McDonald, H.B., Goldstein, L.S.B., and Theurkauf, W.E. Anastral meiotic spindle morphogenesis: Role of the Non-Claret Disjunctional kinesin-like protein. 134:455.
Matunis, M.J., Coutavas, E., and Blobel, G. A novel ubiquitin-like modification modulates the partitioning of the Ran-GTPase–activating protein RanGAP1 between the cytosol and the nuclear pore complex. 135:1457.
Maughan, D. *See* Kreuz, Simcox, and Maughan.
Maul, G.G. *See* Ishov and Maul.
Maxfield, F.R. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
Mayer, H. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Mayhew, M.W. *See* Prekeris, Mayhew, Cooper, and Terrian.
Mayor, S. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
McAvoy, M.J. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
McBride, H.M., Goping, I.S., and Shore, G.C. The human mitochondrial import receptor, hTom20p, prevents a cryptic matrix targeting sequence from gaining access to the protein translocation machinery. 134:307.
McCaffery, J.M. *See* Denker, McCaffery, Palade, Insel, and Farquhar.
— *See* Rowe, Aridor, McCaffery, Plutner, Nuoffer, and Balch.
— *See* Wendland, McCaffery, Xiao, and Emr.
McCallum, P.D. *See* Debec, Kalpin, Daily, McCallum, Rothwell, and Sullivan.
McCarter, J. *See* Iwasaki, McCarter, Francis, and Schedl.
McCartney, B.M., and Fehon, R.G. Distinct cellular and subcellular patterns of expression imply distinct functions for the *Drosophila* homologues of moesin and the neurofibromatosis 2 tumor suppressor, merlin. 133:843.
McCracken, A.A., and Brodsky, J.L. Assembly of ER-associated protein degradation in vitro: Dependence on cytosol, calnexin, and ATP. 132:291.
McCrea, P.D. *See* Tao, Edwards, Tubb, Wang, Bryan, and McCrea.
McDonald, H.B. *See* Matthies, McDonald, Goldstein, and Theurkauf.
McDowell, S.D. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
McGraw, T.E. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
McGrew, L.L. *See* Torres, Yang-Snyder, Purcell, DeMarais, McGrew, and Moon.
McIntosh, J.R. *See* Vaisberg, Grissom, and McIntosh.
McKeown-Longo, P.J. *See* Hocking, Smith, and McKeown-Longo.
McKinnon, C.A. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
McMahon, G.A. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
McNagny, K.M. *See* Vainio, Dunon, Aïssi, Dangy, McNagny, and Imhof.
McNeil, J.A. *See* Clemson, McNeil, Willard, and Lawrence.
McNew, J.A. *See* Dyer, McNew, and Goodman.
McNiven, M.A. *See* Henley and McNiven.
Mead, K. *See* Piperno, Mead, and Henderson.
Mehlen, P. *See* Kretz-Remy, Mehlen, Mirault, and Arrigo.
Melia, S.M. *See* Chilcoat, Melia, Haddad, and Turkewitz.
Melton, D.W. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Mendoza, L.M. *See* Moy, Mendoza, Schulz, Swanson, Glabe, and Vacquier.
Menegazzi, R., Busetto, S., Dri, P., Cramer, R., and Patriarca, P. Chloride ion efflux regulates adherence, spreading, and respiratory burst of neutrophils stimulated by tumor necrosis factor-α (TNF) on biologic surfaces. 135:511.

Mennitt, P.A. *See* Cardone, Smith, Mennitt, Mochly-Rosen, Silver, and Mostov.
Mercer, B. *See* Li, Miano, Mercer, and Olson.
Merlie, J.P. *See* Tansey, Chu, and Merlie.
Mertens, C., Kuhn, C., and Franke, W.W. Plakophilins 2a and 2b: Constitutive proteins of dual location in the karyoplasm and the desmosomal plaque. 135:1009.
Mertens, G., van der Schueren, B., van den Berghe, H., and David, G. Heparan sulfate expression in polarized epithelial cells: The apical sorting of glypican (GPI-anchored proteoglycan) is inversely related to its heparan sulfate content. 132:487.
Mesaeli, N. *See* Opas, Szewczenko-Pawlikowski, Jass, Mesaeli, and Michalak.
Metzig, B. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Meyer, D.I. *See* Bush and Meyer.
Meyer, D.W. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Miano, J.M. *See* Li, Miano, Mercer, and Olson.
Miao, B. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Michaelis, S. *See* Brizzio, Gammie, Nijbroek, Michaelis, and Rose.
Michalak, M. *See* Opas, Szewczenko-Pawlikowski, Jass, Mesaeli, and Michalak.
Michalopoulos, G.K. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Midwinter, K. *See* Brock, Midwinter, Lewis, and Martin.
Mikoshiba, K. *See* Muto, Kume, Inoue, Okano, and Mikoshiba.
Miller, F.D. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Miller, K.E., and Joshi, H.C. Tubulin transport in neurons. 133:1355.
Miller, K.G. *See* Hopmann, Cooper, and Miller.
Miller, O.L., Jr. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Miller, S.G. *See* Chavez, Miller, and Moore.
Mills, P.L. *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
Milner, D.J., Weitzer, G., Tran, D., Bradley, A., and Capetanaki, Y. Disruption of muscle architecture and myocardial degeneration in mice lacking desmin. 134:1255.
Minamide, L.S. *See* Abe, Obinata, Minamide, and Bamburg.
Mineo, C. *See* Smart, Mineo, and Anderson.
Miner, J.H., and Sanes, J.R. Molecular and functional defects in kidneys of mice lacking collagen α3(IV): Implications for alport syndrome. 135:1403.
Mir, B. *See* Nikolic, Mac Nulty, Mir, and Wiche.
Mirault, M.-E. *See* Kretz-Remy, Mehlen, Mirault, and Arrigo.
Missotten, M. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Misteli, T. *See* Sönnichsen, Watson, Clausen, Misteli, and Warren.
Mitchell, B.F. *See* Rupp, O'Toole, Gardner, Mitchell, and Porter.
Mitchison, T.J. *See* Field, Al-Awar, Rosenblatt, Wong, Alberts, and Mitchison.
Mittal, B. *See* Ramanathan, Wilkemeyer, Mittal, Perides, and Charness.
Miyake, A. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Miyamoto, S., Teramoto, H., Gutkind, J.S., and Yamada, K.M. Integrins can collaborate with growth factors for phosphorylation of receptor tyrosine kinases and MAP kinase activation: Roles of integrin aggregation and occupancy of receptors. 135:1633.
Miyauchi, Y. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Mochly-Rosen, D. *See* Cardone, Smith, Mennitt, Mochly-Rosen, Silver, and Mostov.
Mohler, W.A. *See* Yang, Rando, Mohler, Rayburn, Blau, and Hynes.
Moldenhauer, G. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Moll, J. *See* Sleeman, Rudy, Hofmann, Moll, Herrlich, and Ponta.
Möller, P. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Monai, N. *See* Spertini, Cordey, Monai, Giuffrè, and Schapira.
Moncman, C.L. *See* Kinose, Wang, Kidambi, Moncman, and Winkelmann.
Monden, M. *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
Mondésert, G., and Reed, S.I. *BED1*, a gene encoding a galactosyltransferase homologue, is required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*. 132:137.
Monneron, A. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Montecucco, C. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Montell, D.J. *See* Murphy and Montell.
Montgomery, A.M.P., Becker, J.C., Siu, C.-H., Lemmon, V.P., Cheresh, D.A., Pancook, J.D., Zhao, X., and Reisfeld, R.A. Human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_V\beta_3$. 132:475.
Moolenaar, W.H. *See* Zondag, Moolenaar, and Gebbink.
Moon, R.T. *See* Torres, Yang-Snyder, Purcell, DeMarais, McGrew, and Moon.
Moore, H.-P.H. *See* Chavez, Miller, and Moore.
Moqrich, A. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Morais Cabral, J.H. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
Morasso, M.I., Markova, N.G., and Sargent, T.D. Regulation of epidermal differentiation by a *Distal-less* homeodomain gene. 135:1879.
Moreau, V., Madania, A., Martin, R.P., and Winsor, B. The *Saccharomyces cerevisiae* actin-related protein Arp2 is involved in the actin cytoskeleton. 134:117.
Morgan, D.O. *See* Jin, Gu, and Morgan.
Moritoh, C. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Morrell, J.C. *See* Gould, Kalish, Morrell, Bjorkman, Urquhart, and Crane.
Morris, E.J., and Geller, H.M. Induction of neuronal apoptosis by camptothecin, an inhibitor of DNA topoisomerase-I: Evidence for cell cycle–independent toxicity. 134:757.
Morris, S.J. *See* Blumenthal, Sarkar, Durell, Howard, and Morris.
Morrow, J.S. *See* Devarajan, Stabach, Mann, Ardito, Kashgarian, and Morrow.
Morton, P.A. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
Moser, B. *See* Weber, Alon, Moser, and Springer.
Moser, H.W. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Mostov, K.E. *See* Cardone, Smith, Mennitt, Mochly-Rosen, Silver, and Mostov.
Mougey, E.B. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Moy, G.W., Mendoza, L.M., Schulz, J.R., Swanson, W.J., Glabe, C.G., and Vacquier, V.D. The sea urchin sperm receptor for egg jelly is a modular protein with extensive homology to the human polycystic kidney disease protein, PKD1. 133:809.
Mueckler, M. *See* Wang, Hansen, Marshall, Holloszy, and Mueckler.
Mulholland, J. *See* Marschall, Jeng, Mulholland, and Stearns.
Müller, C. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Müller, H.-A.J., and Wieschaus, E. *armadillo*, *bazooka*, and *stardust* are critical for early stages in formation of the zonula adherens and maintenance of the polarized blastoderm epithelium in *Drosophila*. 134:149.
Muller, M.T. *See* Rattner, Hendzel, Furbee, Muller, and Bazett-Jones.
Müller, S. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Müller, W. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Mulligan, G.J. *See* Novitch, Mulligan, Jacks, and Lassar.
Mundigl, O. *See* Takei, Mundigl, Daniell, and De Camilli.
Murakoshi, H. *See* Scannevin, Murakoshi, Rhodes, and Trimmer.
Muratova, O. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Muresan, V., Godek, C.P., Reese, T.S., and Schnapp, B.J. Plus-end motors override minus-end motors during transport of squid axon vesicles on microtubules. 135:383.
Murphy, A.M., and Montell, D.J. Cell type–specific roles for Cdc42, Rac, and RhoL in *Drosophila* oogenesis. 133:617.
Murphy-Ullrich, J.E. *See* Chen, Murphy-Ullrich, and Wells.
Murray, A. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Murray, A.W. *See* Wells and Murray.
Murray, J. *See* Liu, Tang, Edmonds, Murray, Levin, and Condeelis
Murray, J.W., Edmonds, B.T., Liu, G., and Condeelis, J. Bundling of actin filaments by elongation factor 1 α inhibits polymerization at filament ends. 135:1309.
Müsch, A., Xu, H., Shields, D., and Rodriguez-Boulan, E. Transport of vesicular stomatitis virus G protein to the cell surface is signal mediated in polarized and nonpolarized cells. 133:543.
Muschler, J. *See* Sastry, Lakonishok, Thomas, Muschler, and Horwitz.
Muto, A., Kume, S., Inoue, T., Okano, H., and Mikoshiba, K. Calcium waves along the cleavage furrows in cleavage-stage *Xenopus* embryos and its inhibition by heparin. 135:181.
Myers, C.D., Goh, P.-Y., Allen, T.StC., Bucher, E.A., and Bogaert, T. Developmental genetic analysis of troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*. 132:1061.

N

Nabeshima, K. *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.

Nabeshima, Y. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Nagahama, M., Orci, L., Ravazzola, M., Amherdt, M., Lacomis, L., Tempst, P., Rothman, J.E., and Söllner, T.H. A v-SNARE implicated in intra-Golgi transport. 133:507.
Nagata, K. *See* Satoh, Hirayoshi, Yokota, Hosokawa, and Nagata.
Naim, H.Y. *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Nakano, A. *See* Sato, Sato, and Nakano.
Nakaseko, Y. *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.
Nakata, K. *See* Tsumaki, Kimura, Matsui, Nakata, and Ochi.
Nakayama, K. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Nakielny, S., and Dreyfuss, G. The hnRNP C proteins contain a nuclear retention sequence that can override nuclear export signals. 134:1365.
Näthke, I.S., Adams, C.L., Polakis, P., Sellin, J.H., and Nelson, W.J. The adenomatous polyposis coli tumor suppressor protein localizes to plasma membrane sites involved in active cell migration. 134:165.
Natsuka, S. *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Neefjes, J. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Nehrbass, U., Rout, M.P., Maguire, S., Blobel, G., and Wozniak, R.W. The yeast nucleoporin Nup188p interacts genetically and physically with the core structures of the nuclear pore complex. 133:1153.
Neisig, A. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Nelson, W.J. *See* Adams, Nelson, and Smith.
— *See* Angres, Barth, and Nelson.
— *See* Näthke, Adams, Polakis, Sellin, and Nelson.
Neupert, W. *See* Schmitt, Neupert, and Langer.
Neuville, P. *See* De Arcangelis, Neuville, Boukamel, Lefebvre, Kedinger, and Simon-Assmann.
Neveu, I., and Arenas, E. Neurotrophins promote the survival and development of neurons in the cerebellum of hypothyroid rats in vivo. 133:631.
Newman, L.S. *See* Simpson, Bright, West, Newman, Darnell, and Robinson.
Ng, D.T.W., Brown, J.D., and Walter, P. Signal sequences specify the targeting route to the endoplasmic reticulum membrane. 134:269.
—, **and Walter, P.** ER membrane protein complex required for nuclear fusion. 132:499.
Nickolenko, J.E. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Niclas, J., Allan, V.J., and Vale, R.D. Cell cycle regulation of dynein association with membranes modulates microtubule-based organelle transport. 133:585.
Niederreither, K. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Nigg, E.A. *See* Lane and Nigg.
Nijbroek, G. *See* Brizzio, Gammie, Nijbroek, Michaelis, and Rose.
Nikolic, B., Mac Nulty, E., Mir, B., and Wiche, G. Basic amino acid residue cluster within nuclear targeting sequence motif is essential for cytoplasmic plectin-vimentin network junctions. 134:1455.
Nishikawa, S.-I. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Nishiyama, T. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Noce, T. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Nocks, A. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Noegel, A.A. *See* Rivero, Furukawa, Noegel, and Fechheimer.
Nomura, M. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Norcott, J.P., Solari, R., and Cutler, D.F. Targeting of P-selectin to two regulated secretory organelles in PC12 cells. 134:1229.
Novitch, B.G., Mulligan, G.J., Jacks, T., and Lassar, A.B. Skeletal muscle cells lacking the retinoblastoma protein display defects in muscle gene expression and accumulate in S and G_2 phases of the cell cycle. 135:441.
Nuoffer, C. *See* Rowe, Aridor, McCaffery, Plutner, Nuoffer, and Balch.
Nürnberg, W. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Nusrat, A. *See* Parkos, Colgan, Liang, Nusrat, Bacarra, Carnes, and Madara.

O

Oatey, P.B. *See* Leiper, Oatey, and Danpure.
Obermann, W.M.J., Gautel, M., Steiner, F., van der Ven, P.F.M., Weber, K., and Fürst, D.O. The structure of the sarcomeric M band: Localization of defined domains of myomesin, M-protein, and the 250-kD carboxy-terminal region of titin by immunoelectron microscopy. 134:1441.
Obermeier, A. *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Obinata, T. *See* Abe, Obinata, Minamide, and Bamburg.
Ochi, T. *See* Tsumaki, Kimura, Matsui, Nakata, and Ochi.
Oda, M.N., Scott, S.V., Hefner-Gravink, A., Caffarelli, A.D., and Klionsky, D.J. Identification of a cytoplasm to vacuole targeting determinant in aminopeptidase I. 132:999.
Odorizzi, G., Pearse, A., Domingo, D., Trowbridge, I.S., and Hopkins, C.R. Apical and basolateral endosomes of MDCK cells are interconnected and contain a polarized sorting mechanism. 135:139.
Oft, M. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Ogawara, M. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Oh, Y.-K., and Swanson, J.A. Different fates of phagocytosed particles after delivery into macrophage lysosomes. 132:585.
Ohkura, H. *See* Glover, Ohkura, and Tavares.
Ohno, H. *See* Marks, Woodruff, Ohno, and Bonifacino.
Ohtsubo, M. *See* Zhu, Ohtsubo, Böhmer, Roberts, and Assoian.
Okano, H. *See* Muto, Kume, Inoue, Okano, and Mikoshiba.
Okazawa, H., Shimizu, J., Kamei, M., Imafuku, I., Hamada, H., and Kanazawa, I. Bcl-2 inhibits retinoic acid–induced apoptosis during the neural differentiation of embryonal stem cells. 132:955.
Okumura, E., Sekiai, T., Hisanaga, S.-I., Tachibana, K., and Kishimoto, T. Initial triggering of M-phase in starfish oocytes: A possible novel component of maturation-promoting factor besides cdc2 kinase. 132:125.
Olson, E.N. *See* Li, Miano, Mercer, and Olson.
Olwin, B.B. *See* Hannon, Kudla, McAvoy, Clase, and Olwin.
Omary, M.B. *See* Liao and Omary.
Oomen, L. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Oorschot, V. *See* Stoorvogel, Oorschot, and Geuze.
Opal, P. *See* Goldman, Khuon, Chou, Opal, and Steinert.
Opas, M., Szewczenko-Pawlikowski, M., Jass, G.K., Mesaeli, N., and Michalak, M. Calreticulin modulates cell adhesiveness via regulation of vinculin expression. 135:1913.
Orci, L. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
— *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
— *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
O'Reilly, M. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Oritani, K., and Kincade, P.W. Identification of stromal cell products that interact with pre-B cells. 134:771.
Orlowski, J. *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Orsulic, S., and Peifer, M. An in vivo structure-function study of armadillo, the β-catenin homologue, reveals both separate and overlapping regions of the protein required for cell adhesion and for wingless signaling. 134:1283.
Ortonne, J.-P. *See* Bertolotto, Bille, Ortonne, and Ballotti.
Osheim, Y.N., Mougey, E.B., Windle, J., Anderson, M., O'Reilly, Ostap, E.M., and Pollard, T.D. Biochemical kinetic characterization of the *Acanthamoeba* myosin-I ATPase. 132:1053.
—, **and Pollard, T.D.** Overlapping functions of myosin-I isoforms? 133:221.
Osumi, T. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Otey, C.A. *See* Hungerford, Compton, Matter, Hoffstrom, and Otey.
O'Toole, E. *See* Rupp, O'Toole, Gardner, Mitchell, and Porter.
Oulad-Abdelghani, M., Bouillet, P., Décimo, D., Gansmuller, A., Heyberger, S., Dollé, P., Bronner, S., Lutz, Y., and Chambon, P. Characterization of a premeiotic germ cell-specific cytoplasmic protein encoded by Stra8, a novel retinoic acid–responsive gene. 135:469.
Ouyang, P., and Sugrue, S.P. Characterization of pinin, a novel protein associated with the desmosome–intermediate filament complex. 135:1027.

P

Pagliusi, S. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.

Pahl, H.L., Sester, M., Burgert, H.-G., and Baeuerle, P.A. Activation of transcription factor NF-κB by the adenovirus E3/19K protein requires its ER retention. 132:511.
— *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Paillart, C., Boudier, J.-L., Boudier, J.-A., Rochat, H., Couraud, F., and Dargent, B. Activity-induced internalization and rapid degradation of sodium channels in cultured fetal neurons. 134:499.
Palacios, I., Weis, K., Klebe, C., Mattaj, I.W., and Dingwall, C. RAN/TC4 mutants identify a common requirement for snRNP and protein import into the nucleus. 133:485.
Palade, G.E. *See* Denker, McCaffery, Palade, Insel, and Farquhar.
Paladini, R.D., Takahashi, K., Bravo, N.S., and Coulombe, P.A. Onset of re-epithelialization after skin injury correlates with a reorganization of keratin filaments in wound edge keratinocytes: Defining a potential role for keratin 16. 132:381.
Pancook, J.D. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Pandiella, A. *See* Cabrera, Díaz-Rodríguez, Becker, Martín-Zanca, and Pandiella.
Panzarella, C. *See* Starich, Lee, Panzarella, Avery, and Shaw.
Park, J. *See* Johnson, Ghosh, Dunn, Garippa, Park, Mayor, Maxfield, and McGraw.
Parkos, C.A., Colgan, S.P., Liang, T., Nusrat, A., Bacarra, A.E., Carnes, D.K., and Madara, J.L. CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia. 132:437.
Parton, R.G. *See* Aniento, Gu, Parton, and Gruenberg.
— *See* Ullrich, Reinsch, Urbé, Zerial, and Parton.
Paschal, B.M. *See* Echeverri, Paschal, Vaughan, and Vallee.
Pasquale, E.B. *See* Soans, Holash, Pavlova, and Pasquale.
Patel, R. *See* Wilding, Wright, Patel, Ellis-Davies, and Whitaker.
Patriarca, P. *See* Menegazzi, Busetto, Dri, Cramer, and Patriarca.
Pavlova, Y. *See* Soans, Holash, Pavlova, and Pasquale.
Payne, G.S. *See* Tan, Howard, and Payne.
Pearse, A. *See* Futter, Pearse, Hewlett, and Hopkins.
— *See* Odorizzi, Pearse, Domingo, Trowbridge, and Hopkins.
Peel, D. *See* Woods, Hough, Peel, Callaini, and Bryant.
Peifer, M. *See* Cox, Kirkpatrick, and Peifer.
— *See* Orsulic and Peifer.
Pelchen-Matthews, A. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Pelletier, S.L. *See* Danieli, Pelletier, Henis, and White.
Pellier, V. *See* Di Paolo, Pellier, Catsicas, Antonsson, Catsicas, and Grenningloh.
Peloquin, J.G. *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
Pepe, A. *See* Mainiero, Pepe, Yeon, Ren, and Giancotti.
Peracchia, C. *See* Li, Liu, Lazrak, Peracchia, Goldberg, Lampe, and Johnson.
Peränen, J., Auvinen, P., Virta, H., Wepf, R., and Simons, K. Rab8 promotes polarized membrane transport through reorganization of actin and microtubules in fibroblasts. 135:153.
Perez, D. *See* Rao, Perez, and White.
Perides, G. *See* Ramanathan, Wilkemeyer, Mittal, Perides, and Charness.
Perrone, T.N. *See* Albers, Perrone, Goodness, Jones, Green, and Davis.
Perrotta, J.A. *See* Brown, Perrotta, and Turner.
Petersen, B.E. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Petersen, O.W. *See* Rønnov-Jessen and Petersen.
Peterson, J. *See* Bender, Lo, Lee, Kokojan, Peterson, and Bender.
Petherbridge, L. *See* Powell, Petherbridge, and Flucher.
Pettersson, U. *See* Bridge, Riedel, Johansson, and Pettersson.
Pfaff, M. *See* Wennerberg, Lohikangas, Gullberg, Pfaff, Johansson, and Fässler.
Pfizenmaier, K. *See* Prestle, Pfizenmaier, Brenner, and Johannes.
Phistry, M. *See* Krämer and Phistry.
Piccolo, S. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
Pierini, L., Holowka, D., and Baird, B. FcεRI-mediated association of 6-μm beads with RBL-2H3 mast cells results in exclusion of signaling proteins from the forming phagosome and abrogation of normal downstream signaling. 134:1427.
Pietrini, G. *See* Borgese, Aggujaro, Carrera, Pietrini, and Bassetti.
Pillus, L. *See* Stone and Pillus.
Piperno, G., Mead, K., and Henderson, S. Inner dynein arms but not outer dynein arms require the activity of kinesin homologue protein KHP1^{FLA10} to reach the distal part of flagella in *Chlamydomonas*. 133:371.
Pitts, J.D. *See* Fincham, Unlu, Brunton, Pitts, Wyke, and Frame.
Plonsky, I., and Zimmerberg, J. The initial fusion pore induced by baculovirus GP64 is large and forms quickly. 135:1831.
Plutner, H. *See* Rowe, Aridor, McCaffery, Plutner, Nuoffer, and Balch.
Poccia, D. *See* Collas, Courvalin, and Poccia.
Pohlmann, R. *See* Kasper, Dittmer, von Figura, and Pohlmann.
Polakis, P. *See* Näthke, Adams, Polakis, Sellin, and Nelson.
Pollard, T.D. *See* Ostap and Pollard.
Pon, L.A. *See* Goodson, Anderson, Warrick, Pon, and Spudich.
Ponta, H. *See* Sleeman, Rudy, Hofmann, Moll, Herrlich, and Ponta.
Porter, M.E. *See* Rupp, O'Toole, Gardner, Mitchell, and Porter.
Porter, R.M., Leitgeb, S., Melton, D.W., Swensson, O., Eady, R.A.J., and Magin, T.M. Gene targeting at the mouse cytokeratin 10 locus: Severe skin fragility and changes of cytokeratin expression in the epidermis. 132:925.
Portnoy, D.A. *See* Smith, Theriot, and Portnoy.
Post, S.R. *See* Hecht, Weiner, Post, and Chun.
Posthuma, G. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Powell, J.A., Petherbridge, L., and Flucher, B.E. Formation of triads without the dihydropyridine receptor α subunits in cell lines from dysgenic skeletal muscle. 134:375.
Pradel, L.-A. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Pralong, W.-F. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Prats, A.-C. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Prats, H. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Prekeris, R., Mayhew, M.W., Cooper, J.B., and Terrian, D.M. Identification and localization of an actin-binding motif that is unique to the epsilon isoform of protein kinase C and participates in the regulation of synaptic function. 132:77.
Prestle, J., Pfizenmaier, K., Brenner, J., and Johannes, F.-J. Protein kinase C μ is located at the Golgi compartment. 134:1401.
Pringle, J.R. *See* Fares, Goetsch, and Pringle.
Privat, A. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Prudovsky, I.A. *See* Garfinkel, Hu, Prudovsky, McMahon, Kapnik, McDowell, and Maciag.
Purcell, S.M. *See* Torres, Yang-Snyder, Purcell, DeMarais, McGrew, and Moon.
Purdue, P.E., and Lazarow, P.B. Targeting of human catalase to peroxisomes is dependent upon a novel COOH-terminal peroxisomal targeting sequence. 134:849.
Puri, K.D. *See* Fuhlbrigge, Alon, Puri, Lowe, and Springer.
Purich, D.L. *See* Zeile, Purich, and Southwick.

Q

Qin, S. *See* Campbell, Qin, Bacon, Mackay, and Butcher.
Quick, M.E. *See* Marshall, Dyer, Quick, and Goodman.

R

Raab, U. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Rachubinski, R.A. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Rader, C. *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
Radhakrishna, H., Klausner, R.D., and Donaldson, J.G. Aluminum fluoride stimulates surface protrusions in cells overexpressing the ARF6 GTPase. 134:935.
Radlgruber, G. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Rafael, J.A., Cox, G.A., Corrado, K., Jung, D., Campbell, K.P., and Chamberlain, J.S. Forced expression of dystrophin deletion constructs reveals structure–function correlations. 134:93.
— *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
Raff, K.D. *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
Raff, M.C. *See* Jacobson, Weil, and Raff.
— *See* Weil, Jacobson, Coles, Davies, Gardner, Raff, and Raff.
Rafuse, V.F., and Landmesser, L. Contractile activity regulates isoform expression and polysialylation of NCAM in cultured myotubes: Involvement of Ca^{2+} and protein kinase C. 132:969.
Raizada, M.K. *See* Lu, Yang, and Raizada.

Rajasekaran, A.K., Hojo, M., Huima, T., and Rodriguez-Boulan, E. Catenins and zonulus occulens-1 form a complex during early stages in the assembly of tight junctions. 132:451.
Ramaekers, F.C.S. *See* van der Loop, Schaart, Timmer, Ramaekers, and van Eys.
Ramanathan, R., Wilkemeyer, M.F., Mittal, B., Perides, G., and Charness, M.E. Alcohol inhibits cell–cell adhesion mediated by human L1. 133:381. Correction. 133:1139.
Ramos, J.W., and DeSimone, D.W. *Xenopus* embryonic cell adhesion to fibronectin: Position-specific activation of RGD/synergy site–dependent migratory behavior at gastrulation. 134:227.
Rando, T.A. *See* Yang, Rando, Mohler, Rayburn, Blau, and Hynes.
Rao, L., Perez, D., and White, E. Lamin proteolysis facilitates nuclear events during apoptosis. 135:1441.
Rao, L.V.M. *See* Sevinsky, Rao, and Ruf.
Rapoport, T.A. *See* Voigt, Jungnickel, Hartmann, and Rapoport.
Rappuoli, R. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Rapraeger, A.C. *See* Lebakken and Rapraeger.
Rashid, D.J. *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Rathjen, F.G. *See* Volkmer, Leuschner, Zacharias, and Rathjen.
Rattner, J.B., Hendzel, M.J., Furbee, C.S., Muller, M.T., and Bazett-Jones, D.P. Topoisomerase II α is associated with the mammalian centromere in a cell cycle– and species-specific manner and is required for proper centromere/kinetochore structure. 134:1097.
Ravazzola, M. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
— *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
— *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Ravid, S. *See* Dembinsky, Rubin, and Ravid.
Rayburn, H. *See* Yang, Rando, Mohler, Rayburn, Blau, and Hynes.
Redick, J. *See* Creutz, Snyder, Daigle, and Redick.
Reed, S.I. *See* Mondésert and Reed.
Reese, T.S. *See* Muresan, Godek, Reese, and Schnapp.
Reichardt, L. *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Reichardt, L.F. *See* Kypta, Su, and Reichardt.
Reichert, F. *See* Saada, Reichert, and Rotshenker.
Reichmann, E. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Reichsman, F., Smith, L., and Cumberledge, S. Glycosaminoglycans can modulate extracellular localization of the *wingless* protein and promote signal transduction. 135:819.
Reid, M.S. *See* Weskamp, Krätzschmar, Reid, and Blobel.
Reinhold, M.I. *See* Lindberg, Gresham, Reinhold, and Brown.
Reinsch, S. *See* Ullrich, Reinsch, Urbé, Zerial, and Parton.
Reisfeld, R.A. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Remington, S. *See* Goulielmos, Gounari, Remington, Müller, Häner, Aebi, and Georgatos.
Ren, Y. *See* Mainiero, Pepe, Yeon, Ren, and Giancotti.
Resnick, N. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Rettig, J. *See* Sakurai, Westenbroek, Rettig, Hell, and Catterall.
Reynolds, K., Zimmer, A.M., and Zimmer, A. Regulation of RARβ2 mRNA expression: Evidence for an inhibitory peptide encoded in the 5′-untranslated region. 134:827.
Reynolds, T. *See* Song, Mach, Chen, Reynolds, and Albright.
Rhodes, K.J. *See* Scannevin, Murakoshi, Rhodes, and Trimmer.
Richards, S.A., Lounsbury, K.M., Carey, K.L., and Macara, I.G. A nuclear export signal is essential for the cytosolic localization of the ran binding protein, RanBP1. 134:1157.
— *See* Carey, Richards, Lounsbury, and Macara.
Ridet, J.-L. *See* Galou, Colucci-Guyon, Ensergueix, Ridet, Giminez y Ribotta, Privat, Babinet, and Dupouey.
Ried, T. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Riedel, K.-U. *See* Bridge, Riedel, Johansson, and Pettersson.
Rieder, C.L. *See* Khodjakov, Cole, Bajer, and Rieder.
— *See* Khodjakov and Rieder.
Rijnboutt, S., Jansen, G., Posthuma, G., Hynes, J.B., Schornagel, J.H., and Strous, G.J. Endocytosis of GPI-linked membrane folate receptor-α. 132:35.
Riley, T. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Rioult, M.G. *See* Rodríguez, Samoff, Rioult, Chung, and Andrews.
Rius, C. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Rival, Y. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
Rivero, F., Furukawa, R., Noegel, A.A., and Fechheimer, M. *Dictyostelium discoideum* cells lacking the 34,000-dalton actin-binding protein can grow, locomote, and develop, but exhibit defects in regulation of cell structure and movement: A case of partial redundancy. 135:965.
Rizzo, M.G. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Robert, A. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Robert-Nicoud, M. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Roberts, J.M. *See* Zhu, Ohtsubo, Böhmer, Roberts, and Assoian.
Roberts, P. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Robertson, E.J. *See* Constam, Calfon, and Robertson.
Robinett, C.C., Straight, A., Li, G., Willhelm, C., Sudlow, G., Murray, A., and Belmont, A.S. In vivo localization of DNA sequences and visualization of large-scale chromatin organization using lac operator/repressor recognition. 135:1685.
Robinson, M.S. *See* Simpson, Bright, West, Newman, Darnell, and Robinson.
Rochat, H. *See* Paillart, Boudier, Boudier, Rochat, Couraud, and Dargent.
Roche, P.A. *See* Warmerdam, Long, and Roche.
Rodel, J.E. *See* Traub, Bannykh, Rodel, Aridor, Balch, and Kornfeld.
Rodgers, W., and Rose, J.K. Exclusion of CD45 inhibits activity of $p56^{lck}$ associated with glycolipid-enriched membrane domains. 135:1515.
Rodríguez, A., Samoff, E., Rioult, M.G., Chung, A., and Andrews, N.W. Host cell invasion by trypanosomes requires lysosomes and microtubule/kinesin–mediated transport. 134:349.
Rodriguez, A.C. *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Rodriguez-Boulan, E. *See* Müsch, Xu, Shields, and Rodriguez-Boulan.
— *See* Rajasekaran, Hojo, Huima, and Rodriguez-Boulan.
Roebuck, A. *See* Irminger-Finger, Hurt, Roebuck, Collart, and Edelstein.
Rohrer, J., Schweizer, A., Russell, D., and Kornfeld, S. The targeting of Lamp1 to lysosomes is dependent on the spacing of its cytoplasmic tail tyrosine sorting motif relative to the membrane. 132:565.
— *See* Schweizer, Kornfeld, and Rohrer.
Romano, J. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
Rønnov-Jessen, L., and Petersen, O.W. A function for filamentous α-smooth muscle actin: Retardation of motility in fibroblasts. 134:67.
Rosado, M. *See* Vincent, Rosado, and Davies.
Roscher, A. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Rose, J.K. *See* Rodgers and Rose.
Rose, M.D. *See* Biggins, Ivanovska, and Rose.
— *See* Brizzio, Gammie, Nijbroek, Michaelis, and Rose.
Rosenbach, T. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Rosenberg, M. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Rosenberg, S. *See* Deng, Curriden, Wang, Rosenberg, and Loskutoff.
Rosenblatt, J. *See* Field, Al-Awar, Rosenblatt, Wong, Alberts, and Mitchison.
Rosenthal, N. *See* Engert, Berglund, and Rosenthal.
Rosenzweig, A. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Ross, A.H., Daou, M.-C., McKinnon, C.A., Condon, P.J., Lachyankar, M.B., Stephens, R.M., Kaplan, D.R., and Wolf, D.E. The neurotrophin receptor, gp75, forms a complex with the receptor tyrosine kinase TrkA. 132:945.
Rossert, J. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Rossetto, O., Gorza, L., Schiavo, G., Schiavo, N., Scheller, R.H., and Montecucco, C. Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues. 132:167.
Roth, A.F., and Davis, N.G. Ubiquitination of the yeast α-factor receptor. 134:661.
Roth, M.G. *See* Ktistakis, Brown, Waters, Sternweis, and Roth.
— *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
— *See* Yoshimori, Keller, Roth, and Simons.
Rothman, J.E. *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
Rothwell, W.F. *See* Debec, Kalpin, Daily, McCallum, Rothwell, and Sullivan.
Rotshenker, S. *See* Saada, Reichert, and Rotshenker.
Rotwein, P. *See* James, Stewart, and Rotwein.

Rougon, G. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
— *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Roussel, P., André, C., Comai, L., and Hernandez-Verdun, D. The rDNA transcription machinery is assembled during mitosis in active NORs and absent in inactive NORs. 133:235.
Rousselle, P. *See* Champliaud, Lunstrum, Rousselle, Nishiyama, Keene, and Burgeson.
Rout, M.P. *See* Nehrbass, Rout, Maguire, Blobel, and Wozniak.
Rowe, T., Aridor, M., McCaffery, J.M., Plutner, H., Nuoffer, C., and Balch, W.E. COPII vesicles derived from mammalian endoplasmic reticulum microsomes recruit COPI. 135:895.
— *See* Bannykh, Rowe, and Balch.
Roy, B. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Roy, S. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Rubin, H. *See* Dembinsky, Rubin, and Ravid.
Rubin, K. *See* Sundberg and Rubin.
Ruco, L. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
Rudy, W. *See* Sleeman, Rudy, Hofmann, Moll, Herrlich, and Ponta.
Ruf, W. *See* Sevinsky, Rao, and Ruf.
Ruhrberg, C., Hajibagheri, M.A.N., Simon, M., Dooley, T.P., and Watt, F.M. Envoplakin, a novel precursor of the cornified envelope that has homology to desmoplakin. 134:715.
Ruiz, P., Brinkmann, V., Ledermann, B., Behrend, M., Grund, C., Thalhammer, C., Vogel, F., Birchmeier, C., Günthert, U., Franke, W.W., and Birchmeier, W. Targeted mutation of plakoglobin in mice reveals essential functions of desmosomes in the embryonic heart. 135:215.
Ruoslahti, E. *See* Frisch, Vuori, Ruoslahti, and Chan-Hui.
— *See* Wada, Kumar, Liu, Ruoslahti, Reichardt, Marvaldi, and Kanwar.
Rupp, G., O'Toole, E., Gardner, L.C., Mitchell, B.F., and Porter, M.E. The *sup-pf-2* mutations of *Chlamydomonas* alter the activity of the outer dynein arms by modification of the γ-dynein heavy chain. 135:1853.
Russell, D. *See* Rohrer, Schweizer, Russell, and Kornfeld.
Ryan, T.A., Li, L., Chin, L.-S., Greengard, P., and Smith, S.J. Synaptic vesicle recycling in synapsin I knock-out mice. 134:1219.
Rybakova, I.N., Amann, K.J., and Ervasti, J.M. A new model for the interaction of dystrophin with F-actin. 135:661.

S

Saada, A., Reichert, F., and Rotshenker, S. Granulocyte macrophage colony stimulating factor produced in lesioned peripheral nerves induces the up-regulation of cell surface expression of MAC-2 by macrophages and Schwann cells. 133:159.
Sabatini, D.D. *See* Simon, Ivanov, Adesnik, and Sabatini.
Sacchi, A. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Sachs, M., Weidner, K.M., Brinkmann, V., Walther, I., Obermeier, A., Ullrich, A., and Birchmeier, W. Motogenic and morphogenic activity of epithelial receptor tyrosine kinases. 133:1095.
Sadoul, R. *See* Allet, Hochmann, Martinou, Berger, Missotten, Antonsson, Sadoul, Martinou, and Bernasconi.
Sagot, I. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Saido, T.C. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Saiganji, A. *See* Sakai, Saiganji, Yurimoto, Takabe, Saiki, and Kato.
Saiki, H. *See* Sakai, Saiganji, Yurimoto, Takabe, Saiki, and Kato.
Saitoh, S., Takahashi, K., Nabeshima, K., Yamashita, Y., Nakaseko, Y., Hirata, A., and Yanagida, M. Aberrant mitosis in fission yeast mutants defective in fatty acid synthetase and acetyl CoA carboxylase. 134:949.
Saitou, M. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Sakaguchi, T., Leser, G.P., and Lamb, R.A. The ion channel activity of the influenza virus M_2 protein affects transport through the Golgi apparatus. 133:733.
Sakai, Y., Saiganji, A., Yurimoto, H., Takabe, K., Saiki, H., and Kato, N. The absence of Pmp47, a putative yeast peroxisomal transporter, causes a defect in transport and folding of a specific matrix enzyme. 134:37.
Sakakibara, A. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
Sakurai, T., Westenbroek, R.E., Rettig, J., Hell, J., and Catterall, W.A. Biochemical properties and subcellular distribution of the BI and rbA isoforms of α_{1A} subunits of brain calcium channel. 134:511.
Salin, P. *See* Castets, Bartoli, Barnier, Baillat, Salin, Moqrich, Bourgeois, Denizot, Rougon, Calothy, and Monneron.
Salomon, D. *See* Simcha, Geiger, Yehuda-Levenberg, Salomon, and Ben-Ze'ev.
Samoff, E. *See* Rodríguez, Samoff, Rioult, Chung, and Andrews.
Sanan, D.A. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Sanders, S.L., and Herskowitz, I. The bud4 protein of yeast, required for axial budding, is localized to the mother/bud neck in a cell cycle–dependent manner. 134:413.
Sanes, J.R. *See* Miner and Sanes.
Santini, F., and Keen, J.H. Endocytosis of activated receptors and clathrin-coated pit formation: Deciphering the chicken or egg relationship. 132:1025.
Santoro, S.A. *See* Xu, Zutter, Santoro, and Clark.
Sapperstein, S.K., Lupashin, V.V., Schmitt, H.D., and Waters, M.G. Assembly of the ER to Golgi SNARE complex requires Uso1p. 132:755.
Saredi, A. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Sargent, T.D. *See* Morasso, Markova, and Sargent.
Sarkar, D.P. *See* Blumenthal, Sarkar, Durell, Howard, and Morris.
Sasaki, T. *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
— *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Sass, R.L. *See* Schroeter, Bretaudiere, Sass, and Goldstein.
Sastry, S.K., Lakonishok, M., Thomas, D.A., Muschler, J., and Horwitz, A.F. Integrin α subunit ratios, cytoplasmic domains, and growth factor synergy regulate muscle proliferation and differentiation. 133:169.
Sato, K. *See* Sato, Sato, and Nakano.
Sato, M., Sato, K., and Nakano, A. Endoplasmic reticulum localization of Sec12p is achieved by two mechanisms: Rer1p-dependent retrieval that requires the transmembrane domain and Rer1p-independent retention that involves the cytoplasmic domain. 134:279.
Sato, N. *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
Satoh, M., Hirayoshi, K., Yokota, S., Hosokawa, N., and Nagata, K. Intracellular interaction of collagen-specific stress protein HSP47 with newly synthesized procollagen. 133:469.
Sato-Harada, R. *See* Hirokawa, Funakoshi, Sato-Harada, and Kanai.
Sauter, M.M., Pelchen-Matthews, A., Bron, R., Marsh, M., LaBranche, C.C., Vance, P.J., Romano, J., Haggarty, B.S., Hart, T.K., Lee, W.M.F., and Hoxie, J.A. An internalization signal in the simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on the cell surface. 132:795.
Scannevin, R.H., Murakoshi, H., Rhodes, K.J., and Trimmer, J.S. Identification of a cytoplasmic domain important in the polarized expression and clustering of the Kv2.1 K^+ channel. 135:1619.
Scardigli, R. *See* Soddu, Blandino, Scardigli, Coen, Marchetti, Rizzo, Bossi, Cimino, Crescenzi, and Sacchi.
Schaart, G. *See* van der Loop, Schaart, Timmer, Ramaekers, and van Eys.
Schadendorf, D., Kern, M.A., Artuc, M., Pahl, H.L., Rosenbach, T., Fichtner, I., Nürnberg, W., Stüting, S., Stebut, E.v., Worm, M., Makki, A., Jurgovsky, K., Kolde, G., and Henz, B.M. Treatment of melanoma cells with the synthetic retinoid CD437 induces apoptosis via activation of AP-1 in vitro, and causes growth inhibition in xenografts in vivo. 135:1889.
Schaer, D. *See* Geering, Beggah, Good, Girardet, Roy, Schaer, and Jaunin.
Schafer, D.A., Jennings, P.B., and Cooper, J.A. Dynamics of capping protein and actin assembly in vitro: Uncapping barbed ends by polyphosphoinositides. 135:169.
Schäfer, S. *See* Keon, Schäfer, Kuhn, Grund, and Franke.
Schapira, M. *See* Spertini, Cordey, Monai, Giuffrè, and Schapira.
Schedl, T. *See* Iwasaki, McCarter, Francis, and Schedl.
Schekman, R. *See* Kuehn, Schekman, and Ljungdahl.
Schekman, R.W. *See* Chuang and Schekman.
— *See* Lupashin, Hamamoto, and Schekman.
Scheller, R.H. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Scherer, P.E., Lederkremer, G.Z., Williams, S., Fogliano, M., Baldini, G., and Lodish, H.F. Cab45, a novel Ca^{2+}-binding protein localized to the Golgi lumen. 133:257.
Scherthan, H., Weich, S., Schwegler, H., Härle, M., Heyting, C., and Cremer, T. Centromere and telomere movements during early meiotic prophase of mouse and man are associated with the onset of chromosome pairing. 134:1109.
— *See* Gotta, Laroche, Formenton, Maillet, Scherthan, and Gasser.
Schiavo, G. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.

Schiavo, N. *See* Rossetto, Gorza, Schiavo, Schiavo, Scheller, and Montecucco.
Schiebel, E. *See* Spang, Geissler, Grein, and Schieble.
Schilke, B., Forster, J., Davis, J., James, P., Walter, W., Laloraya, S., Johnson, J., Miao, B., and Craig, E. The cold sensitivity of a mutant of *Saccharomyces cerevisiae* lacking a mitochondrial heat shock protein 70 is suppressed by loss of mitochondrial DNA. 134:603.
Schirmacher, P. *See* Blessing, Schirmacher, and Kaiser.
Schmitt, H.D. *See* Sapperstein, Lupashin, Schmitt, and Waters.
Schmitt, M., Neupert, W., and Langer, T. The molecular chaperone Hsp78 confers compartment-specific thermotolerance to mitochondria. 134:1375.
Schnapp, B.J. *See* Muresan, Godek, Reese, and Schnapp.
Schnell, D.J. *See* Ma, Kouranov, LaSala, and Schnell.
Scholey, J.M. Kinesin-II, a membrane traffic motor in axons, axonemes, and spindles. 133:1.
— *See* Wedaman, Meyer, Rashid, Cole, and Scholey.
Scholze, A. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Schornagel, J.H. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Schröck, E. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Schroeder, H., Leventis, R., Shahinian, S., Walton, P.A., and Silvius, J.R. Lipid-modified, cysteinyl-containing peptides of diverse structures are efficiently S-acylated at the plasma membrane of mammalian cells. 134:647.
Schroer, T.A. *See* Gaglio, Saredi, Bingham, Hasbani, Gill, Schroer, and Compton.
Schroeter, J.P., Bretaudiere, J.-P., Sass, R.L., and Goldstein, M.A. Three-dimensional structure of the Z band in a normal mammalian skeletal muscle. 133:571.
Schulz, J.R. *See* Moy, Mendoza, Schulz, Swanson, Glabe, and Vacquier.
Schulz, M.-F. *See* Wang, Pralong, Schulz, Rougon, Aubry, Pagliusi, Robert, and Kiss.
Schulze-Osthoff, K. *See* Grimm, Bauer, Baeuerle, and Schulze-Osthoff.
Schütte, K. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Schwall, R.H., Chang, L.Y., Godowski, P.J., Kahn, D.W., Hillan, K.J., Bauer, K.D., and Zioncheck, T.F. Heparin induces dimerization and confers proliferative activity onto the hepatocyte growth factor antagonists NK1 and NK2. 133:709.
Schwartz, M.A. *See* Lewis, Cheresh, and Schwartz.
Schwarz, H. *See* Fialka, Schwarz, Reichmann, Oft, Busslinger, and Beug.
Schwarzbauer, J.E. *See* Sechler, Takada, and Schwarzbauer.
Schwegler, H. *See* Scherthan, Weich, Schwegler, Härle, Heyting, and Cremer.
Schwei, M.J. *See* Yu, Schwei, and Baas.
Schweizer, A., Kornfeld, S., and Rohrer, J. Cysteine^{34} of the cytoplasmic tail of the cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting. 132:577.
— *See* Rohrer, Schweizer, Russell, and Kornfeld.
Scidmore, M.A., Fischer, E.R., and Hackstadt, T. Sphingolipids and glycoproteins are differentially trafficked to the *Chlamydia trachomatis* inclusion. 134:363.
Scott, R.E. *See* Wang, Xie, and Scott.
Scott, S.V., and Theg, S.M. A new chloroplast protein import intermediate reveals distinct translocation machineries in the two envelope membranes: Energetics and mechanistic implications. 132:63.
— *See* Oda, Scott, Hefner-Gravink, Caffarelli, and Klionsky.
Sechler, J.L., Takada, Y., and Schwarzbauer, J.E. Altered rate of fibronectin matrix assembly by deletion of the first type III repeats. 134:573.
Seidah, N.G. *See* De Bie, Marcinkiewicz, Malide, Lazure, Nakayama, Bendayan, and Seidah.
Sekiai, T. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Sekimata, M., Tsujimura, K., Tanaka, J., Takeuchi, Y., Inagaki, N., and Inagaki, M. Detection of protein kinase activity specifically activated at metaphase-anaphase transition. 132:635.
Sellers, J.R. *See* Kelley, Sellers, Gard, Bui, Adelstein, and Baines.
Sellin, J.H. *See* Näthke, Adams, Polakis, Sellin, and Nelson.
Senger, D.L. *See* Campenot, Lund, and Senger.
Sester, M. *See* Pahl, Sester, Burgert, and Baeuerle.
Seth, P. *See* Slack, Belliveau, Rosenberg, Atwal, Lochmüller, Aloyz, Haghighi, Lach, Seth, Cooper, and Miller.
Sevinsky, J.R., Rao, L.V.M., and Ruf, W. Ligand-induced protease receptor translocation into caveolae: A mechanism for regulating cell surface proteolysis of the tissue factor–dependent coagulation pathway. 133:293.
Shahinian, S. *See* Schroeder, Leventis, Shahinian, Walton, and Silvius.
Shaw, A.S. *See* Timson Gauen, Linder, and Shaw.
Shaw, J.E. *See* Starich, Lee, Panzarella, Avery, and Shaw.
Shelby, R.D., Hahn, K.M., and Sullivan, K.F. Dynamic elastic behavior of α-satellite DNA domains visualized in situ in living human cells. 135:545.
Sheppard, D. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Shields, D. *See* Austin and Shields.
— *See* Müsch, Xu, Shields, and Rodriguez-Boulan.
Shigeno, C. *See* Shukunami, Shigeno, Atsumi, Ishizeki, Suzuki, and Hiraki.
Shimizu, J. *See* Okazawa, Shimizu, Kamei, Imafuku, Hamada, and Kanazawa.
Shiroishi, T. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Shivers, B.D. *See* Deshmukh, Vasilakos, Deckwerth, Lampe, Shivers, and Johnson.
Shore, G.C. *See* McBride, Goping, and Shore.
Shpetner, H., Joly, M., Hartley, D., and Corvera, S. Potential sites of PI-3 kinase function in the endocytic pathway revealed by the PI-3 kinase inhibitor, wortmannin. 132:595.
Shukunami, C., Shigeno, C., Atsumi, T., Ishizeki, K., Suzuki, F., and Hiraki, Y. Chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 in vitro: Differentiation-dependent gene expression of parathyroid hormone (PTH)/PTH-related peptide receptor. 133:457.
Shulga, N., Roberts, P., Gu, Z., Spitz, L., Tabb, M.M., Nomura, M., and Goldfarb, D.S. In vivo nuclear transport kinetics in *Saccharomyces cerevisiae*: A role for heat shock protein 70 during targeting and translocation. 135:329.
Sibon, O.C.M. *See* Ludérus, van Steensel, Chong, Sibon, Cremers, and de Lange.
Sicks, S. *See* Frisch, Vuori, Kelaita, and Sicks.
Siegrist, S.E. *See* Stronach, Siegrist, and Beckerle.
Silver, P.A. *See* Koepp, Wong, Corbett, and Silver.
Silver, R.B. *See* Cardone, Smith, Mennitt, Mochly-Rosen, Silver, and Mostov.
Silvius, J.R. *See* Schroeder, Leventis, Shahinian, Walton, and Silvius.
Simcha, I., Geiger, B., Yehuda-Levenberg, S., Salomon, D., and Ben-Ze'ev, A. Suppression of tumorigenicity by plakoglobin: An augmenting effect of N-cadherin. 133:199.
Simcox, A. *See* Kreuz, Simcox, and Maughan.
Simon, J.-P., Ivanov, I.E., Adesnik, M., and Sabatini, D.D. The production of post-Golgi vesicles requires a protein kinase C–like molecule, but not its phosphorylating activity. 135:355.
Simon, M. *See* Ruhrberg, Hajibagheri, Simon, Dooley, and Watt.
Simon, P.M. *See* Zhai, Kronebusch, Simon, and Borisy.
Simon-Assmann, P. *See* De Arcangelis, Neuville, Boukamel, Lefebvre, Kedinger, and Simon-Assmann.
Simons, K. *See* Eaton, Wepf, and Simons.
— *See* Peränen, Auvinen, Virta, Wepf, and Simons.
— *See* Yoshimori, Keller, Roth, and Simons.
Simpson, F., Bright, N.A., West, M.A., Newman, L.S., Darnell, R.B., and Robinson, M.S. A novel adaptor-related protein complex. 133:749.
Siu, C.-H. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Slack, R.S., Belliveau, D.J., Rosenberg, M., Atwal, J., Lochmüller, H., Aloyz, R., Haghighi, A., Lach, B., Seth, P., Cooper, E., and Miller, F.D. Adenovirus-mediated gene transfer of the tumor suppressor, p53, induces apoptosis in postmitotic neurons. 135:1085.
Sleeman, J., Rudy, W., Hofmann, M., Moll, J., Herrlich, P., and Ponta, H. Regulated clustering of variant CD44 proteins increases their hyaluronate binding capacity. 135:1139.
Sloane, B.F. *See* Gogos, Thompson, Lowry, Sloane, Weintraub, and Horwitz.
Slot, J.W. *See* Glickman, Morton, Slot, Kornfeld, and Geuze.
— *See* Martin, Tellam, Livingstone, Slot, Gould, and James.
Small, J.V. *See* Anderson, Wang, and Small.
Smart, E.J., Mineo, C., and Anderson, R.G.W. Clustered folate receptors deliver 5-methyltetrahydrofolate to cytoplasm of MA104 cells. 134:1169.
Smit, J.J.M. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
Smith, B.L. *See* Cardone, Smith, Mennitt, Mochly-Rosen, Silver, and Mostov.
Smith, C. *See* Bou-Gharios, Garrett, Rossert, Niederreither, Eberspaecher, Smith, Black, and de Crombrugghe.
Smith, E.F., and Lefebvre, P.A. *PF*16 encodes a protein with armadillo repeats and localizes to a single microtubule of the central apparatus in *Chlamydomonas* flagella. 132:359.
Smith, G.A., Theriot, J.A., and Portnoy, D.A. The tandem repeat domain in the *Listeria monocytogenes* ActA protein controls the rate of actin-based motility, the percentage of moving bacteria, and the localization of vasodilator-stimulated phosphoprotein and profilin. 135:647.
Smith, L. *See* Reichsman, Smith, and Cumberledge.

Smith, R.K. *See* Hocking, Smith, and McKeown-Longo.
Smith, S. *See* Tilney, Connelly, Smith, and Guild.
Smith, S.J. *See* Adams, Nelson, and Smith.
— *See* Ryan, Li, Chin, Greengard, and Smith.
Snyder, S.L. *See* Creutz, Snyder, Daigle, and Redick.
Soans, C., Holash, J.A., Pavlova, Y., and Pasquale, E.B. Developmental expression and distinctive tyrosine phosphorylation of the Eph-related receptor tyrosine kinase Cek9. 135:781.
Soddu, S., Blandino, G., Scardigli, R., Coen, S., Marchetti, A., Rizzo, M.G., Bossi, G., Cimino, L., Crescenzi, M., and Sacchi, A. Interference with p53 protein inhibits hematopoietic and muscle differentiation. 134:193.
Sohn, K., Orci, L., Ravazzola, M., Amherdt, M., Bremser, M., Lottspeich, F., Fiedler, K., Helms, J.B., and Wieland, F.T. A major transmembrane protein of Golgi-derived COPI-coated vesicles involved in coatomer binding. 135:1239.
Solari, R. *See* Norcott, Solari, and Cutler.
Söling, H.-D. *See* Majoul, Bastiaens, and Söling.
Söllner, T.H. *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
Sollner-Webb, B. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Sonderegger, P. *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
— *See* Kunz, Ziegler, Kunz, and Sonderegger.
Song, K., Mach, K.E., Chen, C.-Y., Reynolds, T., and Albright, C.F. A novel suppressor of *ras1* in fission yeast, byr4, is a dosage-dependent inhibitor of cytokinesis. 133:1307.
Sönnichsen, B., Watson, R., Clausen, H., Misteli, T., and Warren, G. Sorting by COP I–coated vesicles under interphase and mitotic conditions. 134:1411.
Southwick, F.S. *See* Zeile, Purich, and Southwick.
Spang, A., Geissler, S., Grein, K., and Schiebel, E. γ-Tubulin–like Tub4p of *Saccharomyces cerevisiae* is associated with the spindle pole body substructures that organize microtubules and is required for mitotic spindle formation. 134:429.
Spector, D. *See* Henderson, Allsopp, Spector, Wang, and Harley.
Spector, D.L. *See* Huang and Spector.
Speicher, M.R. *See* Eils, Dietzel, Bertin, Schröck, Speicher, Ried, Robert-Nicoud, Cremer, and Cremer.
Spertini, O., Cordey, A.-S., Monai, N., Giuffrè, L., and Schapira, M. P-Selectin glycoprotein ligand 1 is a ligand for L-selectin on neutrophils, monocytes, and $CD34^+$ hematopoietic progenitor cells. 135:523.
Spiess, E. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Spitz, L. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Spring, F. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Spring, J. *See* Hagios, Koch, Spring, Chiquet, and Chiquet-Ehrismann.
Springer, T.A. *See* Alon, Fuhlbrigge, Finger, and Springer.
— *See* Clark, Alon, and Springer.
— *See* Dustin, Ferguson, Chan, Springer, and Golan.
— *See* Fuhlbrigge, Alon, Puri, Lowe, and Springer.
— *See* Weber, Alon, Moser, and Springer.
Spudich, J.A. *See* Goodson, Anderson, Warrick, Pon, and Spudich.
Srivastava, S. *See* Verschoor, Srivastava, Grassucci, and Frank.
Stabach, P.R. *See* Devarajan, Stabach, Mann, Ardito, Kashgarian, and Morrow.
Stamenkovic, I. *See* Bartolazzi, Nocks, Aruffo, Spring, and Stamenkovic.
Stappenbeck, T.S. *See* Bornslaeger, Corcoran, Stappenbeck, and Green.
Starich, T.A., Lee, R.Y.N., Panzarella, C., Avery, L., and Shaw, J.E. *eat-5* and *unc-7* represent a multigene family in *Caenorhabditis elegans* involved in cell–cell coupling. 134:537.
Stearns, T. *See* Marschall, Jeng, Mulholland, and Stearns.
Stebut, E.v. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Steiner, F. *See* Obermann, Gautel, Steiner, van der Ven, Weber, and Fürst.
Steinert, P.M. *See* Goldman, Khuon, Chou, Opal, and Steinert.
Steinfeld, R., Van Den Berghe, H., and David, G. Stimulation of fibroblast growth factor receptor-1 occupancy and signaling by cell surface–associated syndecans and glypican. 133:405.
Stenbeck, G. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
Stendahl, O. *See* Zimmerli, Majeed, Gustavsson, Stendahl, Sanan, and Ernst.
Stephens, R.M. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Sternweis, P.C. *See* Ktistakis, Brown, Waters, Sternweis, and Roth.
Stevens, R.L. *See* Friend, Ghildyal, Austen, Gurish, Matsumoto, and Stevens.
Stevens, T.H. *See* Cooper and Stevens.
Stewart, C.E.H. *See* James, Stewart, and Rotwein.
Stierhof, Y.-D. *See* Häusler, Stierhof, Wirtz, and Clayton.
Stoffel, W. *See* Wahle and Stoffel.
Stoffler, D. *See* Erickson and Stoffler.
Stolz, L. *See* Marfatia, Morais Cabral, Lin, Hough, Bryant, Stolz, and Chishti.
Stone, E.M., and Pillus, L. Activation of an MAP kinase cascade leads to Sir3p hyperphosphorylation and strengthens transcriptional silencing. 135:571.
Stoolman, L.M. *See* Knibbs, Craig, Natsuka, Chang, Cameron, Lowe, and Stoolman.
Stoorvogel, W., Oorschot, V., and Geuze, H.J. A novel class of clathrin-coated vesicles budding from endosomes. 132:21.
Straight, A. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Sträter, J. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Strom, S.C. *See* Block, Locker, Bowen, Petersen, Katyal, Strom, Riley, Howard, and Michalopoulos.
Stronach, B.E., Siegrist, S.E., and Beckerle, M.C. Two muscle-specific LIM proteins in *Drosophila*. 134:1179.
Strous, G.J. *See* Rijnboutt, Jansen, Posthuma, Hynes, Schornagel, and Strous.
Stüting, S. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Su, A.L. *See* Sydor, Su, Wang, Xu, and Jay.
Su, H. *See* Kypta, Su, and Reichardt.
Subramani, S. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Sudlow, G. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Sugrue, S.P. *See* Ouyang and Sugrue.
Sullivan, K.F. *See* Shelby, Hahn, and Sullivan.
Sullivan, W. *See* Debec, Kalpin, Daily, McCallum, Rothwell, and Sullivan.
Sundberg, C., and Rubin, K. Stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors. 132:741.
Sundberg, H.A., Goetsch, L., Byers, B., and Davis, T.N. Role of calmodulin and Spc110p interaction in the proper assembly of spindle pole body components. 133:111.
— *See* Friedman, Sundberg, Huang, and Davis.
Suomalainen, M., Hultenby, K., and Garoff, H. Targeting of moloney murine leukemia virus gag precursor to the site of virus budding. 135:1841.
Sutoh, K. *See* Aizawa, Sutoh, and Yahara.
Suzuki, F. *See* Shukunami, Shigeno, Atsumi, Ishizeki, Suzuki, and Hiraki.
Svitkina, T.M., Verkhovsky, A.B., and Borisy, G.G. Plectin sidearms mediate interaction of intermediate filaments with microtubules and other components of the cytoskeleton. 135:991.
Swanson, J.A. *See* Araki, Johnson, and Swanson.
— *See* Oh and Swanson.
Swanson, W.J. *See* Moy, Mendoza, Schulz, Swanson, Glabe, and Vacquier.
Swartzman, E.E., Viswanathan, M.N., and Thorner, J. The *PAL1* gene product is a peroxisomal ATP-binding cassette transporter in the yeast *Saccharomyces cerevisiae*. 132:549.
Sweet, D.J., and Gerace, L. A GTPase distinct from ran is involved in nuclear protein import. 133:971.
Sweet, M.T., Jones, K., and Allis, C.D. Phosphorylation of linker histone is associated with transcriptional activation in a normally silent nucleus. 135:1219.
Swensson, O. *See* Porter, Leitgeb, Melton, Swensson, Eady, and Magin.
Sydor, A.M., Su, A.L., Wang, F.-S., Xu, A., and Jay, D.G. Talin and vinculin play distinct roles in filopodial motility in the neuronal growth cone. 134:1197.
Sympson, C.J., Talhouk, R.S., Alexander, C.M., Chin, J.R., Clift, S.M., Bissell, M.J., and Werb, Z. Targeted expression of stromelysin-1 in mammary gland provides evidence for a role of proteinases in branching morphogenesis and the requirement for an intact basement membrane for tissue-specific gene expression: Correction. 132:753.
Szewczenko-Pawlikowski, M. *See* Opas, Szewczenko-Pawlikowski, Jass, Mesaeli, and Michalak.

T

Tabak, H.F. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.

— *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Tabb, M.M. *See* Shulga, Roberts, Gu, Spitz, Tabb, Nomura, and Goldfarb.
Tachibana, K. *See* Okumura, Sekiai, Hisanaga, Tachibana, and Kishimoto.
Tahara, M. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Takabe, K. *See* Sakai, Saiganji, Yurimoto, Takabe, Saiki, and Kato.
Takada, Y. *See* Sechler, Takada, and Schwarzbauer.
Takahashi, K. *See* Paladini, Takahashi, Bravo, and Coulombe.
— *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.
Takai, Y., Ogawara, M., Tomono, Y., Moritoh, C., Imajoh-Ohmi, S., Tsutsumi, O., Taketani, Y., and Inagaki, M. Mitosis-specific phosphorylation of vimentin by protein kinase C coupled with reorganization of intracellular membranes. 133:141.
— *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
— *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Takeda, S. *See* Funakoshi, Takeda, and Hirokawa.
Takei, K., Mundigl, O., Daniell, L., and De Camilli, P. The synaptic vesicle cycle: A single vesicle building step involving clathrin and dynamin. 133:1237.
Takeichi, M. *See* Uchida, Honjo, Johnson, Wheelock, and Takeichi.
Taketani, Y. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Takeuchi, Y. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Taki, T. *See* Yoshida, Fujisawa-Sehara, Taki, Arai, and Nabeshima.
Talhouk, R.S. *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Tan, P.K., Howard, J.P., and Payne, G.S. The sequence NPFXD defines a new class of endocytosis signal in *Saccharomyces cerevisiae.* 135:1789.
Tanaka, J. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Tang, J. *See* Liu, Tang, Edmonds, Murray, Levin, and Condeelis.
Tang, X. *See* Benian, Tinley, Tang, and Borodovsky.
Tanimura, A., and Turner, R.J. Calcium release in HSY cells conforms to a steady-state mechanism involving regulation of the inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal [Ca^{2+}]. 132:607.
Tansey, M.G., Chu, G.C., and Merlie, J.P. ARIA/HRG regulates AChR ϵ subunit gene expression at the neuromuscular synapse via activation of phosphatidylinositol 3-kinase and Ras/MAPK pathway. 134:465.
Tao, Y.S., Edwards, R.A., Tubb, B., Wang, S., Bryan, J., and McCrea, P.D. β-catenin associates with the actin-bundling protein fascin in a noncadherin complex. 134:1271.
Tarone, G. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Tasaka, K. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Tavares, Ž.A. *See* Glover, Ohkura, and Tavares.
Tekotte, H. *See* Zabel, Doye, Tekotte, Wepf, Grandi, and Hurt.
Telford, J.L. *See* Lupetti, Heuser, Manetti, Massari, Lanzavecchia, Bellon, Dallai, Rappuoli, and Telford.
Tellam, J. *See* Martin, Tellam, Livingstone, Slot, Gould, and James.
Tempst, P. *See* Nagahama, Orci, Ravazzola, Amherdt, Lacomis, Tempst, Rothman, and Söllner.
Terakawa, S. *See* Igarashi, Kozaki, Terakawa, Kawano, Ide, and Komiya.
Teramoto, H. *See* Miyamoto, Teramoto, Gutkind, and Yamada.
Terrian, D.M. *See* Prekeris, Mayhew, Cooper, and Terrian.
Thalhammer, C. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Theg, S.M. *See* Scott and Theg.
Theriot, J.A. *See* Smith, Theriot, and Portnoy.
Theurkauf, W.E. *See* Matthies, McDonald, Goldstein, and Theurkauf.
Thomas, D.A. *See* Sastry, Lakonishok, Thomas, Muschler, and Horwitz.
Thompson, R. *See* Gogos, Thompson, Lowry, Sloane, Weintraub, and Horwitz.
Thorner, J. *See* Swartzman, Viswanathan, and Thorner.
Tian, G. *See* Lewis, Tian, Vainberg, and Cowan.
Ticchioni, M. *See* Deckert, Ticchioni, and Bernard.
Tilney, L.G., Connelly, P., Smith, S., and Guild, G.M. F-Actin bundles in *Drosophila* bristles are assembled from modules composed of short filaments. 135:1291.
—, **Tilney, M.S., and Guild, G.M.** Formation of actin filament bundles in the ring canals of developing *Drosophila* follicles. 133:61.
Tilney, M.S. *See* Tilney, Tilney, and Guild.
Timmer, E.D.J. *See* van der Loop, Schaart, Timmer, Ramaekers, and van Eys.
Timson Gauen, L.K., Linder, M.E., and Shaw, A.S. Multiple features of the p59^{fyn} src homology 4 domain define a motif for immune-receptor tyrosine-based activation motif (ITAM) binding and for plasma membrane localization. 133:1007.
Tinley, T.L. *See* Benian, Tinley, Tang, and Borodovsky.
Toh-e, A. *See* Matsui, Matsui, Akada, and Toh-e.
Tokito, M.K. *See* Holleran, Tokito, Karki, and Holzbaur.
Tokunaga, M. *See* Masumoto, Sasaki, Tahara, Mammoto, Ikebuchi, Tasaka, Tokunaga, Takai, and Miyake.
Tomatis, D. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Tomé, F. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Tomono, Y. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Tooze, S.A. *See* Dittié, Hajibagheri, and Tooze.
Tora, L. *See* Jordan, Mannervik, Tora, and Carmo-Fonseca.
Torres, M.A., Yang-Snyder, J.A., Purcell, S.M., DeMarais, A.A., McGrew, L.L., and Moon, R.T. Activities of the *Wnt*-1 class of secreted signaling factors are antagonized by the *Wnt*-5A class and by a dominant negative cadherin in early *Xenopus* development. 133:1123.
Touriol, C. *See* Vagner, Touriol, Galy, Audigier, Gensac, Amalric, Bayard, Prats, and Prats.
Tran, D. *See* Milner, Weitzer, Tran, Bradley, and Capetanaki.
Traub, L.M., Bannykh, S.I., Rodel, J.E., Aridor, M., Balch, W.E., and Kornfeld, S. AP-2–containing clathrin coats assemble on mature lysosomes. 135:1801.
Tremblay, J.P. *See* Vilquin, Kinoshita, Roy, Goulet, Engvall, Tomé, Fardeau, and Tremblay.
Trielli, M.O., Andreassen, P.R., Lacroix, F.B., and Margolis, R.L. Differential taxol-dependent arrest of transformed and nontransformed cells in the G1 phase of the cell cycle, and specific-related mortality of transformed cells. 135:689.
Trimmer, J.S. *See* Scannevin, Murakoshi, Rhodes, and Trimmer.
Trowbridge, I.S. *See* Odorizzi, Pearse, Domingo, Trowbridge, and Hopkins.
Troyanovsky, R.B. *See* Chitaev, Leube, Troyanovsky, Eshkind, Franke, and Troyanovsky.
Troyanovsky, S.M. *See* Chitaev, Leube, Troyanovsky, Eshkind, Franke, and Troyanovsky.
Tschochner, H. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
Tschopp, J. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Tsujimura, K. *See* Sekimata, Tsujimura, Tanaka, Takeuchi, Inagaki, and Inagaki.
Tsukamoto, T. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Tsukita, S. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
— *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
Tsumaki, N., Kimura, T., Matsui, Y., Nakata, K., and Ochi, T. Separable *cis*-regulatory elements that contribute to tissue- and site-specific α2(XI) collagen gene expression in the embryonic mouse cartilage. 134:1573.
Tsutsumi, O. *See* Takai, Ogawara, Tomono, Moritoh, Imajoh-Ohmi, Tsutsumi, Taketani, and Inagaki.
Tubb, B. *See* Tao, Edwards, Tubb, Wang, Bryan, and McCrea.
Tulp, A. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Turkewitz, A.P. *See* Chilcoat, Melia, Haddad, and Turkewitz.
Turner, C.E. *See* Brown, Perrotta, and Turner.
Turner, R.J. *See* Tanimura and Turner.

U

Uchida, N., Honjo, Y., Johnson, K.R., Wheelock, M.J., and Takeichi, M. The catenin/cadherin adhesion system is localized in synaptic junctions bordering transmitter release zones. 135:767.
Ullrich, A. *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Ullrich, O., Reinsch, S., Urbé, S., Zerial, M., and Parton, R.G. Rab11 regulates recycling through the pericentriolar recycling endosome. 135:913.
Uman, P. *See* Xu, Harder, Uman, and Craig.
Ungewickell, E. *See* Holstein, Ungewickell, and Ungewickell.

Ungewickell, H. *See* Holstein, Ungewickell, and Ungewickell.
Unlu, M. *See* Fincham, Unlu, Brunton, Pitts, Wyke, and Frame.
Urbé, S. *See* Ullrich, Reinsch, Urbé, Zerial, and Parton.
Urquhart, A.J. *See* Gould, Kalish, Morrell, Bjorkman, Urquhart, and Crane.

V

Vachon, P.H., Loechel, F., Xu, H., Wewer, U.M., and Engvall, E. Merosin and laminin in myogenesis; specific requirement for merosin in myotube stability and survival. 134:1483.
Vacquier, V.D. *See* Moy, Mendoza, Schulz, Swanson, Glabe, and Vacquier.
Vagner, S., Touriol, C., Galy, B., Audigier, S., Gensac, M.-C., Amalric, F., Bayard, F., Prats, H., and Prats, A.-C. Translation of CUG- but not AUG-initiated forms of human fibroblast growth factor 2 is activated in transformed and stressed cells. 135:1391.
Vaidya, A.B. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Vainberg, I.E. *See* Lewis, Tian, Vainberg, and Cowan.
Vainio, O., Dunon, D., Aïssi, F., Dangy, J.-P., McNagny, K.M., and Imhof, B.A. HEMCAM, an adhesion molecule expressed by c-kit$^+$ hemopoietic progenitors. 135:1655.
Vaisberg, E.A., Grissom, P.M., and McIntosh, J.R. Mammalian cells express three distinct dynein heavy chains that are localized to different cytoplasmic organelles. 133:831.
Vale, R.D. Switches, latches, and amplifiers: Common themes of G proteins and molecular motors. 135:291.
— *See* Niclas, Allan, and Vale.
Valle, D. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Vallee, R.B. *See* Echeverri, Paschal, Vaughan, and Vallee.
Valtz, N., and Herskowitz, I. Pea2 protein of yeast is localized to sites of polarized growth and is required for efficient mating and bipolar budding. 135:725.
Vance, P.J. *See* Sauter, Pelchen-Matthews, Bron, Marsh, LaBranche, Vance, Romano, Haggarty, Hart, Lee, and Hoxie.
van den Berg, M. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
van den Berghe, H. *See* Mertens, van der Schueren, van den Berghe, and David.
Van Den Berghe, H. *See* Steinfeld, Van Den Berghe, and David.
van der Bijl, P., Lopes-Cardozo, M., and van Meer, G. Sorting of newly synthesized galactosphingolipids to the two surface domains of epithelial cells. 132:813.
— *See* Burger, van der Bijl, and van Meer.
— *See* Zinkl, Zuk, van der Bijl, van Meer, and Matlin.
van der Klei, I. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
van der Loop, F.T.L., Schaart, G., Timmer, E.D.J., Ramaekers, F.C.S., and van Eys, G.J.J.M. Smoothelin, a novel cytoskeletal protein specific for smooth muscle cells. 134:401.
van der Schueren, B. *See* Mertens, van der Schueren, van den Berghe, and David.
van der Valk, M.A. *See* Dunia, Smit, van der Valk, Bloemendal, Borst, and Benedetti.
van der Ven, P.F.M. *See* Obermann, Gautel, Steiner, van der Ven, Weber, and Fürst.
van Erck, M.G.M. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
van Eys, G.J.J.M. *See* van der Loop, Schaart, Timmer, Ramaekers, and van Eys.
Van Haastert, P.J.M., Bishop, J.D., and Gomer, R.H. The cell density factor CMF regulates the chemoattractant receptor cAR1 in *Dictyostelium*. 134:1543.
van Hinsbergh, V.W.M. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
van Meer, G. *See* Burger, van der Bijl, and van Meer.
— *See* van der Bijl, Lopes-Cardozo, and van Meer.
— *See* Zinkl, Zuk, van der Bijl, van Meer, and Matlin.
van Steensel, B. *See* Ludérus, van Steensel, Chong, Sibon, Cremers, and de Lange.
van Veldhoven, P.P. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Vasilakos, J. *See* Deshmukh, Vasilakos, Deckwerth, Lampe, Shivers, and Johnson.
Vaughan, K.T. *See* Echeverri, Paschal, Vaughan, and Vallee.
Veenhuis, M. *See* Distel, Erdmann, Gould, Blobel, Crane, Cregg, Dodt, Fujiki, Goodman, Just, Kiel, Kunau, Lazarow, Mannaerts, Moser, Osumi, Rachubinski, Roscher, Subramani, Tabak, Tsukamoto, Valle, van der Klei, van Veldhoven, and Veenhuis.
Verkhovsky, A.B. *See* Svitkina, Verkhovsky, and Borisy.
Vermeer, M.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Verschoor, A., Srivastava, S., Grassucci, R., and Frank, J. Native 3D structure of eukaryotic 80S ribosome: Morphological homology with the *E. coli* 70S ribosome. 133:495.
Verwoerd, D. *See* Wubbolts, Fernandez-Borja, Oomen, Verwoerd, Janssen, Calafat, Tulp, Dusseljee, and Neefjes.
Viard, I. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Vilquin, J.-T., Kinoshita, I., Roy, B., Goulet, M., Engvall, E., Tomé, F., Fardeau, M., and Tremblay, J.P. Partial laminin α2 chain restoration in α2 chain-deficient *dy/dy* mouse by primary muscle cell culture transplantation. 133:185.
Vincent, I., Rosado, M., and Davies, P. Mitotic mechanisms in Alzheimer's disease? 132:413.
Virta, H. *See* Peränen, Auvinen, Virta, Wepf, and Simons.
Visa, N., Izaurralde, E., Ferreira, J., Daneholt, B., and Mattaj, I.W. A nuclear cap-binding complex binds Balbiani ring pre-mRNA cotranscriptionally and accompanies the ribonucleoprotein particle during nuclear export. 133:5.
Visser, G.D. *See* Chi, Adam, Visser, and Adam.
Viswanathan, M.N. *See* Swartzman, Viswanathan, and Thorner.
Vitale, N. *See* Chasserot-Golaz, Vitale, Sagot, Delouche, Dirrig, Pradel, Henry, Aunis, and Bader.
Vithalani, K. *See* Larochelle, Vithalani, and De Lozanne.
Vogel, F. *See* Ruiz, Brinkmann, Ledermann, Behrend, Grund, Thalhammer, Vogel, Birchmeier, Günthert, Franke, and Birchmeier.
Vogel, S.S., Blank, P.S., and Zimmerberg, J. Poisson-distributed active fusion complexes underlie the control of the rate and extent of exocytosis by calcium. 134:329.
Voigt, S., Jungnickel, B., Hartmann, E., and Rapoport, T.A. Signal sequence–dependent function of the TRAM protein during early phases of protein transport across the endoplasmic reticulum membrane. 134:25.
Volkmer, H., Leuschner, R., Zacharias, U., and Rathjen, F.G. Neurofascin induces neurites by heterophilic interactions with axonal NrCAM while NrCAM requires F11 on the axonal surface to extend neurites. 135:1059.
Volpin, D. *See* Braghetta, Fabbro, Piccolo, Marvulli, Bonaldo, Volpin, and Bressan.
von der Mark, K. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
von Figura, K. *See* Kasper, Dittmer, von Figura, and Pohlmann.
von Reyher, U. *See* Günthert, Sträter, von Reyher, Henne, Joos, Koretz, Moldenhauer, Krammer, and Möller.
Voorn-Brouwer, T. *See* Elgersma, Kwast, Klein, Voorn-Brouwer, van den Berg, Metzig, America, Tabak, and Distel.
Vornehm, S.I. *See* Erdmann, Müller, Bahrami, Vornehm, Mayer, Bruckner, von der Mark, and Burkhardt.
Vuori, K. *See* Frisch, Vuori, Kelaita, and Sicks.
— *See* Frisch, Vuori, Ruoslahti, and Chan-Hui.

W

Wada, J., Kumar, A., Liu, Z., Ruoslahti, E., Reichardt, L., Marvaldi, J., and Kanwar, Y.S. Cloning of mouse integrin α_v cDNA and role of the α_v-related matrix receptors in metanephric development. 132:1161.
Waddell, T.K. *See* Demaurex, Downey, Waddell, and Grinstein.
— *See* Fukushima, Waddell, Grinstein, Goss, Orlowski, and Downey.
Waddle, J.A., Karpova, T.S., Waterston, R.H., and Cooper, J.A. Movement of cortical actin patches in yeast. 132:861.
Wahle, S., and Stoffel, W. Membrane topology of the high-affinity L-glutamate transporter (GLAST-1) of the central nervous system. 135:1867.
Walsh, F.S. *See* Fazeli, Wells, Hobbs, and Walsh.
Walsh, K. *See* Andrés and Walsh.
Walter, P. *See* Ng, Brown, and Walter.
— *See* Ng and Walter.
Walter, W. *See* Schilke, Forster, Davis, James, Walter, Laloraya, Johnson, Miao, and Craig.
Walther, I. *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Walton, P.A. *See* Schroeder, Leventis, Shahinian, Walton, and Silvius.

Wang, C., Pralong, W.-F., Schulz, M.-F., Rougon, G., Aubry, J.-M., Pagliusi, S., Robert, A., and Kiss, J.Z. Functional *N*-methyl-D-aspartate receptors in O-2A glial precursor cells: A critical role in regulating polysialic acid–neural cell adhesion molecule expression and cell migration. 135:1565.
Wang, F.-S. *See* Sydor, Su, Wang, Xu, and Jay.
Wang, H., Xie, Z., and Scott, R.E. Differentiation modulates the balance of positive and negative Jun/AP-1 DNA binding activities to regulate cellular proliferative potential: Different effects in nontransformed and transformed cells. 135:1151.
Wang, K. *See* Corrado, Rafael, Mills, Cole, Faulkner, Wang, and Chamberlain.
Wang, N. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Wang, R.-H. *See* Lazarovits, Naim, Rodriguez, Wang, Fire, Bird, Henis, and Roth.
Wang, S. *See* Deng, Curriden, Wang, Rosenberg, and Loskutoff.
— *See* Tao, Edwards, Tubb, Wang, Bryan, and McCrea.
Wang, S.-S. *See* Henderson, Allsopp, Spector, Wang, and Harley.
Wang, S.X. *See* Kinose, Wang, Kidambi, Moncman, and Winkelmann.
Wang, W., Hansen, P.A., Marshall, B.A., Holloszy, J.O., and Mueckler, M. Insulin unmasks a COOH-terminal Glut4 epitope and increases glucose transport across T-tubules in skeletal muscle. 135:415.
Wang, X.M., Peloquin, J.G., Zhai, Y., Bulinski, J.C., and Borisy, G.G. Removal of MAP4 from microtubules in vivo produces no observable phenotype at the cellular level. 132:345.
Wang, Y.-L. *See* Anderson, Wang, and Small.
Wang, Y.-l. *See* Wheatley and Wang.
Wang, Z.-Z., Hardy, S.F., and Hall, Z.W. Membrane tethering enables an extracellular domain of the acetylcholine receptor α subunit to form a heterodimeric ligand-binding site. 135:809.
Warmerdam, P.A.M., Long, E.O., and Roche, P.A. Isoforms of the invariant chain regulate transport of MHC class II molecules to antigen processing compartments. 133:281.
Warren, G. *See* Sönnichsen, Watson, Clausen, Misteli, and Warren.
Warrick, H.M. *See* Goodson, Anderson, Warrick, Pon, and Spudich.
Waters, M.G. *See* Harris and Waters.
— *See* Ktistakis, Brown, Waters, Sternweis, and Roth.
— *See* Sapperstein, Lupashin, Schmitt, and Waters.
Waterston, R.H. *See* Hoppe and Waterston.
— *See* Waddle, Karpova, Waterston, and Cooper.
Watson, R. *See* Sönnichsen, Watson, Clausen, Misteli, and Warren.
Watt, F.M. *See* Ruhrberg, Hajibagheri, Simon, Dooley, and Watt.
Weber, C., Alon, R., Moser, B., and Springer, T.A. Sequential regulation of α4β1 and α5β1 integrin avidity by CC chemokines in monocytes: Implications for transendothelial chemotaxis. 134:1063.
Weber, K. *See* Obermann, Gautel, Steiner, van der Ven, Weber, and Fürst.
Wedaman, K.P., Meyer, D.W., Rashid, D.J., Cole, D.G., and Scholey, J.M. Sequence and submolecular localization of the 115-kD accessory subunit of the heterotrimeric kinesin-II ($KRP_{85/95}$) complex. 132:371.
Wegehingel, S. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
Wegener, A.-M.K. *See* Dietrich, Neisig, Hou, Wegener, Gajhede, and Geisler.
Weich, H.A. *See* Koolwijk, van Erck, de Vree, Vermeer, Weich, Hanemaaijer, and van Hinsbergh.
Weich, S. *See* Scherthan, Weich, Schwegler, Härle, Heyting, and Cremer.
Weidner, K.M. *See* Sachs, Weidner, Brinkmann, Walther, Obermeier, Ullrich, and Birchmeier.
Weil, M., Jacobson, M.D., Coles, H.S.R., Davies, T.J., Gardner, R.L., Raff, K.D., and Raff, M.C. Constitutive expression of the machinery for programmed cell death. 133:1053.
— *See* Jacobson, Weil, and Raff.
Wein, H., Foss, M., Brady, B., and Cande, W.Z. DSK1, a novel kinesin-related protein from the diatom *Cylindrotheca fusiformis* that is involved in anaphase spindle elongation. 133:595.
Weiner, J.A. *See* Hecht, Weiner, Post, and Chun.
Weintraub, H. *See* Gogos, Thompson, Lowry, Sloane, Weintraub, and Horwitz.
Weis, K. *See* Palacios, Weis, Klebe, Mattaj, and Dingwall.
Weisman, L.S. *See* Hill, Catlett, and Weisman.
Weiss, E., and Winey, M. The *Saccharomyces cerevisiae* spindle pole body duplication gene *MPS1* is part of a mitotic checkpoint. 132:111.
Weiss, M.C. *See* Imaizumi-Scherrer, Faust, Bénichou, Hellio, and Weiss.
Weitzer, G. *See* Milner, Weitzer, Tran, Bradley, and Capetanaki.
Wellems, T.E. *See* Guinet, Dvorak, Fujioka, Keister, Muratova, Kaslow, Aikawa, Vaidya, and Wellems.
Wells, A. *See* Chen, Murphy-Ullrich, and Wells.
Wells, D.J. *See* Fazeli, Wells, Hobbs, and Walsh.
Wells, W.A.E., and Murray, A.W. Aberrantly segregating centromeres activate the spindle assembly checkpoint in budding yeast. 133:75.
Wendland, B., McCaffery, J.M., Xiao, Q., and Emr, S.D. A novel fluorescence-activated cell sorter–based screen for yeast endocytosis mutants identifies a yeast homologue of mammalian eps15. 135:1485.
Wennerberg, K., Lohikangas, L., Gullberg, D., Pfaff, M., Johansson, S., and Fässler, R. β1 integrin–dependent and –independent polymerization of fibronectin. 132:227.
Wepf, R. *See* Eaton, Wepf, and Simons.
— *See* Peränen, Auvinen, Virta, Wepf, and Simons.
— *See* Zabel, Doye, Tekotte, Wepf, Grandi, and Hurt.
Werb, Z. *See* Alexander, Howard, Bissell, and Werb.
— *See* Sympson, Talhouk, Alexander, Chin, Clift, Bissell, and Werb.
Weskamp, G., Krätzschmar, J., Reid, M.S., and Blobel, C.P. MDC9, a widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains. 132:717.
West, M.A. *See* Simpson, Bright, West, Newman, Darnell, and Robinson.
Westenbroek, R.E. *See* Sakurai, Westenbroek, Rettig, Hell, and Catterall.
Westlin, W.F. *See* Yoshida, Westlin, Wang, Ingber, Rosenzweig, Resnick, and Gimbrone.
Wewer, U.M. *See* Vachon, Loechel, Xu, Wewer, and Engvall.
Wheatley, S.P., and Wang, Y.-l. Midzone microtubule bundles are continuously required for cytokinesis in cultured epithelial cells. 135:981. Correction. 135:1679.
Wheelock, M.J. *See* Islam, Carey, Wolf, Wheelock, and Johnson.
—*See* Uchida, Honjo, Johnson, Wheelock, and Takeichi.
Whitaker, M. *See* Wilding, Wright, Patel, Ellis-Davies, and Whitaker.
White, E. *See* Rao, Perez, and White.
White, J.M. *See* Danieli, Pelletier, Henis, and White.
Whitney, J.A. *See* Balda, Whitney, Flores, González, Cereijido, and Matter.
Whittingham, D.G. *See* Kono, Jones, Bos-Mikich, Whittingham, and Carroll.
Wiche, G. *See* Nikolic, Mac Nulty, Mir, and Wiche.
Wickner, W. *See* Xu and Wickner.
Wieland, F.T. *See* Faulstich, Auerbach, Orci, Ravazzola, Wegehingel, Lottspeich, Stenbeck, Harter, Wieland, and Tschochner.
— *See* Sohn, Orci, Ravazzola, Amherdt, Bremser, Lottspeich, Fiedler, Helms, and Wieland.
Wieschaus, E. *See* Müller and Wieschaus.
Wieslander, L. *See* Baurén, Jiang, Bernholm, Gu, and Wieslander.
Wigger, F. *See* Götz, Scholze, Clement, Joester, Schütte, Wigger, Frank, Spiess, Ekblom, and Faissner.
Wilde, A., and Brodsky, F.M. In vivo phosphorylation of adaptors regulates their interaction with clathrin. 135:635.
Wilding, M., Wright, E.M., Patel, R., Ellis-Davies, G., and Whitaker, M. Local perinuclear calcium signals associated with mitosis-entry in early sea urchin embryos. 135:191.
Wilkemeyer, M.F. *See* Ramanathan, Wilkemeyer, Mittal, Perides, and Charness.
Willard, H.F. *See* Clemson, McNeil, Willard, and Lawrence.
Willhelm, C. *See* Robinett, Straight, Li, Willhelm, Sudlow, Murray, and Belmont.
Williams, B.C., Gatti, M., and Goldberg, M.L. Bipolar spindle attachments affect redistributions of ZW10, a *Drosophila* centromere/kinetochore component required for accurate chromosome segregation. 134:1127.
Williams, D.B. *See* Kim, Lingwood, Williams, Furuya, Manolson, and Grinstein.
Williams, S. *See* Scherer, Lederkremer, Williams, Fogliano, Baldini, and Lodish.
Williamson, T.L. *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.
Windle, J. *See* Osheim, Mougey, Windle, Anderson, O'Reilly.
Winey, M. *See* Weiss and Winey.
Winkelmann, D.A. *See* Kinose, Wang, Kidambi, Moncman, and Winkelmann.
Winsor, B. *See* Moreau, Madania, Martin, and Winsor.
Wirtz, E. *See* Häusler, Stierhof, Wirtz, and Clayton.
Witke, W. *See* Barkalow, Witke, Kwiatkowski, and Hartwig.
Wolf, D.E. *See* Ross, Daou, McKinnon, Condon, Lachyankar, Stephens, Kaplan, and Wolf.
Wolf, G.T. *See* Islam, Carey, Wolf, Wheelock, and Johnson.
Wong, D.H. *See* Koepp, Wong, Corbett, and Silver.
Wong, M.-H., and Filbin, M.T. Dominant-negative effect on adhesion by myelin Po protein truncated in its cytoplasmic domain. 134:1531.
Wong, M.L. *See* Field, Al-Awar, Rosenblatt, Wong, Alberts, and Mitchison.
Wong, P.C. *See* Xu, Marszalek, Lee, Wong, Folmer, Crawford, Hsieh, Griffin, and Cleveland.

Woodruff, L. *See* Marks, Woodruff, Ohno, and Bonifacino.
Woods, D.F., Hough, C., Peel, D., Callaini, G., and Bryant, P.J. Dlg protein is required for junction structure, cell polarity, and proliferation control in *Drosophila* epithelia. 134:1469.
Worm, M. *See* Schadendorf, Kern, Artuc, Pahl, Rosenbach, Fichtner, Nürnberg, Stüting, Stebut, Worm, Makki, Jurgovsky, Kolde, and Henz.
Wozniak, R.W. *See* Nehrbass, Rout, Maguire, Blobel, and Wozniak.
Wright, E.M. *See* Wilding, Wright, Patel, Ellis-Davies, and Whitaker.
Wright, J.M. *See* Liesi and Wright.
Wu, J.F. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Wu, J.-R. *See* Lawlis, Keezer, Wu, and Gilbert.
Wu, J.-X. *See* Huang, Wu, Fan, and Adamson.
Wu, X. *See* Jung, Wu, and Hammer.
Wubbolts, R., Fernandez-Borja, M., Oomen, L., Verwoerd, D., Janssen, H., Calafat, J., Tulp, A., Dusseljee, S., and Neefjes, J. Direct vesicular transport of MHC class II molecules from lysosomal structures to the cell surface. 135:611.
Wyke, J.A. *See* Fincham, Unlu, Brunton, Pitts, Wyke, and Frame.

X

Xia, Y., Gil, S.G., and Carter, W.G. Anchorage mediated by integrin α6β4 to laminin 5 (epiligrin) regulates tyrosine phosphorylation of a membrane-associated 80-kD protein. 132:727.
Xiao, Q. *See* Wendland, McCaffery, Xiao, and Emr.
Xie, Z. *See* Wang, Xie, and Scott.
Xu, A. *See* Sydor, Su, Wang, Xu, and Jay.
Xu, H. *See* Müsch, Xu, Shields, and Rodriguez-Boulan.
— *See* Vachon, Loechel, Xu, Wewer, and Engvall.
Xu, J., and Clark, R.A.F. Extracellular matrix alters PDGF regulation of fibroblast integrins. 132:239.
Xu, J., Zutter, M.M., Santoro, S.A., and Clark, R.A.F. PDGF induction of α_2 integrin gene expression is mediated by protein kinase C-ζ. 134:1301.
Xu, J.-Q., Harder, B.A., Uman, P., and Craig, R. Myosin filament structure in vertebrate smooth muscle. 134:53.
Xu, Z., Marszalek, J.R., Lee, M.K., Wong, P.C., Folmer, J., Crawford, T.O., Hsieh, S.-T., Griffin, J.W., and Cleveland, D.W. Subunit composition of neurofilaments specifies axonal diameter. 133:1061.
— **and Wickner, W.** Thioredoxin is required for vacuole inheritance in *Saccharomyces cerevisiae*. 132:787.
— *See* Marszalek, Williamson, Lee, Xu, Hoffman, Becher, Crawford, and Cleveland.

Y

Yagi, T. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yahara, I. *See* Aizawa, Sutoh, and Yahara.
Yamada, K.M. *See* Miyamoto, Teramoto, Gutkind, and Yamada.
Yamamoto, A., Guacci, V., and Koshland, D. Pds1p is required for faithful execution of anaphase in the yeast, *Saccharomyces cerevisiae*. 133:85. Correction. 134:267.
—, **Guacci, V., and Koshland, D.** Pds1p, an inhibitor of anaphase in budding yeast, plays a critical role in the APC and checkpoint pathway(s). 133:99.
Yamamoto, T. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yamanashi, Y. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yamashita, Y. *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.
Yan, G. *See* Zhang, Zapp, Yan, and Green.
Yanagida, M. *See* Saitoh, Takahashi, Nabeshima, Yamashita, Nakaseko, Hirata, and Yanagida.
Yang, H. *See* Lu, Yang, and Raizada.
Yang, J.T., Rando, T.A., Mohler, W.A., Rayburn, H., Blau, H.M., and Hynes, R.O. Genetic analysis of α_4 integrin functions in the development of mouse skeletal muscle. 135:829.
Yang-Snyder, J.A. *See* Torres, Yang-Snyder, Purcell, DeMarais, McGrew, and Moon.
Yap, A.S. *See* Brieher, Yap, and Gumbiner.
Yasuda, M. *See* Yasunaga, Yagi, Hanzawa, Yasuda, Yamanashi, Yamamoto, Aizawa, Miyauchi, and Nishikawa.
Yasunaga, M., Yagi, T., Hanzawa, N., Yasuda, M., Yamanashi, Y., Yamamoto, T., Aizawa, S., Miyauchi, Y., and Nishikawa, S.-I. Involvement of fyn tyrosine kinase in progression of cytokinesis of B lymphocyte progenitor. 132:91.
Yazaki, K. *See* Kanai, Kanai-Azuma, Noce, Saido, Shiroishi, Hayashi, and Yazaki.
Yehuda-Levenberg, S. *See* Simcha, Geiger, Yehuda-Levenberg, Salomon, and Ben-Ze'ev.
Yeon, M. *See* Mainiero, Pepe, Yeon, Ren, and Giancotti.
Yokota, S. *See* Satoh, Hirayoshi, Yokota, Hosokawa, and Nagata.
Yonemura, S. *See* Ando-Akatsuka, Saitou, Hirase, Kishi, Sakakibara, Itoh, Yonemura, Furuse, and Tsukita.
— *See* Hirao, Sato, Kondo, Yonemura, Monden, Sasaki, Takai, Tsukita, and Tsukita.
Yoshida, M., Westlin, W.F., Wang, N., Ingber, D.E., Rosenzweig, A., Resnick, N., and Gimbrone, M.A., Jr. Leukocyte adhesion to vascular endothelium induces E-selectin linkage to the actin cytoskeleton. 133:445.
Yoshida, S., Fujisawa-Sehara, A., Taki, T., Arai, K., and Nabeshima, Y. Lysophosphatidic acid and bFGF control different modes in proliferating myoblasts. 132:181.
Yoshimori, T., Keller, P., Roth, M.G., and Simons, K. Different biosynthetic transport routes to the plasma membrane in BHK and CHO cells. 133:247.
Young, S.G. *See* Huang, Wu, Cass, Erle, Corry, Young, Farese, and Sheppard.
Yu, Q.-C. *See* Allen, Yu, and Fuchs.
— *See* Dowling, Yu, and Fuchs.
Yu, W., Schwei, M.J., and Baas, P.W. Microtubule transport and assembly during axon growth. 133:151.
Yurimoto, H. *See* Sakai, Saiganji, Yurimoto, Takabe, Saiki, and Kato.

Z

Zabel, U., Doye, V., Tekotte, H., Wepf, R., Grandi, P., and Hurt, E.C. Nic96p is required for nuclear pore formation and functionally interacts with a novel nucleoporin, Nup188p. 133:1141.
Zacharias, U. *See* Volkmer, Leuschner, Zacharias, and Rathjen.
Zanetti, A. *See* Del Maschio, Zanetti, Corada, Rival, Ruco, Lampugnani, and Dejana.
Zanin, M.K.B. *See* Balsamo, Leung, Ernst, Zanin, Hoffman, and Lilien.
Zanone, R. *See* French, Hahne, Viard, Radlgruber, Zanone, Becker, Müller, and Tschopp.
Zapp, M.L. *See* Zhang, Zapp, Yan, and Green.
Zeile, W.L., Purich, D.L., and Southwick, F.S. Recognition of two classes of oligoproline sequences in profilin-mediated acceleration of actin-based *Shigella* motility. 133:49.
Zeng, Y. *See* Arnoult, Zeng, and Florman.
Zerial, M. *See* Ullrich, Reinsch, Urbé, Zerial, and Parton.
Zhai, Y., Kronebusch, P.J., Simon, P.M., and Borisy, G.G. Microtubule dynamics at the G_2/M transition: Abrupt breakdown of cytoplasmic microtubules at nuclear envelope breakdown and implications for spindle morphogenesis. 135:201.
— *See* Wang, Peloquin, Zhai, Bulinski, and Borisy.
Zhang, C.X., Lee, M.P., Chen, A.D., Brown, S.D., and Hsieh, T.-S. Isolation and characterization of a *Drosophila* gene essential for early embryonic development and formation of cortical cleavage furrows. 134:923.
Zhang, G., Zapp, M.L., Yan, G., and Green, M.R. Localization of HIV-1 RNA in mammalian nuclei. 135:9.
Zhang, H. *See* Lastres, Letamendía, Zhang, Rius, Almendro, Raab, López, Langa, Fabra, Letarte, and Bernabéu.
Zhang, J.W., and Lazarow, P.B. Peb1p (Pas7p) is an intraperoxisomal receptor for the NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase: Peb1p itself is targeted to peroxisomes by an NH_2-terminal peptide. 132:325.
Zhao, X. *See* Montgomery, Becker, Siu, Lemmon, Cheresh, Pancook, Zhao, and Reisfeld.
Zhidkova, N.I. *See* Belkin, Zhidkova, Balzac, Altruda, Tomatis, Maier, Tarone, Koteliansky, and Burridge.
Zhu, X., Ohtsubo, M., Böhmer, R.M., Roberts, J.M., and Assoian, R.K. Adhesion-dependent cell cycle progression linked to the expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of the retinoblastoma protein. 133:391.
Ziegler, U. *See* Buchstaller, Kunz, Berger, Kunz, Ziegler, Rader, and Sonderegger.
— *See* Kunz, Ziegler, Kunz, and Sonderegger.
Zimmer, A. *See* Reynolds, Zimmer, and Zimmer.
Zimmer, A.M. *See* Reynolds, Zimmer, and Zimmer.

Zimmerberg, J. *See* Plonsky and Zimmerberg.
— *See* Vogel, Blank, and Zimmerberg.
Zimmerli, S., Majeed, M., Gustavsson, M., Stendahl, O., Sanan, D.A., and Ernst, J.D. Phagosome–lysosome fusion is a calcium-independent event in macrophages. 132:49.
Zinkl, G.M., Zuk, A., van der Bijl, P., van Meer, G., and Matlin, K.S. An antiglycolipid antibody inhibits Madin-Darby canine kidney cell adhesion to laminin and interferes with basolateral polarization and tight junction formation. 133:695.
Zioncheck, T.F. *See* Schwall, Chang, Godowski, Kahn, Hillan, Bauer, and Zioncheck.
Zirbel, R.M. *See* Kurz, Lampel, Nickolenko, Bradl, Benner, Zirbel, Cremer, and Lichter.
Zondag, G.C.M., Moolenaar, W.H., and Gebbink, M.F.B.G. Lack of association between receptor protein tyrosine phosphatase RPTPμ and cadherins. 134:1513.
Zuk, A. *See* Zinkl, Zuk, van der Bijl, van Meer, and Matlin.
Zutter, M.M. *See* Xu, Zutter, Santoro, and Clark.

SUBJECT INDEX VOLUMES 132–135

α_{1A}
rbA and BI isoforms of, expressed in rat and rabbit brain, form calcium channels having α_{1A} subunits with distinct molecular mass, pharmacology, and subcellular localization, 134:511
subunits, interaction with proteins involved in synaptic exocytosis, 134:511
Acanthamoeba, biochemical kinetic characterization of myosin-I ATPase of, 132:1053
Acetylcholine receptor
clustering of induced by agrin, mammalian muscle, requires tyrosine phosphorylation, 132:937
expression of, at developing neuromuscular junction, 134:465
α subunit of, formation of heterodimeric binding site by, effect of tethering on, 135:809
Acetylcholine receptor–inducing activity (ARIA), signaling, pathways in, 134:465
Acrosome reaction, regulated by glycoprotein agonist of zona pellucida (ZP3), during mammalian fertilization, 134:637
ActA protein, proline-rich repeats in, of *Listeria monocytogenes*, control actin-based motility, bacterial motility and VASP and profilin binding, 135:647
Actin
chaperonin-mediated folding of tubulin and, 132:1
during cell–cell adhesion, spatial organization and strength of cadherin, catenin and, 135:1899
effects of Rab8 on reorganization of microtubules and, 135:153
formation and elongation of bundles of, in bristles of *Drosophila*, 135:1291
formation of filament bundles of, ring canals of developing *Drosophila* follicles, 133:61
kinetics of capping protein on barbed ends of, effects of polyphosphoinositides on capping protein of, regulation of polymerization of, 135:169
leukocyte adhesion to vascular endothelium induces E-selectin linkage to cytoskeleton of, 133:445
motility in *Shigella* based on, two classes of oligoproline sequences in profilin-mediated acceleration, 133:49
movement of cortical patches of, yeast, 132:861
myosin I proteins required for polarization of cytoskeleton of, synthetic lethality screen identifies novel yeast myosin I gene, *MYO5*, 133:1277
organization of, bristle morphology, viability, affected by actin-capping protein mutations in *Drosophila*, 133:1293
overexpression of cofilin stimulates bundling of filaments of, membrane ruffling, cell movement, *Dictyostelium*, 132:335
purse-string assembly of, during wound healing in chick embryo, roles of cadherin, myosin and Rho in, 135:1097
relationship of dystrophin to, 135:661
role of glycolipid-based domains, protein kinases and cytoskeleton of, endocytosis of GPI-anchored proteins in human lymphocytes, 133:791
in *Saccharomyces*, function in vacuole inheritance of myosin and, 135:1535
α-smooth muscle, retards migratory behavior in fibroblasts, 134:67
Actin cytoskeleton
linkage of E-cadherin to, in strengthening cell adhesion, 134:549
translocation of Src kinase to cell periphery mediated by, controlled by G proteins, 135:1551
Actin filaments, coordinate forward translocation of cell body with protrusion in lamellipodium, 134:1209
Actin polymerization
effected by elongation factor 1 α, 135:1309
effects of capping protein and polyphosphoinositides on, 135:169
α-actin promoter, neural CAM expression in skeletal muscle controlled by, 135:241
Actin-based motility, controlled by proline-rich repeats in ActA protein of *Listeria monocytogenes*, 135:647
Actin-binding domain I, not essential for dystrophin function, 134:873
Actin-binding motif is unique to epsilon isoform of protein kinase C, participates in regulation of synaptic function, 132:77
Actin-binding protein, function of 34k-Dalton, effects on filopodia in *Dictyostelium*, 135:965
Actin-binding protein elongation factor, effect of F-actin on binding of aminoacyl-tRNA to, pH sensitivity of, 135:953
Actin-depolymerizing factor, along with cofilin is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 132:871
Actin-related proteins, centractin and spectrin interact to mediate dynactin associations, 135:1815
Activated receptors, endocytosis of, clathrin-coated pit formation, 132:1025
Active transport, of photoactivated tubulin molecules in growing axons revealed by new electron microscopic analysis, 133:1347
Activin-A, mesoderm inducing factor, activates position-specific recognition of synergy site of fibronectin in vivo, 134:227
Activity zones, of synaptic vesicle recycling pathways, release characteristics of, 135:797
Ad5, replication domains, located adjacent to nuclear domain (ND)10s, 134:815
γ-Adaptin, AP-1 adaptor complex binds to immature secretory granules from PC12 cells, regulated by ADP-ribosylation factor, 132:523
Adaptor complexes, novel adaptor-related protein complex, 133:749
Adaptors, in vivo phosphorylation of, interactions of clathrin-coated vesicles with, 135:635
Adenomatous polyposis coli protein, increased levels at crypt/villus boundary in intestine, 134:165
Adenomatous polyposis coli tumor suppressor protein, localization to plasma membrane sites involved in active cell migration, 134:165
Adenovirus, activation of transcription factor NF-κB by E3/19K protein in, requires ER retention, 132:511
Adherens junction
endothelial, polymorphonuclear leukocyte adhesion triggers disorganization of, 135:497
estrogen-dependent c-JunER protein causes reversible loss of mammary epithelial cell polarity involving destabilization of, 132:1115
key role in cell–cell adhesion and in organizing cells into epithelia, 134:133
role of Armadillo in assembly of, 134:1283
symplekin in, 134:1003
Adherens junction components, E-cadherin, α- and β-catenin, colocalize with DP-NTP and desmosomal cadherins at cell–cell borders, 134:985
Adhesion
activities of *Wnt*-1 class of secreted signaling factors are antagonized by *Wnt*-5A class and by dominant negative cadherin in early *Xenopus* development, 133:1123
alcohol inhibits cell–cell, mediated by human L1, 133:381
anchorage mediated by integrin α6β4 to laminin 5 regulates tyrosine phosphorylation of membrane-associated protein, epiligrin, 132:727
antiglycolipid antibody inhibits Madin-Darby canine kidney cell, to laminin, interferes with basolateral polarization and tight junction formation, 133:695
of cadherin-associated catenin proteins, in synaptic junctions, 135:767
cadherin-dependent, 134:1519
catenins and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 132:451
C-cadherin-mediated, 135:487
CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 132:437
cell cycle progression dependent on, linked to expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of retinoblastoma protein, 133:391
characterization of HEMCAM molecule of, expressed by hemopoietic progenitors, 135:1655
cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 132:1161
defects in cell–cell, when Armadillo is depleted from onset of embryogenesis, 134:133
deficiency of Src family kinases p59/61^{hck} and p58$^{c\text{-}fgr}$ results in defective functions of neutrophils dependent on, 133:895
E-cadherin–mediated, 134:549
effect of nonepithelial cadherin by epithelial cell to disrupt cell–cell, 135:1643
glycosylation of CD44 implicated in CD44-mediated cell adhesion, hyaluronan, 132:1199
heterophilic interactions of DM-GRASP, GRASP-NgCAM interactions involved in neurite extension, 133:657
homophilic, role of truncated myelin Po protein in, 134:1531
of leukocytes to vascular endothelium induces E-selectin linkage to actin cytoskeleton, 133:445
loss of, early event in CD95-mediated apoptosis, 134:1089

mice expressing mutant desmosomal cadherin exhibit abnormalities in desmosomes, proliferation, and epidermal differentiation, 133:1367
in neurite outgrowth, interaction between NgCAM and axonin-1, 135:1593
neuroglian-mediated, induces assembly of membrane skeleton at cell contact sites, 133:647
polymorphonuclear leukocyte, effects on endothelial adherens junctions, 135:497
spatial organization and strength of cadherin, catenin and actin during cell–cell, 135:1899
stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 132:741
strength of, between cell and substrate, important contributor to cell migration rate, 134:1551
tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 132:681
urokinase type plasminogen activator receptor (uPAR)–mediated, 134:1563
Adhesion molecules
characterization of neuronal cell, at node of Ranvier, 135:1355
L1, and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, human neural cell, 132:475
Adhesion receptors, visualization of CD2 interaction with LFA-3 and determination of two-dimensional dissociation constant for, contact area, 132:465
Adhesion-triggering, versus chemotactic responses, in lymphoid cells, 134:255
Adipocytes, separate intracellular compartment of glucose transporter (GLUT-4) in, 134:625
ADP ribosylation factor
activation of, required for formation of coatomer-coated vesicles from Golgi-enriched membranes, 134:295
AP-1 adaptor complex regulated by, binds to immature secretory granules from PC12 cells, 132:523
effector function of, 134:935
ADP ribosylation factor GTP-binding protein, and Rho-related GTPases, specifically regulate distinctive actin-based cell surface structures, 134:935
a-factor, effects of Ste6 in cell fusion during mating of *Saccharomyces* on, 135:741
a-factor mating pheromone, in yeast, cell fusion during mating requires high levels of, 135:1727
Agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine phosphorylation, 132:937
AGT, dimerization of, in peroxisomal and mitochondrial import processes, mistargeting of, 135:939
Alcohol, inhibits cell–cell adhesion mediated by human L1, 133:381
Alport syndrome, related to collagen α3(IV) deficiency in mice, 135:1403
Aluminum fluoride, treatment of ADP ribosylation factor (ARF)6-transfected cells with, results in formation of peripheral plasma membrane protrusions, 134:935
Alzheimer's disease, mitotic mechanisms in, 132:413
Amino acid permeases, roles of Shr3p and COPII in packaging of, 135:585
Aminoacyl-tRNA, effect of F-actin on binding of EF-1α to, pH sensitivity of, 135:953
Aminopeptidase I, identification of cytoplasm to vacuole targeting determinant in, 132:999
Aminopterin, import of DHFR hybrid protein into glycosomes in vivo not inhibited by, folate analogue, 132:311
Amnion, contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, human, 132:1189
Anaphase
detection of protein kinase activity specifically activated at transition with metaphase, 132:635
DSK1 involved in spindle elongation during, novel kinesin-related protein from diatom *Cylindrotheca fusiformis*, 133:595
force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but only at kinetochore during, 132:1093
Pds1p is inhibitor of, budding yeast, plays critical role in APC and checkpoint pathways, 133:99
Pds1p required for execution of, yeast, *Saccharomyces cerevisiae*, 133:85
ZW10 in regulation of chromosome separation or movement at onset of, 134:1127
Anaphase B, later phase of separation for centrosomes associated with spindles, does not occur in free centrosome separation, 134:103
Angiotensin II, stimulation of transporter genes in brain neurons, MAP kinase signaling pathway activated by, 135:1609
Anisotropy, influence of M-phase chromatin on, microtubule asters, 133:125
Ankyrins
identification of small cytoplasmic, binds βIΣ* spectrin, associates with Golgi apparatus, Ank_{G119}, kidney and muscle, 133:819
neuroglian-mediated cell adhesion induces assembly of membrane skeleton at cell contact sites, 133:647
Annexin, identification, localization, and functional implications of an abundant, Nematode, 132:1079
Annexin II, in exocytosis, catecholamine secretion requires translocation of p36 to subplasmalemmal region in chromaffin cells, 133:1217
Anoikis
roles of Jun-N-terminal kinases and ICE-related cysteine protease in, 135:1377
subset of apoptosis, regulated by focal adhesion kinase, 134:793
Antigens, isoforms of invariant chain regulate transport of MHC class II molecules to processing compartments of, 133:281
Antisense RNA, inhibition of laminin α1-chain expression leads to alteration of basement membrane assembly and cell differentiation, 133:417
AP-1, CD437 activates and regulates, inhibits growth and induces apoptosis in human melanoma cells, 135:1889
AP-1 adaptor complex, binds to immature secretory granules from PC12 cells, regulated by ADP-ribosylation factor, 132:523
AP-1 DNA binding, attributed to Jun proteins, in transformed and nontransformed cells, 135:1151
AP-2 adaptor, clathrin coat assembly on dense lysosomes initiated by, 135:1801
Apical endosomes, of MDCK cells, interconnection of basolateral endosomes and, polarized sorting mechanism of, 135:139
Apoptosis
Bcl-2 inhibits retinoic acid–induced, neural differentiation of embryonal stem cells, 132:955
in fibroblasts, role of $pp125^{FAK}$ to suppress, 135:1383
in human melanoma cells, activation and regulation of AP-1 by CD437 inhibits growth and induces, 135:1889
induced by p53 in postmitotic neurons, 135:1085
induction of DNA synthesis and, cardiac myocytes by E1A oncoprotein, 133:325
ligand expression in immune-privileged tissues and coexpression in adult tissues characterized by cell turnover by, Fas and Fas ligand in embryos and adult mice, 133:335
of mammary epithelial cells in mice, effect of metalloproteinase inhibitor on, 135:1669
NGF and ICE family proteases inhibit metabolic and genetic events associated with neuronal, 135:1341
p53-dependent, lamin degradation facilitates nuclear events during, 135:1441
presence of DNA breaks with single-base 3′ overhangs during, 135:1369
roles of Jun-N-terminal kinases and ICE-related cysteine protease in, 135:1377
transcription factor NF-κB in, 134:13
Apoptotic activity, of cysteine protease, link between processing and, 135:479
Arm, required for zonula adherens formation, 134:149
armadillo, zonula adherens formation, and maintenance of polarized blastoderm epithelium in *Drosophila*, 134:149
Armadillo
absence of, prevents assembly of adherens junctions, 134:133
in *Drosophila* embryogenesis, required for adherens junction assembly, cell polarity, and morphogenesis, 134:133
role in assembly of adherens junction, 134:1283
role in Wingless signaling, 134:1283
Armadillo repeats, *PF16* encodes a protein with, localizes to single microtubule of central apparatus in *Chlamydomonas* flagella, 132:359
arp2-H330L mutation, affects polarization of actin distribution, 134:117
Arp2p, cortical localization, and colocalization with actin patches, 134:117
Assembly, of desmoplakin into desmosomes, requires NH2-terminal sequences, 134:985
Astrocytes, disrupted glial fibrillary acidic protein network in, vimentin knockout mice, 133:853
ATA, inhibitor of putative apoptotic endonuclease, inhibits camptothecin-induced neuronal death, 134:757
ATP
function in clathrin dissociation of auxilin, 135:925
interactions of PDZ domain-proteins with, 135:753
oligomerization and maturation of Na,K-ATPase, interaction of cytoplasmic NH_2 terminus of β subunit with α subunit, 133:1193

ATP hydrolysis, assembly of ER-associated protein degradation in vitro is dependent on cytosol, calnexin, and, 132:291
ATPase
biochemical kinetic characterization of myosin-I ATPase, *Acanthamoeba*, 132:1053
chloroplast protein import intermediate reveals distinct translocation machineries in two envelope membranes, 132:63
ATP-binding cassette transporter, *PAL1* gene product is, peroxisomes, *Saccharomyces cerevisiae*, 132:549
Autocrine mechanism, differentially expressed fibroblast growth factors regulate skeletal muscle development through paracrine and, 132:1151
Autophagy, ultrastructural and biochemical characterization of, higher plant cells subjected to carbon deprivation, control by supply of mitochondria with respiratory substrates, 133:1251
Auxilin, effects of hsc70 and ATP in J and clathrin domains of, 135:925
Axis formation, binding to cadherins antagonizes signaling activity of β-catenin during, *Xenopus*, 132:1105
Axon growth, microtubule transport and assembly during, 133:151
Axon vesicles, of squid, plus-end motors override minus-end motors during transport, 135:383
Axonin-1
association with kinases, intracellular signaling during neurite fasciculation triggered by, effects of clustering with Ng-CAM, 135:253
role in neurite outgrowth, interaction between NgCAM and, 135:1593
Axons
active transport of photoactivated tubulin molecules in growing, revealed by new electron microscopic analysis, 133:1347
characterization of cell adhesion molecules at node of Ranvier of myelinated, 135:1355
effect of NF-H on neurofilament regulation of, 135:711
selective stabilization of Tau in, microtubule-associated protein 2C in cell bodies and dendrites, polarized localization of cytoskeletal proteins in mature neurons, 132:667
subunit composition of neurofilaments specifies diameter of, 133:1061
tubulin transport in neurons, 133:1355
B lymphoblasts, biogenesis of MHC class II compartment in human I-Cell Disease, 132:769
B lymphocyte progenitor, cytokinesis progression of, involvement of Fyn tyrosine kinase, 132:91
Bacterial motility, controlled by proline-rich repeats in ActA protein of *Listeria monocytogenes*, 135:647
Baculovirus GP64, formation and kinetics of fusion pore induced by, 135:1831
Balbiani ring, nuclear cap-binding complex binds pre-mRNA in, cotranscriptionally, accompanies ribonucleoprotein particle during nuclear export, 133:5
Basal cells, in mice, *distal-less* homeodomain gene regulates differentiation of, 135:1879
Basement membranes, inhibition of laminin α1-chain expression leads to alteration of assembly of, cell differentiation, 133:417
Basic fibroblast growth factor
cooperative effect of TNFα, VEGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 132:1177
and lysophosphatidic acid control different modes in proliferating myoblasts, 132:181
Basic helix-100p-helix (bHLH), expression of, converts variety of cultured cells into skeletal muscle, 134:837
Basolateral endosomes, of MDCK cells, interconnection of apical endosomes and, polarized sorting mechanism of, 135:139
bazooka, zonula adherens formation, and maintenance of polarized blastoderm epithelium in *Drosophila*, 134:149
Bcl-2
down-regulates activity of transcription factor NF-κB induced upon apoptosis, 134:13
inhibits NF-κB activity in apoptosis by down-modulating transactivating potential of nuclear p65, 134:13
inhibits retinoic acid–induced apoptosis during neural differentiation of embryonal stem cells, 132:955
Bead binding, vitronectin, and integrin-associated protein (IAP), 134:1313
Beaded-chain filaments, filensin and phakinin form, coassemble de novo in cultured cells, 132:643
Becker muscular dystrophy, 134:93, 134:873
BED1 gene, required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 132:137
Bem1p
associations among PH and SH3 domain–containing proteins and Rho-type GTPases in yeast, 133:879
yeast *src* homology region 3 domain-binding proteins involved in bud formation, 133:865
BHK cells, different biosynthetic transport routes to plasma membrane in CHO cells and, 133:247
Binding
characteristics of L- and P-selectin and neutrophils, 135:523
effect of Jun proteins on AP-1 DNA, transformed and nontransformed cells, 135:1151
heterodimeric site formed by acetylcholine receptor α subunit, effect of tethering on, 135:809
homophilic mechanism of C-cadherin, 135:487
of hyaluronic acid, effected by variants of CD44, 135:1139
Biosynthesis, of COOH-terminal variant isoforms of proprotein convertase, 135:1261
Biotinylated tubulin, microtubule transport and assembly during axon growth, 133:151
1,2-bis-(o-Aminophenoxy)-ethane-N,N,N′,N′-tetra acetic acid, biochemically distinct steps revealed with NEM, GTPγS, and, assembly of nuclear pore, 132:5
BK2, mAb, precipitates cadherins independently of receptor-like protein tyrosine phosphatase (RPTPμ), 134:1513
Blunt-ended DNA, presence in apoptotic cells, 135:1369
BMP-6, expression in transgenic mice of, inhibition or stimulation of cell proliferation depending on the pattern of expression of, in formation of psoriasis, 135:227
Boc-Asp(OBzl)-CMK, specific IL-1β converting enzyme inhibitor, completely blocks CD95-triggered apoptosis, 134:1089
Bone diseases, expression of TGF-β2 in osteoblasts results in osteoporosis-like phenotype, 132:195
Bone formation, chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 in vitro, gene expression of parathyroid hormone/PTH-related peptide receptor, 133:457
Bone morphogenetic proteins
-6, expression in transgenic mice of, inhibition or stimulation of cell proliferation depending on the pattern of expression of, in formation of psoriasis, 135:227
processing of, limited by local availability of subtilisin-like proprotein convertases, 134:181
Boss protein, mutations in *Drosophila hook* gene inhibit endocytosis of boss transmembrane ligand into multivesicular bodies, 133:1205
Bouquet, chromosomal, pairing of mouse chromosome-8–specific probe in, 134:1109
BrdU, effect on mtDNA to analyze spatiotemporal constraints of replication, 135:883
Bristles, morphology of, actin organization, viability, affected by actin-capping protein mutations in *Drosophila*, 133:1293
BUB2, cell cycle arrest and survival of *tub4-1* cells dependent on, 134:429
Budding
associations among PH and SH3 domain–containing proteins and Rho-type GTPases in yeast, 133:879
axial, of Bud4p, 134:413
BED1 required for polarized growth and efficient emergence of, *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 132:137
from endosomes, novel class of clathrin-coated vesicles, 132:21
random patterns, in haploid and diploid *arp2-H330L* mutant cells, 134:117
in *Saccharomyces*, effect of mutants defective in Pea2p and Spa2p on, 135:725
yeast *src* homology region 3 domain-binding proteins involved in, 133:865
Budding site, targeting of Moloney murine leukemia virus Gag precursor to, 135:1841
Budding zone, of endoplasmic reticulum export complexes, COPII-coated, 135:19
Bud4p, expressed and localized to mother/bud neck in all cell types, 134:413
Bullous congenital ichthyosiform erythroderma, gene targeting at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 132:925
Bundles, of actin filaments, formation and elongation of, in bristles of *Drosophila*, 135:1291
Bundling, microtubule, by non-Claret disjunctional kinesin-like protein, 134:455
byr4, is novel suppressor of *ras1* in fission yeast, dosage-dependent inhibitor of cytokinesis, 133:1307
Cα, catalytic subunit, high expression of throughout all skeletal muscles of early fetus, 134:1241
C protein, in heterogeneous nuclear RNP (hnRNP) protein, bear a retention sequence, 134:1365

Ca^{2+}, extracellular, reduction of, key factor in hemichannel opening, 134:1019
Cab45, is novel Ca^{2+}-binding protein localized to Golgi lumen, 133:257
Cadherin
activities of *Wnt*-1 class of secreted signaling factors are antagonized by *Wnt*-5A class and by dominant negative, early *Xenopus* development, 133:1123
binding to, antagonizes signaling activity of β-catenin during axis formation in *Xenopus*, 132:1105
binding of plakoglobin to desmosomal, patterns of binding sites and topogenic potential, 133:359
catenin proteins associated with adhesion of, in synaptic junctions, 135:767
in cell–cell adhesion, 134:549
complexes, receptor-like protein tyrosine phosphatase (RPTPμ), lack of association with, 134:1513
during cell–cell adhesion, spatial organization and strength of catenin, actin and, 135:1899
mice expressing mutant desmosomal, exhibit abnormalities in desmosomes, proliferation, and epidermal differentiation, 133:1367
role in actin cable assembly during wound healing of, 135:1097
role in control of morphogenetic movements during development, 134:801
in squamous carcinoma cells, effect of N-cadherin on fibroblastic phenotype and E- and P-, 135:1643
Caenorhabditis, characteristic variations in hydrophobicity along myosin heavy chain rods of, 135:371
Caenorhabditis elegans
eat-5 and *unc-7* genes of, 134:537
gene *unc-89* encodes giant modular protein composed of Ig and signal transduction domains in, required for muscle M-line assembly, 132:835
identification, localization, and functional implications of an abundant Nematode annexin, 132:1079
ion channel subunit required for mechanotransduction in, sequence and transmembrane topology of MEC-4, 133:1071
tropinin T mutations in striated and nonstriated muscle cells of, 132:1061
Calcium
catenins and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 132:451
cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first mitotic division, 132:915
changes in cleavage-stage *Xenopus* embryos, associated with cleavage furrows, inhibition by heparin, 135:181
cortical granule exocytosis triggered by, 134:329
effects of calreticulin on homeostasis of, in endoplasmic reticulum, 135:1913
involved in contractile activity that regulates isoform expression and polysialylation of NCAM in cultured myotubes, protein kinase C, 132:969
phagosome–lysosome fusion is independent of, macrophages, 132:49
release in HSY cells conforms to steady-state mechanism involving regulation of inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$, 132:607
signaling near nucleus before nuclear envelope breakdown, entry into mitosis regulated by, 135:191
Calcium binding protein, Cab45 is novel, localized to Golgi lumen, 133:257
Calcium channel function, role in migration deficiency of weaver granule neurons, 134:477
Calcium channels
brain, α_{1A} subunits of, 134:511
intracellular, role in regulating muscle contraction, 134:885
Calmodulin
role of interaction of Spc110p and, proper assembly of spindle pole body components, 133:111
striatin binds, in presence of Ca^{2+}, 134:1051
Calnexin, assembly of ER-associated protein degradation in vitro is dependent on cytosol, ATP, and, 132:291
Calreticulin, regulates vinculin expression to control cell adhesiveness in endoplasmic reticulum, calcium homeostasis effected by, 135:1913
cAMP
in regulation of myosin II heavy chain-protein kinase C membrane association, 134:911
regulation of tyrosinase gene expression by, 134:747
cAMP receptor, modulation of, by conditioned medium factor (CMF), 134:1543
Camptothecin, neuronal apoptosis induced by, 134:757
Cap-binding proteins, nuclear cap-binding complex binds Balbiani ring pre-mRNA cotranscriptionally and accompanies ribonucleoprotein particle during nuclear export, 133:5
Capping protein
actin organization, bristle morphology, and viability are affected by actin-capping protein mutations in *Drosophila*, 133:1293
on barbed ends of actin, dynamics of, effects of polyphosphoinositides on, 135:169
and gelsolin, coordinated regulation of platelet actin filament barbed ends by, 134:389
translocation of to actin cytoskeleton, depends on gelsolin, 134:389
cAR1, cAMP receptor gene, disruption of, abolishes cAMP and conditioned medium factor (CMF) binding, 134:1543
Carbohydrate starvation, characterization of autophagy in higher plant cells subjected to carbon deprivation, control by supply of mitochondria with respiratory substrates, 133:1251
Carbon, characterization of autophagy in higher plant cells subjected to deprivation of, control by supply of mitochondria with respiratory substrates, 133:1251
Carboxypeptidase Y, Vps10p cycles between late-Golgi and prevacuolar compartments, sorting receptor for multiple yeast vacuolar hydrolases, 133:529
Carcinogenesis, epithelial, focal adhesion kinase in, 134:793
Carcinoma cells
binding of plakoglobin to desmosomal cadherins, patterns of binding sites and topogenic potential, 133:359
chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 in vitro, gene expression of parathyroid hormone/PTH-related peptide receptor, 133:457
Cartilage, *Col11a2* gene regulation in, 134:1573
Cartilage matrix metabolism, differential effect of parathyroid hormone on, 135:1179
Catalase peroxisomal targeting sequence, 134:849
Cataracts, human multidrug resistance 3–P-glycoprotein expression in transgenic mice induces lens membrane alterations leading to, 132:701
Catecholamine, secretion of requires translocation of p36 to subplasmalemmal region in chromaffin cells, annexin II in exocytosis, 133:1217
β-catenin
in adherens junction, 134:1283
association with fascin, in noncadherin complex, 134:1271
binding to cadherins antagonizes signaling activity of, axis formation in *Xenopus*, 132:1105
phosphorylation of, and regulation of cell function, 134:1519
in signaling, 134:1283
tyrosine phosphorylation/dephosphorylation of, in controlling cadherin function, 134:801
Catenins
adhesion of cadherin-associated proteins of, in synaptic junctions, 135:767
complexes formed with adenomatous polyposis coli protein, 134:165
during cell–cell adhesion, spatial organization and strength of cadherin, actin and, 135:1899
mediation of cadherin association with actin containing cytoskeleton mediated by, 134:801
role in cell and developmental signaling pathways, 134:1271
and Zonula Occludens-1 form complex during early stages in assembly of tight junctions, 132:451
CATGTG motifs, two surrounding the TATA box, involved in regulation of tyrosinase gene expression by cAMP in B16 melanoma cells, 134:747
Cathepsin B, direct role in myoblast growth and fusion, 134:837
Cation channels, sperm, ZP3-dependent activation of, 134:637
Caveolae
clustered folate receptors in, 134:1169
ligand-induced protease receptor translocation into, mechanism for regulating cell surface proteolysis of tissue factor–dependent coagulation pathway, 133:293
CC chemokines, differentially and selectively regulate avidity of integrins sharing common β subunits, 134:1063
C-cadherin, homophilic binding mechanism of, lateral dimerization in function of, adhesion mediated by, 135:487
CCV, effect of in vivo phosphorylation on, interactions of adaptors with, 135:635
CD2, visualization of interaction with LFA-3, determination of two-dimensional dissociation constant for adhesion receptors in contact area, 132:465
CD3γ, role in T cell receptor assembly, 132:299
CD44
CD95-mediated shedding of, 134:1089
glycosylation of, implicated in CD44-mediated cell adhesion to hyaluronan, 132:1199

–hyaluronan, interactions, in lymphocyte migration through secondary lymphoid organs, 134:1075
interaction between ERM proteins and, effect of phosphatidylinositol on, regulation mechanism and involvement of Rho in interaction between ERM proteins and, 135:37
proteins, binding properties of hyaluronic acid effected by variants of, 135:1139
CD45, exclusion of, effect on Lck in glycolipid-enriched membrane domains, 135:1515
CD47, mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 132:437
CD437, synthetic retinoid, activation and regulation of AP-1 by, inhibits growth and induces apoptosis in human melanoma cells, 135:1889
CDC2, phosphorylation, and radiation arrest, 134:963
Cdc42
cell type–specific roles for Rac, RhoL and, *Drosophila* oogenesis, 133:617
effect on actin polymerization in formation of wing hairs in *Drosophila*, 135:1277
cdc2 kinase, novel component of maturation-promoting factor besides, triggering of M-phase in starfish oocytes, 132:125
Ced-3/interleukin-1β converting enzyme family, role of Ced-3/ICE-family proteases in staurosporine-induced programmed cell death, 133:1041
Cek9, spatial and temporal expression of, tyrosine phosphorylation of, 135:781
Cell adhesion, β4 integrin required for, 134:559
Cell adhesiveness, in endoplasmic reticulum, calreticulin regulation of vinculin expression controls, 135:1913
Cell culture transplantation, partial laminin α2 chain restoration in α2 chain-deficient *dy/dy* mouse by primary muscle, 133:185
Cell cycle
aberrantly segregating centromeres activate spindle assembly checkpoint in budding yeast, 133:75
accumulation in S and G_2 phases of skeletal muscle cells due to lack of retinoblastoma protein, 135:441
G1 arrest in transformed and nontransformed cells caused by Taxol, 135:689
increases in calcium near nucleus are required to trigger entry into mitosis, 135:191
induction of DNA synthesis and apoptosis in cardiac myocytes by E1A oncoprotein, 133:325
mitosis-specific phosphorylation of vimentin by protein kinase C coupled with reorganization of intracellular membranes, 133:141
myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis, 132:657
spindle pole body component is phosphoprotein that is modified in a manner dependent on, *Saccharomyces cerevisiae*, 132:903
Cell cycle progression
adhesion-dependent, linked to expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of retinoblastoma protein, 133:391
14-3-3 proteins associate with phosphorylated simple epithelial keratins during, act as solubility cofactor, 133:345
Cell cycle regulation
of dynein association with membranes modulates microtubule-based organelle transport, 133:585
novel suppressor of *ras1* in fission yeast is dosage-dependent inhibitor of cytokinesis, *byr4*, 133:1307
Pds1p plays critical role in APC and checkpoint pathways, inhibitor of anaphase in budding yeast, 133:99
Pds1p required for execution of anaphase in yeast, *Saccharomyces cerevisiae*, 133:85
Cell death
in human melanoma cells, activation and regulation of AP-1 by CD437 inhibits growth and induces, 135:1889
induced by overexpression of p53 in postmitotic neurons, 135:1085
NGF and ICE family proteases inhibit metabolic and genetic events associated with neuronal, 135:1341
Cell differentiation
inhibition of laminin α1-chain expression leads to alteration of basement membrane assembly and, 133:417
regulated by p53 through transcriptional activity, 134:193
Cell division, in *Saccharomyces*, function of myosin and actin in rapid and directed vacuole movement during, 135:1535
Cell fusion
during mating of *Saccharomyces*, effects of Ste6 on **a**-factor in, 135:741
myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede, myogenesis, 132:657
in yeast, reduced levels of **a**-factor mating pheromone results in defective mutants during, 135:1727
Cell migration, active, localization of adenamatous polyposis coli tumor suppressor protein to plasma membrane sites involved in, 134:165
Cell movement, overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and, *Dictyostelium*, 132:335
Cell spreading, syndecan-1 mediates, transfected human lymphoblastoid cells, Raji cells, 132:1209
Cell surface transport, different biosynthetic transport routes to plasma membrane in BHK and CHO cells, 133:247
Cell survival, β4 integrin required for, 134:559
Cell–cell adhesion, mediated specifically by expression of E-cadherin, 134:549
Cell–cell coupling, *eat-5* and *unc-7* involved in, 134:537
Cellubrevin, partial colocalization between glucose transporter (GLUT-4) and, in adipocytes, 134:625
Centractin, functions of, dynactin associations mediated by interaction between spectrin and, 135:1815
Central apparatus, *PF*16 localizes to single microtubule of, *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 132:359
Central nervous system
membrane topology of L-glutamate transporter in, 135:1867
striatin exclusively expressed in, 134:1051
Centromeres
activate spindle assembly checkpoint in budding yeast, aberrantly segregating, 133:75
behavior after kinetochore destruction, 135:315
distinguished from remainder of chromosome by intensity of topoisomerase II α reactivity, 134:1097
distribution patterns in mouse and human meiocytes, 134:1109
elastic behavior of α-satellite DNA domain of, 135:545
Centrosomes
free, in normal and aphidicolin-treated *Drosophila* embryos, 134:103
maturation of, in mitotic human cells, polo-like kinase1 required for, 135:1701
Cerebellum
neurotrophins promote survival and development of neurons in, hypothyroid rats in vivo, 133:631
terminals of climbing and parallel fibers in, 134:511
Cerulenin, lethal effect of, in nuclear division and septation/cytokinesis, 134:949
cGMP, in regulation of myosin II heavy chain-protein kinase C membrane association, 134:911
Chagas' disease, 134:349
Chaperone proteins, Hsp70 class, prevent irreversible inactivation of proteins, allowing their reactivation under normal growth conditions, 134:1375
Chaperones, roles in protein degradation, 132:255
Chaperonins, mediates folding of actin and tubulin, 132:1
Charcot-Marie-Tooth, type 1B, peripheral neuropathy, and myelin Po protein, 134:1531
Chemokine, and classical chemoattractant receptors, biology of, and differential requirements for adhesion-triggering versus chemotactic responses in lymphoid cells, 134:255
Chemotaxis, transendothelial, facilitated by sequential modulation, 134:1063
Chick embryo, wound healing, actin purse-string assembly in, roles of cadherin, myosin and Rho in, 135:1097
Chick laminin-4, contains β2 chain, motor axon growth promoted by, effect of LRE in, 135:1583
Chimeric proteins, endocytosis of, 134:339
Chironomus tentans
dynamic transcription-dependent organization of pre-mRNA splicing factors in polytene nuclei, 133:929
nuclear cap-binding complex binds Balbiani ring pre-mRNA cotranscriptionally and accompanies ribonucleoprotein particle during nuclear export, 133:5
Chitin synthase proteins, temporal and spatial characterization of Chs2p and Chs3p in *Saccharomyces*, 135:597
Chlamydia trachomatis, vesicular sorting in *C. trachomatis*–infected cells, 134:363
Chlamydomonas
analysis of dynein regulatory mutations in, 135:1865
inner dynein arms but not outer dynein arms require activity of kinesin homologue protein KHP1^{FLA10} to reach distal part of flagella in, 133:371
Chlamydomonas reinhardtii, *PF*16 localizes to single microtubule of central apparatus in flagella of, encodes a protein with armadillo repeats, 132:359

Chloride ion, efflux of, regulatory role in polymorphonuclear leukocytes, induced by tumor necrosis factor, 135:511
Chloroplast protein import apparatus, 134:315
Chloroplast protein import intermediate, reveals distinct translocation machineries in two envelope membranes, 132:63
CHO cells, different biosynthetic transport routes to plasma membrane in BHK cells and, 133:247
Cholchicine, disrupts adhesion of full-length Po protein, 134:1531
Cholera toxin, in vero cells, transport of external Lys-Asp-Glu-Leu protein from plasma membrane to endoplasmic reticulum, KDEL, 133:777
Chondrocytes, bovine and human, effects of parathyroid hormone on collagen gene expression in, 135:1179
Chromaffin cells, catecholamine secretion requires translocation of p36 to subplasmalemmal region in, annexin II in exocytosis, 133:1217
Chromatin
  influence on anisotropy of microtubule asters, M-phase, 133:125
  organization of, lac operator-repressor recognition allows visualization of, 135:1685
Chromosome
  kinetochore forces acting on, 135:315
  structure of, dictates site-specific initiation of DNA replication, in *Xenopus*, 135:1207
  territories of, 3-dimensional arrangement of genes positioned peripherally in, gene expression is independent of position on, 135:1195
Chromosome alignment, molecular characterization of 50-kD subunit of dynactin reveals function for complex in spindle organization and, mitosis, 132:617
Chromosome territories, of active and inactive human X interphase, three-dimensional analysis of, 135:1427
Chromosome transport, force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but only at kinetochore during anaphase, 132:1093
Chrondrogenesis, chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 in vitro, gene expression of parathyroid hormone/PTH-related peptide receptor, 133:457
Chs2p, in *Saccharomyces*, temporal and spatial characterization of Chs3p and, 135:597
Chs3p, in *Saccharomyces*, temporal and spatial characterization of Chs2p and, 135:597
Chymase, alterations of phenotypes of, changes in mice during mastocytosis, 135:279
c-JunER protein, causes reversible loss of mammary epithelial cell polarity involving destabilization of adherens junctions, estrogen-dependent, 132:1115
Clathrin, single vesicle budding step involving dynamin and, synaptic vesicle cycle, 133:1237
Clathrin coat protein, AP-2 adaptor initiates assembly on dense lysosomes, 135:1801
Clathrin domain, functions of auxilin at, cooperation effects of J domain and, 135:925
Clathrin-coated pit formation, endocytosis of activated receptors and, 132:1025
Clathrin-coated vesicles
  budding from endosomes, 132:21
  effect of in vivo phosphorylation on, interactions of adaptors with, 135:635
Cleavage
  detection of protein kinase activity specifically activated at metaphase-anaphase transition, 132:635
  in living normal rat kidney cells, correlation of midzone microtubule bundles and TD60 with, 135:981
  regulated by cysteine protease, 135:479
Cleavage furrows
  calcium waves along, effects of heparin on, 135:181
  cellular blastoderm, *dah* involved in, 134:923
  in mutant embryos, disruption by novel mutant that arrests embryonic development during transition from syncytial to cellular blastoderm, 134:923
Clostridium botulinum, neurotoxin C1, causes growth cone collapse and inhibited exonal growth, 134:205
Clustering, telomere, in yeast, 134:1349
Coagulation pathway, mechanism for regulating cell surface proteolysis of tissue factor–dependent, ligand-induced protease receptor translocation into caveolae, 133:293
Coat and bud assembly, requirements of first stage of post-Golgi vesicle formation, role of PKC in, 135:355
Coat proteins, novel adaptor-related protein complex, 133:749
Coated pits
  cytoplasmic recognition feature for, 134:339
  endocytosis of activated receptors and clathrin-coated pit formation, 132:1025
Coatomer
  physical and functional interactions between seven COP subunits, 135:53
  role of p23 to bind, in formation of COPI vesicles on Golgi, 135:1239
Codon, CUD start, role in translational activation of FGF-2 in cell transformation and stress response, 135:1391
Cofilin
  along with actin-depolymerizing factor is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 132:871
  overexpression stimulates bundling of actin filaments, membrane ruffling, and cell movement in *Dictyostelium*, 132:335
Col11a2, tissue-specific expression of, 134:1573
Colchicine, reduction in synthesis and mRNA levels for α-tubulin by, 135:1525
Collagen
  α3(IV), molecular and functional consequences in mice kidneys deficient in, related to Alport syndrome, 135:1403
  (V), in chicken, role in regulation of corneal fibril diameter, 135:1415
  intracellular interaction of collagen-specific stress protein HSP47 specific for, newly synthesized procollagen, 133:469
  type XI, expression of, 134:1573
Collagen chains
  α1, effects of parathyroid hormone on, in bovine and human chondrocytes, 135:1179
  α1(VI), in transgenic mice, specific regions involved in regulation of transcription of, temporal and spatial expression patterns of, 135:1163
Collagen gene, proα2(I), regulation of high levels of expression of, by far-upstream enhancer, 134:1333
Collagen gene expression, effects of parathyroid hormone on, in bovine and human chondrocytes, 135:1179
Colon cancer
  familial and sporadic, adenomatous polyposis coli gene mutations linked to polyp formation, 134:165
  suppression of tumorigenicity by plakoglobin, augmenting effect of N-cadherin, 133:199
Colon carcinoma cells, early detachment of, during CD95(APO-1/Fas)-mediated apoptosis, 134:1089
Conditioned medium factor (CMF), regulates chemoattractant receptor cAR1 in *Dictyostelium*, 134:1543
Conjugation, of *Saccharomyces*, effects of Ste6 on a-factor in cell fusion during, 135:741
Connexin43 (Cx43), nonjunctional, plasma membrane hemichannels, 134:1019
Constitutive cells, biosynthetic regulated secretory pathway in secretory, 133:1177
Contractile potential, and smoothelin expression, 134:401
Contractility, Rho-stimulated, drives formation of stress fibers and focal adhesions, 133:1403
Convertase, subtilisin-like protein, expression during embryogenesis, 134:181
COOH-terminal, isoforms of proprotein convertase varying in, sort to different subcellular locations, 135:1261
COOH-terminal Glut4 epitope, unmasked by insulin, 135:415
COOH-terminal region, truncation mutants of, of fusion protein, importance in various stages of fusion, 135:73
COOH-terminal sequence, extreme, required for targeting of human catalase to yeast peroxisomes, 134:849
βCOP, involved in pH-dependent formation of transport vesicles destined for late endosomes, 133:29
δCOP, characterization of, interactions within coatomer complex, homologies to clathrin adaptor complexes, lethal knock-out of, 135:53
COP I vesicles, sorting by, under interphase and mitotic conditions, 134:1411
COPI, recruitment by COPII of, in transport mechanism from endoplasmic reticulum to Golgi, regulation by GTPases of, 135:895
COPI-coated vesicles, role of p23 to bind coatomer involved in formation of, 135:1239
COPII
  cargo export in endoplasmic reticulum budding zones containing, 135:19
  on endoplasmic reticulum, packaging amino acid permeases into transport vesicles by Shr3p and, 135:585
  recruitment of COPI by, in transport mechanism from endoplasmic reticulum to Golgi, regulation by GTPases of, 135:895
COPs, physical and functional interactions between coatomer subunits, 135:53
Cornea, role of collagen (V) in regulation of fibril diameter of the, 135:1415
Cortical cytoskeleton, regulation of orientation of centrosome separation by, 134:103
Cortical granule exocytosis, at fertilization in mouse eggs, function of Rabphilin-3A in, 135:1741

Cortical granules, have spare active fusion complexes randomly distributed as Poisson process, 134:329
Crossbridges, side-polar nonhelical arrangement of myosin filaments in, 134:53
Cryoelectron microscopy, of D1 and cytochrome b559 polypeptides, biochemical characterization, localization, two-dimensional crystals of photosystem II, 132:823
Crystals, two-dimensional of photosystem II, biochemical characterization, cryoelectron microscopy, localization of D1 and cytochrome b559 polypeptides, 132:823
CS-1, melanoma cells, transfection with expression vectors coding for integrins β3 or β5, 134:1323
cut6, impaired nuclear division in, 134:949
Cyclin B, cytoplasmic localization of in damage-arrested cells, 134:963
Cyclins, adhesion-dependent cell cycle progression linked to expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of retinoblastoma protein, 133:391
Cycloheximide, constitutive expression of machinery for programmed cell death, 133:1053
Cylindrotheca fusiformis, DSK1 is novel kinesin-related protein from diatom, anaphase spindle elongation, 133:595
Cysteine^{34}, of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 132:577
Cysteine protease, NEDD-2, link between processing and apoptotic activity of, cleavage events regulated by, 135:479
Cysteine residues, posttranslational acylation on, by integral membrane proteins and reversibly membrane-associated proteins in eukaryotic cells, 134:647
Cysteine-rich protein (CRP), of family of LIM proteins, required for terminal differentiation in vertebrate muscle development, 134:1179
Cytochalasin D, blocks aluminum fluoride–induced, ADP ribosylation factor–dependent, formation of protrusive structures, 134:935
Cytochrome b559, biochemical characterization, cryoelectron microscopy, and localization of polypeptide D1 and, two-dimensional crystals of photosystem II, 132:823
Cytokeratins, gene targeting at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 132:925
Cytokinesis
  actin-depolymerizing factor/cofilin is a phosphorylation-regulated protein essential for development, *Xenopus laevis*, 132:871
  involvement of Fyn tyrosine kinase in progression of, B lymphocyte progenitor, 132:91
  in living normal rat kidney cells, correlation of midzone microtubule bundles and TD60 with cleavage during, 135:981
  novel member of *rho* family of small GTP-binding proteins required for, 133:1321
  novel suppressor of *ras1* in fission yeast is dosage-dependent inhibitor of, *byr4*, 133:1307
Cytoplasm, identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 132:999
Cytoplasmic domain
  of integrin-associated protein (IAP), irrelevant for vitronectin bead binding, 134:1313
  internalization signal in, simian immunodeficiency virus transmembrane protein, modulates expression of envelope glycoproteins on cell surface, 132:795
  involved in Rer1p-independent retention in ER localization of Sec12p, 134:279
  synergy of integrin α subunit ratios, growth factors and, regulate muscle proliferation and differentiation, 133:169
Cytoplasmic tail
  cysteine^{34} of, cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 132:577
  β_{PS}, directs localization of CD2 to muscle termini in vivo, 134:217
  targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 132:565
Cytoskeleton
  cortical actin, β-catenin and fascin modulate dynamics of, 134:1271
  epidermal growth factor–induced alterations in, can be reproduced by displacement of select actin-modifying proteins from PIP2-bound state, 134:689
  plectin-mediated interaction of microtubules with intermediate filaments in, 135:991
  selective stabilization of Tau in axons and microtubule-associated protein 2C in cell bodies and dendrites contributes to polarized localization of proteins of, mature neurons, 132:667
  smoothelin novel protein in, 134:401
Cytosol, assembly of ER-associated protein degradation in vitro is dependent on calnexin, ATP, and, 132:291
Cytosolic transport, recombinant p58, p54, and p62 deplete activity, 134:589
Cytotoxin, oligomeric and subunit structure of *Helicobacter pylori* vacuolating, 133:801
dah mutant embryos, abnormal nuclear morphology in, correlates with cytoskeletal structure defects, 134:923
Dendrites
  selective stabilization of microtubule-associated protein 2C in cell bodies and, tau in axons, polarized localization of cytoskeletal proteins in mature neurons, 132:667
  striatin in spines, 134:1051
Dense-core secretory granules, targeting of membrane proteins to, 134:1229
Desmin, absence of, leads to muscle degeneration, 134:1255
Desmin null mice, disruption of muscle architecture and degeneration in, 134:1255
Desmoglein, mice expressing mutant desmosomal cadherin exhibit abnormalities in desmosomes, proliferation, and epidermal differentiation, 133:1367
Desmoplakin
  COOH-terminal domain of, associates with vimentin or keratin intermediate filaments, 134:985
  required for intermediate filament anchorage to desmosomes, 134:985
Desmosomal plaque, intermediate filaments linked by desmoplakin to, 134:985
Desmosomes
  binding of plakoglobin to cadherins in, patterns of binding sites and topogenic potential, 133:359
  envoplakin links cornified envelope to, 134:715
  localization of plakophilin 2 in, 135:1009
  mice expressing mutant cadherin in, exhibit abnormalities in desmosomes, proliferation, and epidermal differentiation, 133:1367
  plakoglobin as essential component of myocardiac, 135:215 Heart, role of plakoglobin in stability of tissue of, functions of desmosomes in, 135:215
  role of pinin in reinforcement of intermediate filament complex, 135:1027
  symplekin in, 134:1003
Detergent solubility
  in MDCK cells, correlation of Kv2.1 K^+ channel with, 135:1619
  traffic, polarity and, glycosylphosphatidylinositol–anchored protein after LDL-deprivation of MDCK cells, 133:1265
Developmental defect, of male *Plasmodium* gametocytes, identification and characterization of, 135:269
Diapedesis, monocyte, VLA-4 in, 134:1063
Diatom spindle kinesin 1, involved in anaphase spindle elongation, novel kinesin-related protein from diatom *Cylindrotheca fusiformis*, 133:595
Dictyostelium
  functions of 34k-Dalton actin-binding protein in, effects on filopodia, 135:965
  overlapping functions of myosin-I isoforms, 133:221
  regulation of cAR1 in, 134:1543
Dictyostelium, mutants lacking multiple classic myosin I isoforms reveal combinations of shared and distinct functions, 133:305
Dictyostelium discoideum
  myosin heavy chain kinase A in, regulates myosin localization during growth and development, 132:101
  myosin II heavy chain–specific protein kinase C isolated from, regulates myosin II assembly and localization in response to cAMP, 134:911
  overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and cell movement in, 132:335
Differentiation
  of basal cells in mice, *distal-less* homeodomain gene regulates, 135:1879
  hematopoietic and muscle, p53 involvement in, 134:193
Dihydrofolate reductase fusion proteins, import of into glycosomes in vivo not inhibited by folate-analogue aminopterin, 132:311
Dihydropyridine receptor, skeletal muscle, role in triad formation, 134:375
Dihydroxyacetone synthase (DHAS), Pmp47 and translocation and folding in, 134:37
Dilation, of fusion pore in HA-mediated cell fusion, 135:63
Di-leucine-based signals, protein targeting mediated by, distinct saturable components of targeting mechanisms of tyrosine-based and, 135:341
Dimerization
  of AGT, in peroxisome-to-mitochondrion mistargeting, 135:939
  heparin induces, confers proliferative activity onto hepatocyte growth factor antagonists NK1 and NK2, 133:709
  requirement for homophilic binding of C-cadherin, 135:487
Discontinuous actin hexagon (*dah*), new female sterile mutants with embryonic lethal phenotype, 134:923

Disintegrin, MDC9 is widely expressed cellular, contains cytoplasmic SH3 ligand domains, 132:717
Dissociation constant, determination of for adhesion receptors in contact area, visualization of CD2 interaction with LFA-3, 132:465
Distal-less homeodomain gene, Dlx3, in basal cells of mice, regulation of epidermal differentiation by, 135:1879
Dlg, Discs large protein of *Drosophila*, required for junction structure, cell polarity, and proliferation control in epithelia, 134:1469
DM-GRASP, heterophilic interactions of, GRASP-NgCAM interactions involved in neurite extension, 133:657
DNA
  interaction between TRF and mammalian telomeric, 135:867
  presence of breaks with single-base 3′ overhangs during apoptosis, 135:1369
DNA binding, p53, and transcriptional activities, both required to induce differentiation in p53-negative K562 cells, 134:193
DNA replication
  initiation of, chromosome structure dictates site-specific, in *Xenopus*, 135:1207
  nuclear proteins of quiescent *Xenopus laevis* cells inhibit, intact and permeabilized nuclei, 133:955
DNA sequences, method for localization and preservation of, 135:1685
DNA synthesis, induction of apoptosis and, cardiac myocytes by E1A oncoprotein, 133:325
DNA-damage checkpoint, Pds1p plays critical role in APC and checkpoint pathways, inhibitor of anaphase in budding yeast, 133:99
DNase I, hypersensitive sites in proα2(I) collagen gene, 134:1333
Drosophila
  actin organization, bristle morphology, and viability are affected by actin-capping protein mutations in, 133:1293
  cell type–specific roles for Cdc42, Rac, and RhoL in oogenesis in, 133:617
  Dlg protein of, required for junction structure, cell polarity, and proliferation control in epithelia, 134:1469
  embryogenesis, apodystrophin-like protein in, 134:923
  formation of actin filament bundles in ring canals of developing follicles of, 133:61
  formation and elongation of actin bundles in bristles of, 135:1291
  free centrosome duplication, in embryos, 134:103
  integrin localization in embryo, 134:217
  mechanism of *wingless* protein transmission from cell to cell in, role of glycosaminoglycans in WG signaling in, 135:819
  mitotic timing during postblastoderm development, controlled by dephosphorylation of inhibitory sites in CDC2, 134:963
  mutations in *hook* gene in, inhibit endocytosis of boss transmembrane ligand into multivesicular bodies, 133:1205
  oocytes, microtubule reorganization during meiotic spindle assembly in, 134:455
  paramyosin/miniparamyocin gene products, large diversity in quantity, localization, and isoform pattern, 134:81
  patterns of expression imply distinct functions for homologues of moesin and neurofibromatosis 2 tumor suppressor in, merlin, 133:843
  restricted temporal and spatial expression of muscle LIM proteins in, 134:1179
  roles of Cdc42 and Rac1 in formation of wing hairs in, 135:1277
  septin complex in, 134:1345
  septin complex forms filaments and exhibits GTPase activity, 133:605
  structure and function of flight muscle in tropomyosin mutants of, 135:673
  ZW10 protein of, 134:1127
DSK2, yeast ubiquitin-like genes involved in duplication of microtubule organizing center, 133:1331
Duchenne muscular dystrophy, 134:93, 134:873
Dynactin
  interaction between spectrin and centractin mediate associations of, 135:1815
  molecular characterization of 50-kD subunit of, reveals function for complex in chromosome alignment and spindle organization during mitosis, 132:617
  role in microtubule organization, 135:399
Dynamins
  association of dynamin-like protein with Golgi apparatus in mammalian cells, 133:761
  single vesicle budding step involving clathrin and, synaptic vesicle cycle, 133:1237
Dynein
  cell cycle regulation of, association with membranes modulates microtubule-based organelle transport, 133:585
  in *Chlamydomonas*, *sup-pf-2* mutations alter activity of outer arms of, 135:1865
  mammalian cells express three distinct heavy chains of, localized to different cytoplasmic organelles, 133:831
  role in microtubule organization, 135:399
Dynein arms, inner dynein arms but not outer dynein arms require activity of kinesin homologue protein KHP1^{FLA10} to reach distal part of flagella in *Chlamydomonas*, 133:371
γ-dynein heavy chain, in *Chlamydomonas*, *sup-pf-2* mutations alter activity of outer dynein arms by modifying, 135:1865
Dysgenesis, muscular, 134:375
Dystrophia muscularis mouse model, partial laminin α2 chain restoration in α2 chain-deficient *dy/dy* mouse by primary muscle cell culture transplantation, 133:185
Dystrophin
  deletion of actin-binding domain of, 134:873
  interaction of F-actin with, relationship with actin cross-linking superfamily, 135:661
  in skeletal muscle, links to dystrophin-associated protein (DAP) complex, by direct association with β-dystroglycan, 134:93
Dystrophin deletion constructs, forced expression, reveals structure–function correlations of COOH terminus, 134:93
E1 homotrimer, role in Semliki Forest virus fusion activity, 134:863
E1A proteins, induction of DNA synthesis and apoptosis in cardiac myocytes by E1A oncoprotein, 133:325
Early growth response–1 gene, UV activates growth factor receptors via reactive oxygen intermediates, 133:211
eat-5 and *unc-7*, multigene family in *Caenorhabditis elegans*, 134:537
E-cadherin–mediated adhesion, kinetic analysis, 134:549
EF1α, actin polymerization effected by, role in altering assembly of F-actin, 135:1309
EF-1α, effect of F-actin on binding of aminoacyl-tRNA to, pH sensitivity of, 135:953
Eg5, role in microtubule organization, 135:399
EGF, multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 132:1011
Egg jelly, sea urchin sperm receptor for, modular protein with extensive homology to human polycystic kidney disease protein, PKD1, 133:809
Elastic behavior, of α-satellite DNA domain of centromeres, 135:545
Elongation factor 1 α, actin polymerization effected by, role in altering assembly of F-actin, 135:1309
Embryogenesis
  expression of *SM22α* promoter in transgenic mice, distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, 132:849
  Fas and Fas ligand in embryos and adult mice, ligand expression in immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover, 133:335
  SPC4, SPC6, and SPC7, coexpressed with bone morphogenetic proteins at distinct sites during, 134:181
Embryonic development, compromised by reduced Armadillo function, 134:133
Embryonic stem cells, Bcl-2 inhibits retinoic acid–induced apoptosis during neural differentiation of, 132:955
Embryos, cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in, mouse, first mitotic division, 132:915
emo-1, homologue of *Caenorhabditis elegans* Sec61p γ, necessary for oogenesis and coupling of ovulation with meiotic maturation, 134:699
end4 mutation, blocks constitutive endocytosis, but not ubiquitination, of α-factor receptor (Ste3p), 134:661
Endocrine cells, tyrosine phosphatase and tyrosine kinase are required for formation of secretory vesicles from TGN of, 135:1471
Endocytosis
  of activated receptors and clathrin-coated pit formation, 132:1025
  of GPI-anchored proteins in human lymphocytes, role of glycolipid-based domains, actin cytoskeleton, and protein kinases, 133:791
  of GPI-linked membrane folate receptor-α, 132:35
  internalization signal in simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on cell surface, 132:795
  kinetics of, unaltered in synapsin I knockout mice, 134:1219
  mutations in *Drosophila hook* gene inhibit, boss transmembrane ligand into multivesicular bodies, 133:1205
  novel genetic screen identified *dim* mutants of, role for She4p and Pan1p in, 135:1485
  potential sites of PI-3 kinase function in, revealed by PI-3 kinase inhibitor, wortmannin, 132:595

in *Saccharomyces*, signaled by clathrin-dependent NPFXD sequence, distinct from Golgi localization sequence, 135: 1789
ubiquitin addition plays initiating role in, 134:661
of VAMP facilitated by synaptic vesicle targeting signal, 132:537
Endoglin, modulates cellular responses to TGF-β1, 133:1109
Endoplasmic reticulum
activation of transcription factor NF-κB by adenovirus E3/19K protein requires retention in, 132:511
assembly of ER to Golgi SNARE complex requires Uso1p, 132:755
calreticulin regulation of vinculin expression controls cell adhesiveness in, 135: 1913
COPII recruitment of COPI in transport mechanism to Golgi from, 135:895
ER membrane protein complex required for nuclear fusion, 132:499
export of cargo through COPII-coated vesicular carriers, organization of export complexes, 135:19
roles of Shr3p and COPII in packaging amino acid permeases into transport vesicles on, 135:585
storage disease causing congenital goiter and hypothyroidism, 133:517
topology of sphingolipid galactosyltransferases in Golgi and, transbilayer movement of monohexosyl sphingolipids required for higher glycosphingolipid biosynthesis, 133:15
transport of external Lys-Asp-Glu-Leu protein from plasma membrane to, KDEL, cholera toxin in vero cells, 133: 777
Endoplasmic reticulum–associated protein degradation, assembly in vitro is dependent on cytosol, calnexin, and ATP, 132:291
Endosomes
containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 132:1011
endosomal βCOP involved in pH-dependent formation of transport vesicles destined for late, 133:29
isoforms of invariant chain regulate transport of MHC class II molecules to antigen processing compartments, 133:281
of MDCK cells, apical and basolateral, polarized sorting mechanism of, 135:139
novel class of clathrin-coated vesicles budding from, 132:21
Endosomes, potential sites of PI-3 kinase function in endocytic pathway revealed by PI-3 kinase inhibitor, wortmannin, 132:595
role of specific domain of transferrin receptor in determining physical characteristics of, 135:1749
synaptic vesicle cycle, single vesicle budding step involving clathrin and dynamin, 133:1237
Endothelial cells
cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular, fibrin matrix, role of urokinase activity, 132: 1177
leukocyte adhesion to vascular, induces E-selectin linkage to actin cytoskeleton, 133:445
polymorphonuclear leukocyte adhesion triggers disorganization of adherens junctions of, 135:497
Entactin, in mammary epithelial cells in mice, relationship of apoptosis and matrix metalloproteinase activity to, 135: 1669
Envelope
chloroplast, import machinery, 134:315
cornified, 134:210-kD novel precursor of, 134:715
nuclear, MHC-B involved in breakdown of, 134:675
Envelope membranes, chloroplast protein import intermediate reveals distinct translocation machineries in, 132:63
Envoplakin, homology to desmoplakin, 134: 715
Eph subfamily, spatial and temporal expression of Cek9 of, 135:781
Epidermal growth factor
population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, TGFα, and, chemically defined (HGM) medium, 132:1133
regulation of intracellular functions of $\alpha_6\beta_4$ integrin by, 134:241
Epidermis
in mice, *distal-less* homeodomain gene regulates differentiation of, 135:1879
mice expressing mutant desmosomal cadherin exhibit abnormalities in desmosomes, proliferation, and differentiation in, 133:1367
severe skin fragility and changes of cytokeratin expression in, gene targeting at mouse cytokeratin 10 locus, 132:925
Epithelial cells
CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal, 132:437
estrogen-dependent c-JunER protein causes reversible loss of polarity of, mammary, destabilization of adherens junctions, 132:1115
heparan sulfate expression in polarized, apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, 132:487
human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 132:1189
inactivation of integrin β6 subunit gene reveals role of integrins in, regulating inflammation in lungs and skin, 133:921
motogenic and morphogenic activity of receptor tyrosine kinases in, 133:1095
14-3-3 proteins associate with phosphorylated simple epithelial keratins during cell cycle progression and act as solubility cofactor, 133:345
re-epithelialization after skin injury correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 132:381
sorting of newly synthesized galactosphingolipids to two surface domains of, 132: 813
ER membrane, TRAM required for protease-resistant insertion of nascent chains into translocation site, 134:25
ERM proteins, interaction between CD44 and effect of phosphatidylinositol on, regulation mechanism and involvement of Rho in interaction between CD44 and, 135:37
Escherichia coli, native 3D structure of eukaryotic 80S ribosome, morphological homology with 70S ribosome of, 133:495
E-selectin
fucosyltransferase FucT-VII regulates synthesis of ligand of, human T cells, 133:911
leukocyte adhesion to vascular endothelium induces linkage of, actin cytoskeleton, 133:445
Eukaryotes
roles of molecular chaperones in protein degradation, 132:255
septins and assembly of neck filaments in, 134:1345
Excitation–contraction coupling, ryanodine receptors in, 134:885
Exocytosis
actin-binding motif is unique to epsilon isoform of protein kinase C, participates in regulation of synaptic function, 132:77
annexin II in, catecholamine secretion requires translocation of p36 to subplasmalemmal region in chromaffin cells, 133:1217
calcium-triggered, rate and extent controlled by Poisson-distributed active fusion complexes, 134:329
of cortical granule, at fertilization in mouse eggs, function of Rabphilin-3A in, 135:1741
formation and kinetics of fusion pore induced by baculovirus GP64 during, 135:1831
Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues, 132:167
Explants, smooth muscle tissue, halt transcription of smoothelin mRNA, 134:401
Export, nuclear protein signals, 134:1157
Export complex, organization on endoplasmic reticulum, COPII-containing budding zones, 135:19
Expression
of α1(VI) collagen chain, in transgenic mice, temporal and spatial patterns of, 135:1163
of genes, independent of position in chromosome territories, 135:1195
Extracellular domain, role of CD3γ in T cell receptor assembly, 132:299
Extracellular matrix, alters PDGF regulation of fibroblast integrins, 132:239
Ezrin, interactions between protein 4.1 and, in PDZ and I3 domains of hDlg, 135: 1125
Ezrin-radixin-moesin family, patterns of expression imply distinct functions for *Drosophila* homologues of moesin and neurofibromatosis 2 tumor suppressor, merlin, 133:843
F-actin
effect on binding of EF-1α to aminoacyl-tRNA, pH sensitivity of, 135:953
effect of EF1α to alter assembly of, 135:1309
in filopodial motility, 134:1197
interaction of dystrophin with, 135:661
FAK, interaction domain on paxillin of, 135: 1109
Fas, and Fas ligand in embryos and adult mice, ligand expression in immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover, 133:335
Fascin, association with β-catenin, in non-cadherin complex, 134:1271
Fatty acid deficiency, and cell division, 134: 949
FBR v-fos, myristylation of, cell-specific differentiation in osteosarcoma caused by, 135:457

Fcγ receptors
for monomeric IgE, endocytosis of activated receptors and clathrin-coated pit formation, 132:1025
role of tyrosine kinases and, Na^+/H^+ exchange activity during phagocytosis in human neutrophils, 132:1037
FcεRI, aggregation of, initiates signaling pathways leading to exocytosis of inflammatory mediators of allergic response, 134:1427
Fetal alcohol syndrome, alcohol inhibits cell–cell adhesion mediated by human L1, 133:381
FGF-2
increases association of FGFR-1 immunoreactivity with nucleus time- and dose-dependently, 134:529
role of CUG start codon in translational activation of, in cell transformation and stress response, 135:1391
Fibril formation, RGD sequence required for initiation of, 134:573
Fibrillogenesis, in chicken cornea, role of collagen (V) in regulation of fibril diameter during, 135:1415
Fibrin, cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular endothelial cells in matrix of, role of urokinase activity, 132:1177
Fibroblast growth factor receptor-1
nuclear, role in regulation of gene activity, 134:529
stimulation of occupancy of, signaling by cell surface–associated syndecans and glypican, 133:405
Fibroblast growth factor receptors
nuclear translocation of, 134:529
signaling pathway, in human endothelial cells, 134:783
Fibroblast growth factors, regulate skeletal muscle development through autocrine and paracrine mechanisms, differentially expressed, 132:1151
Fibroblasts
electroporated with α-smooth muscle actin–specific mAb, exhibit increased motility, 134:67
extracellular matrix alters PDGF regulation of integrins in, 132:239
pp125^{FAK} functions to suppress apoptosis in, 135:1383
role of Rab8 in, reorganization of actin and microtubules in, 135:153
α-smooth muscle actin restrains migratory behavior in, 134:67
stimulation of β_1 integrins on, induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 132:741
Fibronectin
effect of CC chemokines on monocyte adhesion to, 134:1063
β1 integrin–dependent and –independent polymerization of, 132:227
matrix assembly, 134:573
novel role for integrin-binding III-10 module in matrix assembly of, 133:431
Filaments
Drosophila septin complex forms, GTPase activity, 133:605
subunit composition of neurofilaments specifies axonal diameter, 133:1061
Filensin, and phakinin form beaded intermediate filaments and coassemble de novo in cultured cells, 132:643
Filipin, altered distribution of folate receptors in cells treated with, 134:1169
Filopodia
in *Dictyostelium*, effects of 34k-Dalton actin-binding protein on, 135:965
motility of, 134:1197
Fission yeast mutants, aberrant mitosis in, 134:949
Flagella
inner dynein arms but not outer dynein arms require activity of kinesin homologue protein KHP1^{FLA10} to reach distal part of, *Chlamydomonas*, 133:371
*PF*16 localizes to single microtubule of central apparatus in, *Chlamydomonas*, encodes a protein with armadillo repeats, 132:359
Flight muscles
in *Drosophila* tropomyosin mutants, structure and function of, 135:673
indirect, and stretch-activation, 134:81
Fluorescence, monitoring kinetics of cell–cell fusion events, 135:63
Fluorescence in situ hybridization (FISH), detects changes in telomere size at single cell level, 134:1
Focal adhesion
paxillin targeting motif for, 135:1109
pp125^{FAK} functions to suppress apoptosis in fibroblasts by regulating, 135:1383
Focal adhesion kinase
colocalization with αvβ5, independent of protein kinase C, 134:1323
controls adhesion-dependent cell survival, 134:793
phosphorylation of, dependent on protein kinase C, 134:1323
Focal adhesion sites, peripheral association of, in cell migration, 134:783
Focal adhesions, Rho-stimulated contractility drives formation of stress fibers and, 133:1403
Folate receptors, clustered, and 5-methyltetrahydrofolate delivery, 134:1169
Folate-analogue aminopterin, import of DHFR hybrid protein into glycosomes in vivo not inhibited by, 132:311
Follicles, formation of actin filament bundles in ring canals of developing, *Drosophila*, 133:61
Fucosyltransferase, FucT-VII regulates E-selectin ligand synthesis in human T cells, 133:911
Fusion
with lysosomal protease, strategy for engineering dominant negative mutations using, 135:1043
of membranes mediated by influenza virus hemagglutinin requires action of at least three hemagglutinin trimers, 133:559
Fusion complexes, active, Poisson-distributed, in exocytosis, 134:329
Fusion defective mutants, in yeast, characterization of, reduced levels of a-factor mating pheromone results in high levels of, 135:1727
Fusion pore
formation and kinetics of, induced by baculovirus GP64, 135:1831
mediated by HA-RBC fusion, fluorescence monitoring of, stages of dilation of, 135:63
Fusion protein, COOH-terminal region of, importance in various stages of fusion, 135:73
Fyn tyrosine kinase, involved in progression of cytokinesis of B lymphocyte progenitor, 132:91
G1 phase, Taxol-dependent arrest of, effect on transformed and nontransformed cells, 135:689
G proteins
actin cytoskeleton-mediated translocation of Src kinase to cell periphery controlled by, 135:1551
G-protein-coupled receptor, ventricular zone gene-1, in neurogenic regions of cerebral cortex, role in LPA signaling, 135:1071
link cell density–sensing conditioned medium factor (CMF) receptor and chemoattractant cAMP receptor, 134:1543
parallels between molecular motors and, 135:291
G91A mutant, of E1 fusion peptide, causes acid shift in pH threshold of cell–cell fusion, 134:863
G91D mutant, of E1 fusion peptide, blocks spike protein membrane fusion activity, 134:863
Gag precursor, of Moloney murine leukemia virus, targeting to virus budding site of, 135:1841
Galactosphingolipids, sorting of newly synthesized, two surface domains of epithelial cells, 132:813
Galactosyltransferase
BED1 is gene encoding homologue of, required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*, 132:137
β1,4, resident Golgi marker protein, protein kinase C μ colocalizes with, 134:1401
Galectin-1, new pathway for protein export in *Saccharomyces cerevisiae*, 133:1017
Gal-NAcPTase–proteoglycan interaction, role in morphogenetic cell and tissue interactions during development, 134:801
Gametocytes, of *Plasmodium*, identification and characterization of developmental defect of, 135:269
Gap junction proteins, role of *Caenorhabditis elegans* gene family in, 134:537
Gap junctional hemichannels, in plasma membranes of cultured cells, 134:1019
Gastrulation
embryonic defects in, from reduced Armadillo function, 134:133
position-specific activation of RGD/synergy site–dependent migratory behavior in, 134:227
Gelsolin
and capping protein, coordinated regulation of platelet actin filament barbed ends by, 134:389
role in actuating epidermal growth factor receptor–mediated cell motility, 134:689
Gene targeting, at mouse cytokeratin 10 locus, severe skin fragility and changes of cytokeratin expression in epidermis, 132:925
Gene trapping, in differentiating cell lines, 134:837
Genes, 3-dimensional arrangement of, in periphery of chromosome territories, independent of gene expression, 135:1195
Genetic screen, yeast endocytosis *dim* mutants identified through novel, 135:1485
Genome integrity, mitochondrial, Hsp necessary for maintenance of under severe temperature stress, 134:1375
GLAST-1, in central nervous system, membrane topology of, 135:1867
Glial fibrillary acidic protein, disrupted network of, astrocytes from vimentin knockout mice, 133:853

Glucocorticoid receptor, fusion of chimera with green fluorescent protein, evidence that RAN/TC4 GTPase mediates essential function independent of nuclear protein import, 133:985
Glucose transport, in adipocytes, insulin stimulates, by translocation of glucose transporter from intracellular site to cell surface, 134:625
Glucose transporter (GLUT-4), segregated from recycling endosomes in insulin-sensitive cells, 134:625
Glut4, in skeletal muscle, insulin-stimulated transport mediated by, localized in transverse tubules, 135:415
Glutamate receptors, role in migration deficiency of weaver granule neurons, 134:477
L-glutamate transporter, in central nervous system, membrane topology of, 135:1867
Glutathione peroxidase, inhibition of IκB-α phosphorylation and degradation and subsequent NF-κB activation by overexpression of, 133:1083
Glutathione-*S*-transferase, fusion to complete intracellular domain of receptor-like protein tyrosine phosphatase (RPTPμ), 134:1513
Glycine 699, alanine mutagenesis of, alters motor activity of skeletal muscle myosin, 134:895
Glycoaminoglycan chains, [^{35}S]SO_4-labeled, biosynthetic regulated secretory pathway in constitutive secretory cells, 133:1177
Glycolipid-enriched membrane domains, Lck and lymphoid-specific tyrosine kinase associated with, relationship between Lck and exclusion of CD45 from, 135:1515
Glycolipids
  antiglycolipid antibody inhibits Madin-Darby canine kidney cell adhesion to laminin, interferes with basolateral polarization and tight junction formation, 133:695
  role of actin cytoskeleton, protein kinases and domains based on, endocytosis of GPI-anchored proteins in human lymphocytes, 133:791
Glycoproteins
  internalization signal in simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope, cell surface, 132:795
  processing of, not disrupted by *Chlamydia trachomatis*, 134:363
  sorting of, in chlamydia-infected cells, 134:363
Glycosaminoglycans, role in *Drosophila wingless* protein transmission, 135:819
Glycosomes, import of DHFR hybrid protein into, in vivo, not inhibited by folate-analogue aminopterin, 132:311
Glycosphingolipid biosynthesis, transbilayer movement of monohexosyl sphingolipids required for, topology of sphingolipid galactosyltransferases in ER and Golgi, 133:15
Glycosylation, of CD44 implicated in CD44-mediated cell adhesion to hyaluronan, 132:1199
Glycosylphosphatidylinositol, traffic, polarity, and detergent solubility of protein anchored with, LDL-deprivation of MDCK cells, 133:1265
Glycosylphosphatidylinositol (GPI)-anchored proteins, in caveolae, clustering of, depends on lipid–lipid interactions, 134:1169
Glycosyl-phosphatidyl-inositol–linked membrane folate receptor-α, endocytosis of, 132:35
Glypican
  apical sorting of, GPI-anchored proteoglycan, inversely related to heparan sulfate content, heparan sulfate expression in polarized epithelial cells, 132:487
  stimulation of fibroblast growth factor receptor-1 occupancy and signaling by cell surface–associated syndecans and, 133:405
Goiter, endoplasmic reticulum storage disease causing hypothyroidism and congenital, 133:517
Golgi
  association of dynamin-like protein with, mammalian cells, 133:761
  Cab45 is novel Ca^{2+}-binding protein localized to lumen of, 133:257
  COPII recruitment of COPI in transport mechanism from endoplasmic reticulum to, 135:895
  distribution of α subunits and βγ subunits of heterotrimeric G proteins on membranes of, exocrine pancreas, 133:1027
  with high basal phospholipase D activity, do not require exogenous ADP ribosylation factor (ARF) for production of coatomer-coated vesicles, 134:295
  identification of small cytoplasmic ankyrin in kidney and muscle that binds βIΣ* spectrin and associates with, Ank_{G119}, 133:819
  ion channel activity of influenza virus M_2 protein affects transport through, 133:733
  localization sequence of, new clathrin-dependent endocytosis signal in *Saccharomyces* distinct from, 135:1789
  mammalian cells express three distinct dynein heavy chains that are localized to different cytoplasmic organelles, 133:831
  p23 receptor binds coatomer involved in formation of COPI vesicles on, 135:1239
  topology of sphingolipid galactosyltransferases in ER and, transbilayer movement of monohexosyl sphingolipids required for higher glycosphingolipid biosynthesis, 133:15
  transport of external Lys-Asp-Glu-Leu protein from plasma membrane to endoplasmic reticulum, KDEL, cholera toxin in vero cells, 133:777
  Vps10p cycles between late-Golgi and prevacuolar compartments in its function as sorting receptor for multiple yeast vacuolar hydrolases, 133:529
  v-SNARE implicated in intra-Golgi transport, 133:507
Golgi apparatus
  assembly of ER to Golgi SNARE complex requires Uso1p, 132:755
  localization of yeast early Golgi mannosyltransferase involves retrograde transport, Och1p, 132:985
Golgi compartment, protein kinase C μ, located in, 134:1401
Golgi complex, pH of, 134:1387
Golgi membranes, targeting and fusion of ER-derived transport vesicles with purified, yeast, 132:277
Golgi vesicle formation, phospholipase D in, 134:295
GOS-28, v-SNARE implicated in intra-Golgi transport, 133:507
gp75, neurotrophin receptor, forms complex with receptor tyrosine kinase TrkA, 132:945
GPI-anchored proteins, endocytosis of, human lymphocytes, role of glycolipid-based domains, actin cytoskeleton, and protein kinases, 133:791
Granule lattice protein 1, in *Tetrahymena*, function and composition of secretory granules effected by, 135:1775
Granulocyte macrophage colony stimulating factor, produced in lesioned peripheral nerves induces up-regulation of cell surface expression of MAC-2 by macrophages and Schwann cells, 133:159
Green fluorescent protein, fusion with glucocorticoid receptor chimera, evidence that RAN/TC4 GTPase mediates essential function independent of nuclear protein import, 133:985
Growth cone, motile behaviors, in response to micro-CALI of talin and vinculin, 134:1197
Growth cone collapse, neurotoxin C1 of *Clostridium botulinum* cause of, 134:205
Growth factor receptors, UV activates, via reactive oxygen intermediates, 133:211
Growth factors
  mechanisms of synergy between integrins and, MAP kinase activation by, phosphorylation triggered by, 135:1633
  synergy of integrin α subunit ratios, cytoplasmic domains and, regulate muscle proliferation and differentiation, 133:169
GTPase
  COPII recruitment of COPI in transport mechanism from endoplasmic reticulum to Golgi regulated by, 135:895
  distinct from Ran involved in nuclear protein import, 133:971
  Drosophila septin complex forms filaments and exhibits activity of, 133:605
  evidence that RAN/TC4 GTPase mediates essential function independent of nuclear protein import, green fluorescent protein-glucocorticoid receptor chimera, 133:985
GTP-binding proteins, novel member of *rho* family of small, required for cytokinesis, 133:1321
GTPγS, biochemically distinct steps revealed with NEM, BAPTA, and, assembly of nuclear pore, 132:5
GUK, conserved domain of Discs large (Dlg) protein, in regulation and maintenance of septate junctions, 134:1469
Haemanthus, force for poleward chromosome motion in cells of, acts along length of chromosome during metaphase but only at kinetochore during anaphase, 132:1093
Hair, on wings of *Drosophila*, roles of Cdc42 and Rac1 in formation of, 135:1277
hDlg
  in PDZ domains, interactions of ATP and protein 4.1 with, 135:753
  protein 4.1 interactions with PDZ and I3 domains of, 135:1125
Healing of wounds, in chick embryo, actin purse-string assembly during, roles of cadherin, myosin and Rho during, 135:1097
Heat shock protein 47, intracellular interaction of collagen-specific stress protein, newly synthesized procollagen, 133:469
Heat shock proteins, refolding activity of Ssa1/2p in yeast cytosol, 135:1229

Heat-sensitive mutant, troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*, 132:1061
Helicobacter pylori, oligomeric and subunit structure of vacuolating cytotoxin in, 133:801
Helix initiation peptides, 1A, disrupt vimentin intermediate filaments in vitro and in vivo, 134:971
Hemagglutinin
escapes endocytosis because of property residing in its transmembrane sequences, 134:339
ion channel activity of influenza virus M_2 protein affects transport through Golgi apparatus, 133:733
membrane fusion mediated by, influenza virus, requires action of at least three hemagglutinin trimers, 133:559
wild type, exclusion from coated pit, 134:339
Hemagglutinin antigen α1-tubulin, isolation and characterization in hamster, repression of synthesis of endogenous α-tubulin caused by overexpression of, 135:1525
HEMCAM, characterization of adhesion molecule, expressed by hemopoietic progenitors, 135:1655
Hemidesmosomes, β4 integrin required for formation of, 134:559
Hemopoietic progenitors, characterization of adhesion molecule HEMCAM expressed by, 135:1655
Heparan sulfate
expression in polarized epithelial cells, apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, 132:487
stimulation of fibroblast growth factor receptor-1 occupancy and signaling by cell surface–associated syndecans and glypican, 133:405
Heparin
calcium waves along cleavage furrows inhibited by, 135:181
induces dimerization and confers proliferative activity onto hepatocyte growth factor antagonists NK1 and NK2, 133:709
tenascin-Y binding to, 134:1499
Hepatocyte growth factor, heparin induces dimerization and confers proliferative activity onto antagonists NK1 and NK2 of, 133:709
Hepatocyte growth factor/scatter factor, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by EGF, TGFα, and, chemically defined (HGM) medium, 132:1133
Hepatocytes, population expansion, clonal growth, and specific differentiation patterns in primary cultures of, induced by HGF/SF, EGF, and TGFα in chemically defined (HGM) medium, 132:1133
HepG2 cells, express endogenous protein kinase μ (PKCμ), 134:1401
Heptanol, in gap junctional communication, 134:1019
Heregulin, signaling, at neuromuscular synapse, 134:465
Herpes simplex virus (HSV)-1, replication compartments of, located at periphery of nuclear domain (ND)10, 134:815
Heterotrimeric G proteins, distribution of α subunits and βγ subunits of, Golgi membranes of exocrine pancreas, 133:1027
High mobility group box, identification of two Sox17 messenger RNA isoforms with and without region of, differential expression in mouse spermatogenesis, 133:667
Histone, δ, phosphorylation of, occurs early in *Tetrahymena* conjugation when micronuclei become transcriptionally active, 135:1219
HIV-1 RNA, nuclear export facilitated by Rev protein, intranuclear distribution and spatial relationships of, 135:9
Homodimers, formed by Pex11p, formed by oxidative metabolism within maturing peroxisomes, 135:123
hook gene, mutations in, *Drosophila*, inhibit endocytosis of boss transmembrane ligand into multivesicular bodies, 133:1205
Hox, regulatory relationship with *Col11a2*, 134:1573
hsc70, effects in J and clathrin domains of auxilin and, 135:925
Hsp70, activity during targeting and translocation phases of nuclear transport, effect on NLS, 135:329
Hsp78, necessary for reactivation of mitochondrial protein synthesis after heat stress, 134:1375
HSY cells, calcium release in, conforms to steady-state mechanism involving regulation of inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$, 132:607
Human, multidrug resistance 3–P-glycoprotein of, expression in transgenic mice induces lens membrane alterations leading to cataract, 132:701
Human diploid fibroblasts, interphase, telomeres distributed in interchromosomal size heterogeneity throughout nucleus, 134:1
Human fibroblast growth factor 2, role of CUG start codon in translational activation of, in cell transformation and stress response, 135:1391
Human mitochondrial import receptor, hTom20p, excludes cryptic matrix-targeting signal from mammalian mitochondria, 134:307
Human polo-like kinase1, maturation of mitotic centrosomes and spindle formation requires, 135:1701
Human umbilical vein endothelial cells, senescent, FGF-1–dependent proliferative and migratory responses impaired in, 134:783
Human X interphase chromosome, territories of, three-dimensional analysis of active and inactive, 135:1427
Hyaluronan
CD44 binding to, mediates rolling interactions of lymphocytes on tonsillar stroma, 134:1075
glycosylation of CD44 implicated in CD44-mediated cell adhesion to, 132:1199
Hyaluronic acid, variants of CD44 effect binding properties of, 135:1139
Hydrolases, Vps10p cycles between late-Golgi and prevacuolar compartments in its function as sorting receptor for multiple yeast vacuolar, 133:529
Hydrophobic regions, along surface of myosin heavy chain rods, characteristic variations in, 135:371
Hydrophobicity, of signal sequence hydrophobic core, correlates with pathway use, 134:269
Hypothyroidism
endoplasmic reticulum storage disease causing congenital goiter and, 133:517
neurotrophins promote survival and development of neurons in cerebellum of rats with, in vivo, 133:631
I band, three-dimensional structure of Z band in normal mammalian skeletal muscle, 133:571
I3 domain, of hDlg, interaction between protein 4.1 and, 135:1125
IAP75, outer envelope import machinery component, cross-links to transit sequence of precursor to small subunit of ribulose-1,5-bisphosphate carboxylase, 134:315
IAP86, outer envelope import machinery component, cross-links to transit sequence of precursor to small subunit of ribulose-1,5-bisphosphate carboxylase, 134:315
IκB-α, inhibition of phosphorylation of, degradation and subsequent NF-κB activation by glutathione peroxidase overexpression, 133:1083
ICE family proteases, inhibits metabolic and genetic events associated with neuronal cell death, functional comparison between ICE family proteases and, 135:1341
I-cell disease
biogenesis of MHC class II compartment in B lymphoblasts of, 132:769
specific lysosomal deficiency of cathepsin B contributes to myopathy of, 134:837
ICE-related cysteine protease, roles in anoikis of Jun-N-terminal kinases and, 135:1377
ICRF-193, topoisomerase II α inhibitor, prevents its normal accumulation at centromere, 134:1097
Ig fusion proteins, soluble, and responsiveness of pre-B cells to IL7, 134:771
Ig superfamily, interaction between neurofascin and NrCAM of, 135:1059
IgE-FcεRI, capacity of ligand-coated, 6-μm diameter beads to bind, 134:1427
III-10 Module, novel role for integrin-binding, fibronectin matrix assembly, 133:431
Immature secretory granules, AP-1 adaptor complex binds to, PC12 cells, regulated by ADP-ribosylation factor, 132:523
Immune-receptor tyrosine-based activation motif, features of $p59^{fyn}$ src homology 4 domain define motif for binding of, plasma membrane localization, 133:1007
Immunoglobulin A, signal transduction by polymeric immunoglobulin receptor suggests role in regulation of receptor transcytosis, 133:997
Immunoglobulin domains, gene *unc-89* encodes giant modular protein composed of domains of signal transduction and, *Caenorhabditus elegans*, required for muscle M-line assembly, 132:835
Import
nuclear protein, role of Ran binding protein (RanBP1) in, 134:1157
protein, across chloroplast envelope, 134:315
Importin-β, dynamic localization of nuclear import receptor and its interactions with transport factors, 133:1163
Inclusion membrane, chlamydial, dissociated from endocytic pathway, 134:363
Inflammatory diseases
CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia, 132:437

inactivation of integrin β6 subunit gene reveals role of epithelial integrins in regulating, lungs and skin, 133:921
Influenza hemagglutinin, kinetics of cell fusion between RBC and, fusion pore dilation mediated by, fluorescence monitoring of, 135:63
Influenza virus
chimeric hemagglutinin proteins, endocytosis of, 134:339
ion channel activity of M_2 protein of, transport through Golgi apparatus, 133:733
membrane fusion mediated by hemagglutinin of, requires action of at least three hemagglutinin trimers, 133:559
Inheritance, in *Saccharomyces*, function of myosin and actin in rapid and directed vacuole movement during cell division, 135:1535
Inositol 1,4,5-trisphosphate, calcium release in HSY cells conforms to steady-state mechanism involving regulation of, receptor Ca^{2+} channel, luminal $[Ca^{2+}]$, 132:607
Insulin
in skeletal muscle, Glut4 mediates transport stimulated by, surface area of transverse tubules increased by, COOH-terminal antigenic epitope unmasked by, 135:415
stimulation of glucose transport in adipocytes by, 134:625
Insulin-like growth factor binding protein-5, modulates muscle differentiation through insulin-like growth factor–dependent mechanism, 133:683
Insulin-like growth factor I, events leading to stimulation of myogenesis by, activates proliferation prior to differentiation, 135:431
Insulin-like growth factors, insulin-like growth factor binding protein-5 modulates muscle differentiation through mechanism dependent on, 133:683
Integral membrane proteins
p56, is lamin B receptor-like, mediates membrane targeting to chromatin in sea urchins, 135:1715
S-acylation of, on cysteine residues near cytoplasmic termini of transmembrane helixes, or on cytoplasmic cysteine residues distant from helix, 134:647
sorting sequence of peroxisomal, PMP47 is contained within short hydrophilic loop, 133:269
Integrin
α6β4, anchorage mediated by, to laminin 5 (epiligrin), regulates tyrosine phosphorylation of membrane-associated protein, 132:727
β1D, displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 132:211
αIIbβ3, ectopically expressed, migration of CHO cells on fibrinogen mediated by, 134:1551
α_4, functions in mouse skeletal muscle of, 135:829
high affinity, dominant effect in inhibiting migration, 134:1551
$\alpha_v\beta_3$, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for, 132:475
β4, in junctional epidermolysis bullosa, 134:559
α5β1, novel role for integrin-binding III-10 module in fibronectin matrix assembly, 133:431
$\alpha_6\beta_4$, regulation by epidermal growth factor, 134:241
regulation by platelet-derived growth factor, 134:1301
β_1, stimulation of, fibroblasts, induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 132:741
Integrin localization, directed by intracellular signals to sites of function in embryonic muscles, 134:217
Integrin α subunit ratios, cytoplasmic domains, growth factor synergy regulate muscle proliferation and differentiation, 133:169
Integrin-associated protein
effects of thrombospondin on vitronectin through, 135:533
IgV domain, required for vitronectin bead binding, 134:1313
Integrins
cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 132:1161
enhance FcεRI-mediated responses in adherent RBL-2H3 cells, 134:1427
extracellular matrix alters PDGF regulation of, fibroblast, 132:239
inactivation of integrin β6 subunit gene reveals role of epithelial, regulating inflammation in lungs and skin, 133:921
β1 integrin–dependent and –independent polymerization of fibronectin, 132:227
interaction with extracellular matrix proteins, activates focal adhesion kinase, 134:793
mechanisms of synergy between growth factors and, MAP kinase activation by, tyrosine phosphorylation triggered by, 135:1633
Intermediate filaments
dynamic roles in physiological activities, 134:971
filensin and phakinin form beaded, coassemble de novo in cultured cells, 132:643
function in cell shape and cytoskeletal integrity, 134:971
in maintenance of mechanical integrity and strength of myofibers, 134:1255
plectin interaction with, 134:1455
plectin-mediated interaction of microtubules with, 135:991
role of pinin in reinforcement of desmosome complex, 135:1027
Internalization, initial step of activity-induced sodium channel down-regulation mediated by, 134:499
Interphase, XIST RNA paints the inactive X chromosome in, novel RNA involved in nuclear/chromosome structure, 132:259
Interphase/mitosis transition, microtubule dynamics before and after nuclear envelope breakdown during, 135:201
Intracellular Ca^{2+}, mediates stimulus–secretion coupling in ZP3-activated pathway in sperm, 134:637
Intracellular functions, $\alpha_6\beta_4$ integrin, regulated by epidermal growth factor, 134:241
Intracellular signals, direct integrin localization to sites of function in embryonic muscles, 134:217
Intranuclear microtubule organizer, role of calmodulin and Spc110p interaction in proper assembly of spindle pole body components, 133:111
Invariant chain, isoforms of, regulate transport of MHC class Ii molecules to antigen processing compartments, 133:281
Involuting marginal zone cells, attachment to fibronectin, mediated by RGD sequence in type III-10 repeat, and by PPRAR sequence in type III-13 of Hep II domain, 134:227
Ion channel
activity of, influenza virus M_2 protein, transport through Golgi apparatus, 133:733
subunit required for mechanotransduction in *Caenorhabditis elegans*, sequence and transmembrane topology of MEC-4, 133:1071
Isoforms
Dictyostelium mutants lacking multiple classic myosin I, combinations of shared and distinct functions, 133:305
invariant chain regulate transport of MHC class II molecules to antigen processing compartments, 133:281
mammalian cells express three distinct dynein heavy chains that are localized to different cytoplasmic organelles, 133:831
overlapping functions of myosin-I, 133:221
J domain, functions of auxilin at, cooperation effects of clathrin domain and, 135:925
Jun proteins, Ap-1 DNA binding activity attributed to, transformed and nontransformed cells effected by, 135:1151
Junctional epidermolysis bullosa, β4 integrin in, 134:559
Jun-N-terminal kinases, roles in anoikis of ICE-related cysteine protease and, 135:1377
Karyogamy, ER membrane protein complex required for nuclear fusion, 132:499
Karyoplasm, localization of plakophilin 2 in, 135:1009
KDEL-receptor, in COP I vesicles, 134:1411
Keratin filaments, envoplakin links cornified envelope to, 134:715
Keratins
nontetrameric species is major soluble form of, *Xenopus* oocytes and rabbit reticulocyte lysates, 132:153
14-3-3 proteins associate with phosphorylated, simple epithelial, cell cycle progression, act as solubility cofactor, 133:345
re-epithelialization after skin injury correlates with reorganization of filaments of, wound edge keratinocytes, potential role for keratin 16, 132:381
Keratocyte, locomotion of, 134:1209
Kidney
identification of small cytoplasmic ankyrin in muscle and, binds βIΣ* spectrin and associates with Golgi apparatus, Ank_{G119}, 133:819
in mice, molecular and functional consequences resulting from collagen α3(IV) deficiency in, related to Alport syndrome, 135:1403
Kinase A, phosphorylation of, influences plectin binding to vimentin and lamin B, 134:1455
Kinase C, phosphorylation of, influences plectin binding to vimentin and lamin B
Kinases, association with axonin-1 and Ng-CAM on dorsal root ganglion neurons, 135:253
Kinesin, role in transport of squid axon vesicles on microtubules, 135:383
Kinesin homologue protein 1, inner dynein arms but not outer dynein arms require activity of kinesin homologue protein $KHP1^{FLA10}$ to reach distal part of flagella in *Chlamydomonas*, 133:371

Kinesin-II, membrane traffic motor in axons, axonemes, and spindles, 133:1
Kinesin II, sequence and localization of 115-kD accessory subunit of heterotrimeric complex of, $KRP_{85/95}$, 132:371
Kinesin-related proteins, DSK1 is novel, diatom *Cylindrotheca fusiformis*, anaphase spindle elongation, 133:595
Kinetochore
component required for accurate chromosome segregation, 134:1127
force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during metaphase but during anaphase only at, 132:1093
localization of topoisomerase II α at region adjacent to, 134:1097
movement and response during prometaphase, force on chromosome depends on movement toward or away from pole, factors that create neutral motion of, centromere behavior after destruction of, 135:315
$KRP_{85/95}$, sequence and localization of 115-kD accessory subunit of heterotrimeric kinesin-II complex, 132:371
Kv2.1 K^+ channel, in MDCK cells, basis for localization of polarized and clustered distribution of, correlation of detergent solubility with, 135:1619
L1, alcohol inhibits cell–cell adhesion mediated by, human, 133:381
Lac operator-repressor staining, visualization of chromatin organization and DNA sequences using, 135:1685
lacZ
expression of, in mesenchymal cells, controlled by far-upstream enhancer of mouse proα2(I) collagen gene, 134:1333
expression of, in notochord, 134:1573
Lamellar ichthyosis, epidermal transglutaminases in, 134:715
Lamin B receptor homologue, p56 mediates membrane targeting to chromatin in sea urchins, 135:1715
Laminin
-1, inhibition of laminin α1-chain expression leads to alteration of basement membrane assembly and cell differentiation, 133:417
5, anchorage mediated by integrin α6β4 to, epiligrin, regulates tyrosine phosphorylation of membrane-associated protein, 132:727
antiglycolipid antibody inhibits Madin-Darby canine kidney cell adhesion to, interferes with basolateral polarization and tight junction formation, 133:695
in myogenesis, 134:1483
neurite outgrowth domain of B2 chain of, antibodies against, 134:477
Laminin α2 chain, partial restoration of, α2 chain-deficient *dy/dy* mouse, primary muscle cell culture transplantation, 133:185
Laminin β2 chain, contained in native chick laminin-4, promotes motor axon growth, 135:1583
Laminins, human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 132:1189
Lamins, nuclear events during p53-dependent apoptosis facilitated by degradation of, 135:1441
Lamp1, targeting to lysomes, dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 132:565
Lampellipodium, protrusion of, and forward translocation of cell body, in rapidly moving keratocyte, 134:1209
LAR-related member of transmembrane tyrosine phosphatase family (LAR-PTP), associates with cadherin–catenin complex, 134:1519
Lck, associated with glycolipid-enriched membrane domains, relationship to CD45 of, 135:1515
Lens
filensin and phakinin form beaded intermediate filaments and coassemble de novo in cultured cells, 132:643
human multidrug resistance 3–P-glycoprotein expression in transgenic mice induces membrane alterations of, leading to cataract, 132:701
Leukocytes
adhesion, locomotion, and chemotaxis, regulation by rhodopsin-related Ga_i-linked 7 transmembrane "serpentine" receptors of chemoattractant subfamily, 134:255
adhesion to vascular endothelium induces E-selectin linkage to actin cytoskeleton, 133:445
characterization of L-selectin ligands on, tethering and rolling adhesions in shear flow on, 135:837
interactions through L-selectin of, adhesions in shear flow effected by, 135:849
Leupeptin, effect on degradation of sodium channel, 134:499
Lever arm, in muscle myosin head domain, 134:895
Ligands
endoglin modulates cellular responses to TGF-β1, 133:1109
expression in immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover, Fas and Fas ligand in embryos and adult mice, 133:335
fucosyltransferase FucT-VII regulates E-selectin ligand synthesis in human T cells, 133:911
Ligation, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 132:475
LIM domains, determinant of localization of paxillin focal adhesions, 135:1109
LIM proteins, role in muscle differentiation in vertebrates, 134:1179
Linker histone, δ, phosphorylation of, occurs early in *Tetrahymena* conjugation when micronuclei become transcriptionally active, 135:1219
Lipopeptide S-acylation, products of, localize preferentially to plasma membrane, 134:647
Localization to ER, of Sec12p, by static retention in ER and dynamic retrieval from early Golgi compartment, 134:279
Low density lipoprotein, traffic, polarity, and detergent solubility of glycosylphosphatidylinositol–anchored protein after deprivation of, MDCK cells, 133:1265
LRE, in β2 chain in native chick laminin-4, structural significance of, 135:1583
lsd1, impaired nuclear division in, 134:949
L-selectin
characterization of ligands for, in mediation of tethering and rolling adhesions in shear flow, 135:837
interaction with neutrophils and, relationship between P-selectin glycoprotein and, 135:523
interactions of leukocytes through, rolling adhesions effected by, 135:849
Lumen, Cab45 is novel Ca^{2+}-binding protein localized to Golgi, 133:257
Lungs
heparin induces dimerization and confers proliferative activity onto hepatocyte growth factor antagonists NK1 and NK2, 133:709
inactivation of integrin β6 subunit gene reveals role of epithelial integrins in regulating inflammation in skin and, 133:921
Lymphoblastoid cells, syndecan-1 mediates cell spreading in transfected human, Raji cells, 132:1209
Lymphocyte function-associated antigen 3, visualization of CD2 interaction with, determination of two-dimensional dissociation constant for adhesion receptors in contact area, 132:465
Lymphocytes, endocytosis of GPI-anchored proteins in human, role of glycolipid-based domains, actin cytoskeleton, and protein kinases, 133:791
Lymphoid cells, adhesion-triggering versus chemotactic responses, differential requirements for, 134:255
Lysophosphatidic acid
and bFGF control different modes in proliferating myoblasts, 132:181
role of ventricular zone gene-1 in signaling of, 135:1071
Lysosomal enzymes, targeting of, overexpression of neither mannose 6-phosphate receptor (MPR)46 nor MPR 300 sufficient for, along intracellular routes, 134:615
Lysosomal proteins, corrected missorting of, by overexpression of mannose 6-phosphate receptor 300, in part due to transport along a route sensitive to mannose 6-phosphate, 134:615
Lysosomes
AP-2-containing clathrin coat protein initiates assembly on dense, 135:1801
availability at cell periphery, required for *Trypanosoma cruzi* entry into host cells, 134:349
cysteine^{34} of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and enzyme sorting of, 132:577
fates of phagocytosed particles after delivery into, macrophages, 132:585
fusion with phagosomes, calcium-independent event in macrophages, 132:49
multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with, 132:1011
protease fusion of, strategy for engineering dominant negative mutations using, 135:1043
recruitment during trypanosome invasion, 134:349
targeting of LAMP1 to, dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 132:565
vesicular transport of MHC class II molecules to plasma membrane from structures of, 135:611
M band, sarcomeric, structure of, 134:1441
MAC-2, granulocyte macrophage colony stimulating factor produced in lesioned peripheral nerves induces up-regulation of cell surface expression of, macrophages and Schwann cells, 133:159

Macrophages
fates of phagocytosed particles after delivery into lysosomes of, 132:585
granulocyte macrophage colony stimulating factor produced in lesioned peripheral nerves induces up-regulation of cell surface expression of MAC-2 by Schwann cells and, 133:159
phagosome–lysosome fusion is calcium-independent event in, 132:49
Macropinocytosis, role of phosphoinositide 3-kinase in completion of, 135:1249
Madin-Darby canine kidney cells
antiglycolipid antibody inhibits adhesion of, to laminin, interferes with basolateral polarization and tight junction formation, 133:695
traffic, polarity, and detergent solubility of glycosylphosphatidylinositol–anchored protein after LDL-deprivation of, 133:1265
Major histocompatibility complex class II molecules
biogenesis of, in human I-Cell Disease B lymphoblasts, 132:769
isoforms of invariant chain regulate transport of, antigen processing compartments, 133:281
Male gametocytes, of *Plasmodium*, identification and characterization of developmental defect of, 135:269
Mammalian cells
association of dynamin-like protein with Golgi apparatus in, 133:761
express three distinct dynein heavy chains that are localized to different cytoplasmic organelles, 133:831
spatiotemporal constraints on replication of mtDNA in, 135:883
Mammalian eps15, homologue identified through novel genetic screen for yeast endocytosis mutants, 135:1485
Mammalian homologues, interspecies diversity of occludin sequence, cDNA cloning of human, mouse, dog, and rat–kangaroo homologues, 133:43
Mammalian telomeres, interaction of TRF with DNA of, 135:867
Mammary carcinogenesis, estrogen-dependent c-JunER protein causes reversible loss of mammary epithelial cell polarity involving destabilization of adherens junctions, 132:1115
Mammary epithelial cells, effect of metalloproteinase inhibitor on apoptosis and entactin of, 135:1669
Mann II, in COP I vesicles, 134:1411
Mannose 6-phosphate receptors
cysteine^{34} of cytoplasmic tail of, reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 132:577
MPR 46 and MPR 300, deficiency of, causes missorting of soluble lysosomal proteins into medium, 134:615
Mannosyltransferase, localization of, yeast early Golgi, involves retrograde transport, Och1p, 132:985
MAP kinase
mechanisms of synergy between integrins and growth factors involving, 135:1633
phosphorylation of Sir3p activated by, in transcriptional silencing regulation, 135:571
signaling pathway involving, in angiotensin II activation of, 135:1609
MAP-homologous protein 1, formation and stabilization of microtubules in *Saccharomyces* depends on, 135:1323
Mast cells, of mice, assessment of protease content during *Trichinella spiralis* infection of, changes in chymase phenotypes of, 135:279
Mastocytosis, of mice, assessment of protease content during *Trichinella spiralis* infection, changes in chymase phenotypes during, 135:279
Matrix assembly, fibronectin, differential requirements for RGD sequence and first type III repeats during, 134:573
Matrix metalloproteinase, functional evaluation of, in mammary gland growth and development in mice, 135:1669
Matrix receptors, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 132:1161
Matrix-targeting signal, excluded from mammalian mitochondria by hTom20p, 134:307
Maturation-promoting factor, novel component of, cdc2 kinase, triggering of M-phase in starfish oocytes, 132:125
MDCK cells
basis for localization of polarized and clustered distribution of Kv2.1 K^+ channel in, 135:1619
expression of wild-type and mutant chicken occludin in, 134:1031
polarized sorting mechanism in apical and basolateral endosomes of, 135:139
transformation of, by activated focal adhesion kinase, 134:793
mdx, transgenic mice, with normal expression of dystrophin protein with deletion in domain I, show mild dystrophic phenotype, 134:873
MEC-4, sequence and transmembrane topology of, ion channel subunit required for mechanotransduction in *Caenorhabditis elegans*, 133:1071
Mechanotransduction, ion channel subunit required for, *Caenorhabditis elegans*, sequence and transmembrane topology of MEC-4, 133:1071
Medium, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes in chemically defined (HGM) medium, induced by HGF/SF, EGF, and TGFα, 132:1133
Meiotic checkpoint, triggering of M-phase in starfish oocytes, novel component of maturation-promoting factor besides cdc2 kinase, 132:125
Melanogenesis, stimulation by cAMP elevating agents, 134:747
Melanoma, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 132:475
Melanoma cells
effects of thrombospondin through integrin-associated protein on vitronectin of, 135:533
in human, CD437 activates and regulates AP-1 which inhibits growth and induces apoptosis in, 135:1889
Membrane, in central nervous system, of L-glutamate transporter domain, integration and topology of, 135:1867
Membrane folate receptor-α, endocytosis of GPI-linked, 132:35
Membrane fusion, mediated by influenza virus hemagglutinin requires action of at least three hemagglutinin trimers, 133:559
Membrane proteins
activation of transcription factor NF-κB by adenovirus E3/19K protein requires ER retention, 132:511
of ER required for nuclear fusion, 132:499
Membrane ruffling, overexpression of cofilin stimulates bundling of actin filaments, cell movement, and, *Dictyostelium*, 132:335
Membrane skeleton, neuroglian-mediated cell adhesion induces assembly of, cell contact sites, 133:647
Membrane targeting, in sea urchins, lamin B receptor-like integral membrane protein mediates, 135:1715
Membrane traffic motors, kinesin II, axons, axonemes, and spindles, 133:1
Membrane trafficking
cell cycle regulation of dynein association with membranes modulates microtubule-based organelle transport, 133:585
identification, localization, and functional implications of an abundant Nematode annexin, 132:1079
signal transduction by polymeric immunoglobulin receptor suggests role in regulation of receptor transcytosis, 133:997
Merlin, patterns of expression imply distinct functions for *Drosophila* homologues of moesin and, neurofibromatosis 2 tumor suppressor, 133:843
Merosin, in myogenesis, 134:1483
Messenger RNA isoforms, identification of two Sox17, with and without high mobility group box region, differential expression in mouse spermatogenesis, 133:667
Messenger RNA precursors, nuclear cap-binding complex binds, Balbiani ring, cotranscriptionally, accompanies ribonucleoprotein particle during nuclear export, 133:5
Metalloprotease/disintegrin/cysteine-rich protein 9, is widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains, 132:717
Metanephroi, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 132:1161
Metaphase
detection of protein kinase activity specifically activated at transition with anaphase, 132:635
force for poleward chromosome motion in *Haemanthus* cells acts along length of chromosome during, but only at kinetochore during anaphase, 132:1093
Metaphase chromosomes, detection frequency of telomeres in, 134:1
Metastasis, and integrin cell surface adhesion receptors, 134:1551
Metazoan, rDNA enhancer from, acts by making more genes transcriptionally active, 133:943
5-methyltetrahydrofolate, delivered to cell cytoplasm of MA104 cells, by potocytosis, 134:1169
MHC, class II molecules of, transport from lysosomal structures to plasma membrane of, 135:611
MHP1, formation and stabilization of microtubules in *Saccharomyces* depends on, 135:1323
Mice
assessment of protease content during *Trichinella spiralis* infection of mast cells of, changes in chymase phenotypes of mast cells, 135:279
chondrogenic differentiation of clonal, embryonic cell line ATDC5 in vitro, gene expression of parathyroid hormone/PTH-related peptide receptor, 133:457

disrupted glial fibrillary acidic protein network in astrocytes from vimentin knockout, 133:853
distal-less homeodomain gene regulates epidermal differentiation in, 135:1879
Fas and Fas ligand in embryos and adult, ligand expression in immune-privileged tissues and coexpression in adult tissues characterized by apoptotic cell turnover, 133:335
functions of α_4 integrin in skeletal muscle of, 135:829
identification of two Sox17 messenger RNA isoforms and their differential expression in spermatogenesis in, with and without high mobility group box region, 133:667
nonsense mutation in dystrophin gene in *mdx*, leads to complete absence of dystrophin in muscle, 134:93
partial laminin α2 chain restoration in α2 chain-deficient *dy/dy* mouse by primary muscle cell culture transplantation, 133:185
renal consequences of collagen α3(IV) deficiency in, related to Alport syndrome, 135:1403
subunit composition of neurofilaments specifies axonal diameter, 133:1061
tissue-specific regulation of transcription of α1(VI) collagen chain in transgenic, 135:1163
transgenic expression of BMP-6 in, psoriasis formed by overexpression of bone morphogenetic protein-6 in, inhibition or stimulation of cell proliferation depending on the pattern of expression of BMP-6 in, 135:227
Microenvironments, lympho-hematopoietic, 134:771
Microphthalmia, role in regulation of mouse tyrosinase gene expression by cAMP, 134:747
Microtubule
dynamics before and after nuclear envelope breakdown, during interphase/mitosis transition, 135:201
sensitivity to G1 arrest caused by Taxol, 135:689
Microtubule assembly
influence of M-phase chromatin on anisotropy of microtubule asters, 133:125
and transport during axon growth, 133:151
Microtubule nucleation, model of γ-tubulin rings, 135:5
Microtubule organizing centers, 134:429, 134:443
Microtubule organizing centers, astral, 134:455
Microtubule transport, and microtubule assembly during axon growth, 133:151
Microtubule-associated protein 4, removal from microtubules in vivo produces no observable phenotype at cellular level, 132:345
Microtubule-associated protein 2C, selective stabilization in cell bodies and dendrites, Tau in axons, polarized localization of cytoskeletal proteins in mature neurons, 132:667
Microtubule/kinesin-mediated transport, required for *Trypanosoma cruzi* entry into host cells, 134:349
Microtubules
association with aberrant kinetochores, 134:1097
cell cycle regulation of dynein association with membranes modulates organelle transport based on, 133:585
effects of Rab8 on reorganization of actin and, 135:153
in living normal rat kidney cells, correlation of TD60 and midzone bundles of, requirements for cytokinesis, 135:981
PF16 localizes to a single, central apparatus in *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 132:359
plectin-mediated interaction of intermediate filaments with, 135:991
removal of MAP4 from, in vivo, no observable phenotype at cellular level, 132:345
roles of Eg5, dynein and dynactin in organization of, opposing motor activities of, 135:399
in *Saccharomyces*, MHP1 necessary in formation and stabilization of, 135:1323
tubulin transport in neurons, 133:1355
yeast ubiquitin-like genes involved in duplication of organizing center of, 133:1331
Migration
cell, modulation of, by integrin-mediated cytoskeletal linkages and ligand-binding affinity, 134:1551
of oligodendrocyte-type 2 astrocytes, NMDA as receptor for, role of PSA-NCAM in, 135:1565
Mini-chromosomes, aberrantly segregating centromeres activate spindle assembly checkpoint in budding yeast, 133:75
Miniparamyosin, in *Drosophila* muscles, 134:81
Minus-end vesicles, of squid axon vesicles, overridden by plus-end motors during transport, 135:383
Mitochondria
control by supply of, respiratory substrates, characterization of autophagy in higher plant cells subjected to carbon deprivation, 133:1251
role of N-myristoylation in association of NADH-cytochrome b_5 reductase with, 135:1501
Mitochondrial import process, AGT mistargeting in, 135:939
Mitochondrial protein sorting, 134:307
Mitogen-activated protein kinase, phosphoprotein stathmin essential for nerve growth factor–stimulated differentiation, 133:1383
Mitoplasts, generation of, for removal of proximal components of outer membrane import machinery, 134:307
Mitosis
calcium signals associated with entry into, in sea urchin embryos, 135:191
cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first division of, 132:915
detection of protein kinase activity specifically activated at metaphase-anaphase transition, 132:635
in human cells, polo-like kinase1 required for entry into, 135:1701
molecular characterization of 50-kD subunit of dynactin reveals function for complex in chromosome alignment and spindle organization during, 132:617
rDNA transcription machinery is assembled during, active NORs and absent in inactive NORs, 133:235
removal of MAP4 from microtubules in vivo produces no observable phenotype at cellular level, 132:345
XIST RNA paints the inactive X chromosome in interphase, novel RNA involved in nuclear/chromosome structure, 132:259
Mitotic checkpoint, *Saccharomyces cerevisiae* spindle pole body duplication gene *MPS1* is part of, 132:111
Mitotic delay, after DNA damage, inhibitory CDC2 phosphorylation partly responsible for, 134:963
Mitotic mechanisms in Alzheimer's disease, 132:413
MK-801, rescues weaver granule neurons similar to verapamil in high concentrations, 134:477
Mlp gene, expression of, is developmentally regulated in *Drosophila*, 134:1179
Moesin, patterns of expression imply distinct functions for *Drosophila* homologues of, neurofibromatosis 2 tumor suppressor, merlin, 133:843
Molecular chaperones
endoplasmic reticulum storage disease causing congenital goiter and hypothyroidism, 133:517
intracellular interaction of collagen-specific stress protein HSP47 with newly synthesized procollagen, 133:469
Molecular motors, parallels between G proteins and, 135:291
Moloney murine leukemia virus. Gag precursor of, targeting to virus budding site of, 135:1841
Monocytes, endoglin modulates cellular responses to TGF-β1, 133:1109
Monosome, 80S, native 3D structure of eukaryotic 80S ribosome, morphological homology with *E. coli* 70S ribosome, 133:495
Morphogenesis
anastral meiotic spindle, 134:455
motogenic and morphogenic activity of epithelial receptor tyrosine kinases, 133:1095
Mossy fiber terminals, in hippocampus, 134:511
Mother/bud neck, localization of Bud4p to, 134:413
Motility
biochemical kinetic characterization of myosin-I ATPase, *Acanthamoeba*, 132:1053
Dictyostelium mutants lacking multiple classic myosin I isoforms reveal combinations of shared and distinct functions, 133:305
Motility
epidermal growth factor receptor–mediated, actuated by gelsolin, 134:689
motogenic and morphogenic activity of epithelial receptor tyrosine kinases, 133:1095
two classes of oligoproline sequences in profilin-mediated acceleration of, actin-based, *Shigella*, 133:49
Motoneurons, striatin conspicuous in, 134:1051
Motor axon, laminin β2 chain in native chick laminin-4 promotes growth of, 135:1583
Motor domains, kinesin II is membrane traffic motor in axons, axonemes, and spindles, 133:1
Motor function, actin, modulated in general by α-smooth muscle actin, 134:67
Motor neuron disease, mediated by NF-H in mice, 135:711
Mouse eggs, at fertilization, function of Rabphilin-3A in calcium-dependent cortical granule exocytosis in, 135:1741

Mouse embryonic fibroblasts, deficient in mannose 6-phosphate receptors, missort soluble lysosomal proteins into medium, 134:615
M-phase, influence of chromatin during, anisotropy of microtubule asters, 133:125
M-protein, localizes to M band, 134:1441
MPS1 gene, of spindle pole body in *Saccharomyces cerevisiae*, part of mitotic checkpoint, 132:111
mRNA, export from nucleus, mediated by heterogeneous nuclear RNP (hnRNP) proteins, 134:1365
mtDNA
  loss of, results in suppression of *ssh1* cold-sensitive phenotype, 134:603
  spatiotemporal constraints on replication of, effect of BrdU on, 135:883
Multidrug resistance 3–P-glycoprotein, expression in transgenic mice induces lens membrane alterations leading to cataract, human, 132:701
Multivesicular bodies
  multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and fuse with lysosomes, 132:1011
  mutations in *Drosophila hook* gene inhibit endocytosis of boss transmembrane ligand into, 133:1205
Muscle
  agrin-induced acetylcholine receptor clustering in mammalian, requires tyrosine phosphorylation, 132:937
  of flight, structure and function of, in *Drosophila* tropomyosin mutants, 135:673
  gene *unc-89* required for assembly of muscle M-line, encodes giant modular protein composed of Ig and signal transduction domains, *Caenorhabditus elegans*, 132:835
  three-dimensional structure of Z band in normal mammalian skeletal, 133:571
Muscle attachments, in *Drosophila* embryo, chimeras containing β_{PS} cytoplasmic tail localize to ends of muscles with endogenous integrins, or in their absence, 134:217
Muscle cells
  β1D integrin displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 132:211
  distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, expression of *SM22α* promoter in transgenic mice, 132:849
  partial laminin α2 chain restoration in α2 chain-deficient *dy/dy* mouse by primary, cell culture transplantation, 133:185
  tropinin T mutations in striated and nonstriated, *Caenorhabditis elegans*, 132:1061
Muscle connective tissue, tenascin-Y in, 134:1499
Muscle differentiation
  insulin-like growth factor binding protein-5 modulates, insulin-like growth factor–dependent mechanism, 133:683
  integrin α subunit ratios, cytoplasmic domains, growth factor synergy regulate muscle proliferation and, 133:169
Muscle gene expression, defects in, due to lack of retinoblastoma protein, 135:441
Muscular dystrophy, congenital, and merosin deficiency, 134:1483
Mutation, dominant negative, lysosomal protease fusion in formation of, 135:1043
Myelin, Po protein, truncated, dominant-negative effect on adhesiveness of full-length Po, 134:1531
Myoblast fusion, enhanced by neural CAM, 135:241
Myoblasts
  dysgenic, show higher incidence of cell fusion than normal counterparts, 134:375
  integrin α subunit ratios, cytoplasmic domains, growth factor synergy regulate muscle proliferation and differentiation, 133:169
  lysophosphatidic acid and bFGF control different modes in proliferating, 132:181
  myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis, 132:657
Myocardium, progressive degeneration and necrosis of, in desmin null mice, 134:1255
Myocytes, induction of DNA synthesis and apoptosis in cardiac, E1A oncoprotein, 133:325
Myofibers, neural CAM expression results in changes to, 135:241
Myofibrils, M band of, 134:1441
Myogenesis
  cell differentiation in, late role of Mlp60A and Mlp84B, 134:1179
  functions of α_4 integrin in, 135:829
  IGF-I activates proliferation and then subsequent events of, 135:431
  insulin-like growth factor binding protein-5 modulates muscle differentiation through insulin-like growth factor–dependent mechanism, 133:683
  lysophosphatidic acid and bFGF control different modes in proliferating myoblasts, 132:181
  myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon, 132:657
Myomesin, localizes to M band, 134:1441
Myopathy, and desmin deficiency, 134:1255
Myosin
  heavy chain, in *Caenorhabditis*, characteristic variations in hydrophobicity along rods of, filament initiation of, 135:371
  recombinant, motor activity of, 134:895
  role in actin cable assembly during wound healing of, 135:1097
  in *Saccharomyces*, function in vacuole inheritance of actin and, 135:1535
Myosin filaments
  square backbone and crossbridges, 134:53
  structure, in vertebrate smooth muscle, 134:53
Myosin heavy chain, *Xenopus*, nonmuscle, isoforms of, 134:675
Myosin heavy chain A (MHC-A), localizes to mitotic spindle, 134:675
Myosin heavy chain B (MHC-B), in mitotic prophase, localizes to nuclear membrane, 134:675
Myosin heavy chain kinase A, regulates myosin localization during growth and development, *Dictyostelium*, 132:101
Myosin I
  Dictyostelium mutants lacking multiple classic isoforms of, combinations of shared and distinct functions, 133:305
  overlapping functions of isoforms of, 133:221
  synthetic lethality screen identifies novel yeast gene of, *MYO5*, myosin I proteins required for polarization of actin cytoskeleton, 133:1277
Myosin II, *Dictyostelium* myosin heavy chain kinase A regulates localization of, growth and development, 132:101
Myosin II heavy chain, phosphorylation by MHC-protein kinase C in cell cortex, leads to myosin II dissociation from cytoskeleton, 134:911
Myosin motor domain, site-directed mutagenesis of, 134:895
Myosin tails, antiparallel arrangement of, and bare ends in, filament from skinned smooth muscles, 134:53
Myosin-I, biochemical kinetic characterization of ATPase of, *Acanthamoeba*, 132:1053
Myotube
  formation, role of merosin in, 134:1483
  primary–secondary interactions regulated by Neural CAM, 135:241
  stability, role of merosin in, 134:1483
Myotubes, contractile activity regulates isoform expression and polysialylation of NCAM in, involvement of Ca^{2+} and protein kinase C, 132:969
Myristic acid, novel role in NADH-cytochrome b_5 reductase targeting for, 135:1501
Myristylation, of FBR v-fos, cell-specific differentiation in osteosarcoma caused by, 135:457
Na^+, triggers activity-dependent endocytotic process that adjusts sodium channel density, 134:499
NADH-cytochrome b_5 reductase, role of N-myristoylation in targeting of, 135:1501
Na^+/H^+ antiport, intracellular pH regulation during spreading of human neutrophils, 133:1391
Na^+/H^+ exchangers, activity of during phagocytosis in human neutrophils, role of Fcγ receptors and tyrosine kinases, 132:1037
Na,K-ATPase, oligomerization and maturation of, interaction of cytoplasmic NH_2 terminus of β subunit with α subunit, 133:1193
N-cadherin
  augmenting effect of, suppression of tumorigenicity by plakoglobin, 133:199
  in squamous carcinoma cells, effect on E- and P-cadherin by, results in scattered fibroblastic phenotype, 135:1643
Neck filaments, septins in assembly of, 134:1345
NEDD-2, cysteine protease, link between processing and apoptotic activity of, cleavage events regulated by, 135:479
Nematodes
  identification, localization, and functional implications of an abundant annexin in, 132:1079
  ryanodine receptors and excitation–contraction coupling in, 134:885
Nephrogenesis, cloning of mouse integrin α_v cDNA, role of α_v-related matrix receptors in metanephric development, 132:1161
Nerve growth factor
  phosphoprotein stathmin essential for differentiation stimulated by, 133:1383
  target-derived influences of, 134:487
Nerve growth factor receptor complex, neurotrophin receptor forms complex with receptor tyrosine kinase TrkA, gp75, 132:945

Nerve growth factor-inducible large external glycoprotein, human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$, 132:475
N-ethylmaleimide, biochemically distinct steps revealed with GTPγS, BAPTA, and, assembly of nuclear pore, 132:5
Neural CAM, role in regulation of primary–secondary myotube interactions and muscle fusion, controlled by skeletal muscle α-actin promoter, 135:241
Neural cell adhesion molecule, contractile activity regulates isoform expression and polysialylation of, cultured myotubes, involvement of Ca^{2+} and protein kinase C, 132:969
Neurite fasciculation, intracellular signaling during, roles of NgCAM and axonin-1 during, 135:253
Neurite outgrowth, interaction between NgCAM and axonin-1 promotes, 135:1593
Neurite outgrowth domain, of B2 chain of laminin, 134:477
Neurites
  extensions of, effects of neurofascin and NrCAM on, 135:1059
  GRASP-NgCAM interactions involved in extension of, heterophilic interactions of DM-GRASP, 133:657
  tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 132:681
Neurofascin
  ankyrin-binding CAM, identified at node of Ranvier, 135:1355
  interaction of NrCAM with, neurite extensions effected by, 135:1059
Neurofibromatosis 2 tumor suppressor, patterns of expression imply distinct functions for *Drosophila* homologues of moesin and, merlin, 133:843
Neurofilament transport, effects of NF-H on, 135:711
Neurofilaments, subunit composition of, specifies axonal diameter, 133:1061
Neurogenic regions, of cerebral cortex, role of ventricular zone gene-1 in signaling of LPA in, 135:1071
Neuroglian, mediates cell adhesion, induces assembly of membrane skeleton at cell contact sites, 133:647
Neuromodulation, MAP kinase signaling pathway activated by angiotensin II mediates, 135:1609
Neuromuscular junction, RIα recruits Cα to, 134:1241
Neuronal cell adhesion molecules, characterization of, at node of Ranvier, 135:1355
Neurons
  neurotrophins promote survival and development of, cerebellum of hypothyroid rats in vivo, 133:631
  phosphoprotein stathmin essential for nerve growth factor–stimulated differentiation, 133:1383
  p53-induced apoptosis in, 135:1085
  tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for, 132:681
  tubulin transport in, 133:1355
Neurotoxin C1, of *Clostridium botulinum*, depletes membrane-bound syntaxin in situ, 134:205
Neurotoxins, activation of sodium channels by, 134:499
Neurotrophic factor, brain-derived, 134:487
Neurotrophin receptor, forms complex with receptor tyrosine kinase TrkA, gp75, 132:945
Neurotrophins
  promote survival and development of neurons in cerebellum of hypothyroid rats in vivo, 133:631
  in survival and development of sensory and sympathetic neurons, 134:487
Neutrophils
  CD47 mediates post-adhesive events required for migration of, polarized intestinal epithelia, 132:437
  deficiency of Src family kinases p59/61^{hck} and p58$^{c\text{-}fgr}$ results in defective functions of adhesion-dependent, 133:895
  interaction with L-selectin and, 135:523
  intracellular pH regulation during spreading of human, 133:1391
  Na^+/H^+ exchange activity during phagocytosis in, human, role of Fcγ receptors and tyrosine kinases, 132:1037
NF-κB, inhibition of IκB-α phosphorylation and degradation and subsequent activation of, glutathione peroxidase overexpression, 133:1083
NF-H, effect on neurofilament transport of, mediation of motor neuron disease in mice by, 135:711
NgCam, association with kinases, intracellular signaling during neurite fasciculation triggered by, effects of clustering with axonin-1, 135:253
NgCAM
  heterophilic interactions of DM-GRASP with, neurite extension, 133:657
  role in neurite outgrowth, interaction between axonin-1 and, 135:1593
NGF, inhibits metabolic and genetic events associated with neuronal cell death, functional comparison between ICE family proteases and, 135:1341
Nic96p
  required for nuclear pore formation and interacts with novel nucleoporin, Nup188p, 133:1141
  yeast nucleoporin Nup188p interacts genetically and physically with core structures of nuclear pore complex, 133:1153
NK1, heparin induces dimerization and confers proliferative activity onto hepatocyte growth factor antagonists NK2 and, 133:709
NLS, import rates of, kinetics of nuclear transport directed by, effect of Hsp70 on, 135:329
N-methyl-D-aspartate, as functional receptors of O-2A cells for migration, 135:1565
N-myristoylation, role in protein targeting for, required for NADH-cytochrome b_5 reductase association with mitochondria, 135:1501
Node of Ranvier, characterization of cell adhesion molecules at, 135:1355
Nomenclature, revision for peroxisome biogenesis factors, 135:1
Non-Claret disjunctional kinesin-like protein, required for normal spindle assembly kinetics, and spindle stabilization during metaphase arrest, 134:455
Novel protein, pinin, reinforcement role in intermediate filament–desmosome complex, 135:1027
NrCAM
  ankyrin-binding CAM, identified at node of Ranvier, 135:1355
  interaction of neurofascin with, neurite extensions effected by, 135:1059
NT-3, cutaneous overexpression of, increases sensory and sympathetic neuron numbers, 134:487
Nuclear division, impaired by deficiency of intracellular fatty acid, 134:949
Nuclear domain 10 (ND10), peripheries of, preferred sites for early steps of transcription and replication of three DNA virus families, 134:815
Nuclear envelope, accumulation of Nup153 in, 134:1141
Nuclear envelope breakdown
  cell cycle-associated change in Ca^{2+} releasing activity leads to generation of Ca^{2+} transients in mouse embryos during first mitotic division, 132:915
  microtubule dynamics before and after, 135:201
Nuclear export
  nuclear cap-binding complex binds Balbiani ring pre-mRNA cotranscriptionally and accompanies ribonucleoprotein particle during, 133:5
  targeting and function in, of Nup153, 134:1141
Nuclear export signals
  mediate nuclear export of shuttling heterogeneous nuclear RNP (hnRNP) proteins, 134:1365
  in Ran binding protein (RanBP1), 134:1157
Nuclear import
  evidence that RAN/TC4 GTPase mediates essential function independent of, protein, green fluorescent protein-glucocorticoid receptor chimera, 133:985
  GTPase distinct from Ran involved in, protein, 133:971
  Ran seems to participate in early step of, at nuclear pore complex, 134:589
  RAN/TC4 mutants identify common requirement for snRNP and protein import into nucleus, 133:485
Nuclear import receptor, dynamic localization of, interactions with transport factors, 133:1163
Nuclear localization, intron-dependent recruitment of pre-mRNA splicing factors to sites of transcription, 133:719
Nuclear localization sequences, targeting signals for nuclear protein import, 134:589
Nuclear localization signal, basic amino acid residue cluster in functionally intact sequence motif within, essential for vimentin binding, 134:1455
Nuclear matrix
  association of telomeric DNA and TRF with, 135:867
  concentration of translocated fibroblast growth factor receptors in, 134:529
Nuclear membrane fusion, ER membrane protein complex required for, 132:499
Nuclear pore complex
  assembly of, steps revealed with NEM, GTPγS, and BAPTA, 132:5
  association of Nup153 constructs with, 134:1141
  identification and localization of novel form of RanGAP1 involved with regulation of transport across, 135:1457
  mediates molecular exchanges between nucleus and cytoplasm, 134:589
  Nic96p required for formation of, interacts with novel nucleoporin, Nup188p, 133:1141
  yeast nucleoporin Nup188p interacts genetically and physically with core structures of, 133:1153

Nuclear protein import
effect of RanBP1 on p97 interactions in, 135:559
pathway of, involves interaction of transport ligand complex with cytoplasmic periphery of nuclear pore complex, 134:589
Nuclear proteins, of quiescent *Xenopus laevis* cells inhibit DNA replication in intact and permeabilized nuclei, 133:955
Nuclear retention sequence (NRS), in heterogeneous nuclear proteins (hnRNP), 134:1365
Nuclear transport, Hsp70 activity during targeting and translocation phases of, 135:329
Nucleolar organizer regions, rDNA transcription machinery is assembled during mitosis in active, absent in inactive NORs, 133:235
Nucleoporin
interacts genetically and physically with core structures of nuclear pore complex, yeast, Nup188p, 133:1153
Nic96p required for nuclear pore formation and interacts with novel, Nup188p, 133:1141
Nucleotide hydrolysis, parallels between molecular motors and G proteins, 135:291
NuMA, localization at mitotic spindle pole, effect of opposing motor activities on, 135:399
Nup153, role in nucleocytoplasmic transport, 134:1141
Nup188p
interacts genetically and physically with core structures of nuclear pore complex, yeast nucleoporin, 133:1153
Nic96p required for nuclear pore formation and interacts with, novel nucleoporin, 133:1141
Occludin
interspecies diversity of sequence of, cDNA cloning of human, mouse, dog, and rat–kangaroo homologues, 133:43
transmembrane component of tight junctions, 134:1031
Och1p, yeast early Golgi mannosyltransferase, localization involves retrograde transport, 132:985
Octanol, in gap junctional communication, 134:1019
O-glycoprotease, treatment of rolling cells with, blocks binding to stromal matrix, 134:1075
Oligodendrocyte-type 2 astrocyte, migration of, NMDA as receptor for, role of PSA-NCAM in migration of, 135:1565
Oligodeoxynucleotide (ODN), antisense, immediately and totally inhibits α-smooth muscle actin synthesis for >48 h, compared with sense ODN, 134:67
Oligomerization, and maturation of Na,K-ATPase, interaction of cytoplasmic NH_2 terminus of β subunit with α subunit, 133:1193
Oligomers, structure of, *Helicobacter pylori* vacuolating cytotoxin, 133:801
Oligoproline sequences, two classes of, profilin-mediated acceleration of actin-based motility, *Shigella*, 133:49
Oocytes
nontetrameric species is major soluble form of keratin in, *Xenopus*, rabbit reticulocyte lysates, 132:153
specificity of *emo-1* (*oz1*) defect to, 134:699
triggering of M-phase in, starfish, novel component of maturation-promoting factor besides cdc2 kinase, 132:125
Oogenesis
cell type–specific roles for Cdc42, Rac, and RhoL in, *Drosophila*, 133:617
emo-1 gene product necessary for, 134:699
Open reading frame (ORF), major, regulation of expression altered by mutation of either upstream (u)ORF2, ORF4, or uORF5 start codon, 134:827
Organelle inheritance, thioredoxin required for vacuole inheritance in *Saccharomyces cerevisiae*, 132:787
Organelle transport, cell cycle regulation of dynein association with membranes modulates microtubule-based, 133:585
Osteoblasts, expression of TGF-β2 in, results in osteoporosis-like phenotype, 132:195
Osteoporosis, expression of TGF-β2 in osteoblasts results in osteoporosis-like phenotype, 132:195
Osteosarcoma, myristylation of FBR v-fos causing, 135:457
oz151, putative *emo-1* null mutation, displays embryonic lethality, 134:699
p23, characterization of, is Golgi receptor for coatomer involved in formation of COPI-coated vesicles, 135:1239
p36, catecholamine secretion requires translocation of, subplasmalemmal region in chromaffin cells, annexin II in exocytosis, 133:1217
p47, novel adaptor-related protein complex, 133:749
p53, expression induced by camptothecin, 134:757
p53
-dependent apoptosis, lamin degradation facilitates nuclear events during, 135:1441
hematopoietic and muscle differentiation, inhibited by interference with, 134:193
postmitotic neuronal apoptosis induced by, 135:1085
p56, lamin B receptor-like integral membrane protein, membrane targeting to chromatin in sea urchins mediated by, 135:1715
p58, p54, immobilized recombinant, deplete nuclear transport activity from cytosol, 134:589
$p59^{fyn}$, features of src homology 4 domain, define motif for immune-receptor tyrosine-based activation motif binding and for plasma membrane localization, 133:1007
p62 complex
assembly of nuclear pore complex glycoproteins, directly implicated in nuclear protein import, 134:589
localizes on both nucleoplasmic and cytoplasmic surfaces of nuclear pore complex near central gated channel, 134:589
p65, transdominant-negative mutant of, partially inhibits apoptosis, 134:13
p97, effect of RanBP1 on interactions with, in nuclear protein import, 135:559
Pairing, meiotic chromosome, 134:1109
PAL1 gene, product of is peroxisomal ATP-binding cassette transporter in *Saccharomyces cerevisiae*, 132:549
Palmitoylation, cysteine[34] of cytoplasmic tail of cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting, 132:577
Pancreas, distribution of α subunits and βγ subunits of heterotrimeric G proteins on Golgi membranes of exocrine, 133:1027
Pan1p, endocytosis role of, identified by novel genetic screen, 135:1485
Pantophysin, cDNA related to synaptophysin, defines constitutive transport vesicles, 134:731
Paracellular permeability, expression of wild-type and mutant occludin increases, 134:1031
Paracrine mechanism, differentially expressed fibroblast growth factors regulate skeletal muscle development through autocrine and, 132:1151
Paramyosin
in *Drosophila* muscles, 134:81
and miniparamyosin, involvement in muscle function and flight, indicated by changes in isoform patterns, 134:81
Parathyroid hormone
collagen gene expression effected by, in bovine and human chondrocytes, 135:1179
gene expression of parathyroid hormone/PTH-related peptide receptor, chondrogenic differentiation of clonal mouse embryonic cell line ATDC5 in vitro, 133:457
Pas7p, is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase, Peb1p targeted to peroxisomes by NH_2-terminal peptide, 132:325
Pas10p, *Saccharomyces cerevisiae* peroxisomal targeting sequence 1 (PTS1) receptor, 134:849
Patches, movement of cortical actin, yeast, 132:861
Pathways, of synaptic vesicles, active and nonactive zones of, 135:797
Paxillin, viniculin- and FAK-interaction domains on, focal adhesion targeting motif of, 135:1109
PC5, proprotein convertase, COOH-terminal variant isoforms of, biosynthesis of, sorting to different subcellular locations by, 135:1261
PC12 cells
AP-1 adaptor complex binds to immature secretory granules from, regulated by ADP-ribosylation factor, 132:523
targeting of P-selectin to regulated secretory organelles of, 134:1229
Pds1p
plays critical role in APC and checkpoint pathways, inhibitor of anaphase in budding yeast, 133:99
required for execution of anaphase in yeast, *Saccharomyces cerevisiae*, 133:85
PDZ domains
of hDlg, interaction between protein 4.1 and, 135:1125
structural organization of, interactions of ATP and protein 4.1 with proteins of, 135:753
Pea2p, in *Saccharomyces*, characterization of mutants defective in Spa2p and, role in polarized growth and budding of, 135:725
Peb1p, is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase, Peb1p targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 132:325
Pericentriolar recycling compartment, interaction between rab11 and transferrin through, 135:913

Peripheral nerves, granulocyte macrophage colony stimulating factor produced in lesioned, induces up-regulation of cell surface expression of MAC-2 by macrophages and Schwann cells, 133:159
Peroxisomal import pathway, dissection of, caused by depletion of Pmp47 in dihydroxyacetone synthase (DHAS), 134:37
Peroxisomal import process, AGT mistargeting in, 135:939
Peroxisomal membrane proteins, sorting sequence of peroxisomal integral membrane protein PMP47 is contained within short hydrophilic loop, 133:269
Peroxisomal targeting sequence 1 (PTS1), pathway, human catalase sorted by, 134:849
Peroxisomal targeting sequence type 2, Peb1p is intraperoxisomal receptor for, NH_2-terminal, thiolase, Peb1p targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 132:325
Peroxisome biogenesis factors, revised nomenclature for, 135:1
Peroxisomes
  interaction between Pex5p and Pex13p in SH3 domain, 135:97
  interaction between Pex13p in SH3 domain and Pex5p, 135:111
  PAL1 gene product is peroxisomal ATP-binding cassette transporter in *Saccharomyces cerevisiae*, 132:549
  Pex13p as docking factor for PTS1 receptor, 135:85
  proliferation caused by Pex11p, fragmentation caused by Pex11p, formation of homodimers in, 135:123
  repeated cycling of Pex5p between cytoplasm and, multiple PEX genes required for distribution and stability of Pex5p, 135:1763
  targeting of human catalase to, depends on novel COOH-terminal peroxisomal targeting sequence, 134:849
PEX genes, revised nomenclature for, 135:1
Pex5p
  interaction between Pex13p in SH3 domain and, 135:111
  interaction between Pex13p in SH3 domain and, as receptor for PTS1 protein, 135:97
  repeated cycling between cytoplasm and peroxisomes by, multiple PEX genes required for distribution and stability of, 135:1763
Pex11p, peroxisomal proliferation caused by, fragmentation of large peroxisomes caused by, homodimers formed by, 135:123
Pex13p
  function of, interaction between Pex5p and SH3 domain of, as peroxisomal receptor for PTS1, deficiency of, 135:111
  peroxisomal protein essential for protein import, SH3 domain of, interaction with Pex5p, 135:97
  peroxisome protein, as docking factor for PTS1 receptor, role of SH3 domain in function of, lack of, 135:85
PF16 gene, localizes to single microtubule of central apparatus in *Chlamydomonas* flagella, encodes a protein with armadillo repeats, 132:359
pH
  during spreading of human neutrophils, intracellular, 133:1391
  endosomal βCOP involved in formation of transport vesicles is dependent on, destined for late endosomes, 133:29
  in Golgi complex, measurement of, 134:1387
pH sensitivity, in interactions between F-actin, EF-1α and aminoacyl-tRNA, 135:953
Phagocytosis
  deficient signaling during FcεRI-mediated, 134:1427
  Dictyostelium mutants lacking multiple classic myosin I isoforms reveal combinations of shared and distinct functions, 133:305
  fates of phagocytosed particles after delivery into macrophage lysosomes, 132:585
  Na^+/H^+ exchange activity during, human neutrophils, role of Fcγ receptors and tyrosine kinases, 132:1037
  role of phosphoinositide 3-kinase in completion of, 135:1249
Phagosomes
  fates of phagocytosed particles after delivery into macrophage lysosomes, 132:585
  fusion with lysosomes, calcium-independent event in macrophages, 132:49
Phakinin, and filensin form beaded intermediate filaments and coassemble de novo in cultured cells, 132:643
Pharynx, *eat-5* required for synchronized muscle contractions of, 134:537
PH-domain, in regulatory region of protein kinase C μ (PKCμ), 134:1401
Phorbol 12-myristate 13-acetate (PMA), induces α_2 integrin mRNA level, through promoter-directed transcription, 134:1301
Phosphatidic acid, production of, involved in coatomer binding to membranes, 134:295
Phosphatidylinositol, effect on CD44/ERM interaction, 135:37
Phosphatidylinositol 4,5-bisphosphate, capping protein released from *n*-octyl β-D glucopyranoside (OG)-permeabilized platelets on treatment with, 134:389
Phosphatidylinositol 3-kinase, signaling pathway in acetylcholine receptor–inducing activity, 134:465
Phospho-epitopes, mitotic mechanisms in Alzheimer's disease, 132:413
Phosphoinositide 3-kinase, role in completion of macropinocytosis and phagocytosis, effect in pseudopod formation signaling, 135:1249
Phospholipase D, mediates ADP ribosylation factor–dependent formation of Golgi coated vesicles, 134:295
Phospholipase (PLC)-γ, epidermal growth factor-induced activation of, causes release of gelsolin from plasma membrane and remodeling of actin cytoskeleton in submembrane region of cell protrusions, 134:689
Phosphoprotein
  modified in cell cycle-dependent manner, spindle pole body component, *Saccharomyces cerevisiae*, 132:903
  stathmin essential for nerve growth factor–stimulated differentiation, 133:1383
Phosphorylation
  of adaptors, in vivo response of clathrin-coated vesicles to, 135:635
  agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine, 132:937
  anchorage mediated by integrin α6β4 to laminin 5 regulates tyrosine, epiligrin, membrane-associated protein, 132:727
  of α-factor receptor (Ste3p), 134:661
  of IκB-α is inhibited, degradation and subsequent NF-κB activation by glutathione peroxidase overexpression, 133:1083
  key mechanism in regulation of gap junction communication, 134:1019
  of linker histone δ, occurs early in *Tetrahymena* conjugation when micronuclei become transcriptionally active, 135:1219
  mitotic mechanisms in Alzheimer's disease, 132:413
  14-3-3 proteins associate with phosphorylated simple epithelial keratins during cell cycle progression and act as solubility cofactor, 133:345
  regulation of dynamic properties of intermediate filaments in vivo by, 134:971
  of retinoblastoma protein linked to adhesion-dependent cell cycle progression, expression of cyclin D1, activation of cyclin E-cdk2, 133:391
  of Sir3p, in transcriptional silencing regulation, activated by MAP kinase, 135:571
  Spc42p is phosphorylated component of spindle pole body with essential function during SPB duplication, *Saccharomyces cerevisiae*, 132:887
  spindle pole body component is phosphoprotein that is modified in cell cycle-dependent manner, *Saccharomyces cerevisiae*, 132:903
  stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine, PDGF β-receptors, 132:741
  of β_4 subunit at multiple tyrosine residues, by activation of epidermal growth factor receptor activation, 134:241
  synergy between integrins and growth factors activate, 135:1633
  TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 132:427
  tyrosine, of β-catenin, mediated by TrkA, 134:1519
  tyrosine, of Cek9, 135:781
  of vimentin by protein kinase C coupled with reorganization of intracellular membranes, mitosis-specific, 133:141
  Xenopus laevis actin-depolymerizing factor/cofilin is a protein regulated by, essential for development, 132:871
Phosphotyrosine, increases in levels of, correlate with changes in subcellular localization of Src and its substrates, 134:783
Photosystem II, two-dimensional crystals of, biochemical characterization, cryoelectron microscopy, localization of D1 and cytochrome b559 polypeptides, 132:823
PI-3 kinase, potential sites of, function in endocytic pathway revealed by PI-3 kinase inhibitor, wortmannin, 132:595
Pinin, characterization of novel protein, reinforcement role in intermediate filament–desmosome complex, 135:1027
PIP2-binding peptide, induces cell motility in absence of epidermal growth factor–receptor signaling, 134:689
PKC, behavior of, during formation stages of post-Golgi vesicles, 135:355
Plakoglobin
  binding to desmosomal cadherins, patterns of binding sites and topogenic potential, 133:359
  essential component of myocardiac desmosomes, effect of null mutation of gene for, 135:215

fascin binding to, 134:1271
suppression of tumorigenicity by, augmenting effect of N-cadherin, 133:199
Plakophilin 2, discovery and localization of, in desmosomes and karyoplasm, 135:1009
Plasma membrane
different biosynthetic transport routes to, BHK and CHO cells, 133:247
distribution of α subunits and βγ subunits of heterotrimeric G proteins on, Golgi, exocrine pancreas, 133:1027
features of $p59^{fyn}$ src homology 4 domain define motif for immune-receptor tyrosine-based activation motif binding and for localization of, 133:1007
transport of external Lys-Asp-Glu-Leu protein from, endoplasmic reticulum, KDEL, cholera toxin in vero cells, 133:777
vesicular transport of MHC class II molecules from lysosomal structures to, 135:611
Plasminogen activator inhibitor-1 (PA 1), governs urokinase receptor–mediated cell adhesion and release, 134:1563
Plasmodium, identification and characterization of developmental defect of male gametocytes of, 135:269
Platelet actin filament, barbed ends, coordinated regulation by gelsolin and capping proteins, 134:389
Platelet-derived growth factor
extracellular matrix alters regulation by, fibroblast integrins, 132:239
regulation of α_2 integrin gene expression by, 134:1301
stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors, 132:741
Pleckstrin homology domains, associations among SH3 domain–containing proteins, Rho-type GTPases and, yeast, 133:879
Plectin
in cytoskeleton, microtubule interaction with intermediate filaments mediated by, 135:991
defect in expression of, in epidermolysis bullosa simplex-muscular dystrophy, 134:1455
Plectin-vimentin network junction, basic amino acid residue cluster within nuclear targeting sequence motif, essential for, 134:1455
Plus-end vesicles, of squid axon vesicles, copurify with tightly bound kinesin motors, override minus-end motors during transport, 135:383
Pmp47, absence of, causes defects in transport and folding of specific matrix enzyme, 134:37
Po, myelin protein, truncated, dominant-negative effect on adhesion, 134:1531
Polarity
antiglycolipid antibody interferes with basolateral tight junction formation and basolateral, inhibits Madin-Darby canine kidney cell adhesion to laminin, 133:695
apicobasal, maintained by Discs large (Dlg) protein, in *Drosophila* epithelia, 134:1469
BED1 required for efficient bud emergence and polarized growth, *Saccharomyces cerevisiae*, gene encoding galactosyltransferase homologue, 132:137
myosin I proteins required for, actin cytoskeleton, synthetic lethality screen identifies novel yeast myosin I gene, *MYO5*, 133:1277
reversible loss in mammary epithelial cells, estrogen-dependent c-JunER protein, destabilization of adherens junctions, 132:1115
traffic, detergent solubility and, glycosylphosphatidylinositol–anchored protein after LDL-deprivation of MDCK cells, 133:1265
transport of vesicular stomatitis virus G protein to cell surface is signal mediated in polarized and nonpolarized cells, 133:543
Polarized membrane transport, promoted by Rab8, through reorganization of actin and microtubules in fibroblasts, 135:153
Polo-like kinase1, in human cells, maturation of mitotic centrosomes and spindle formation requires, 135:1701
Polyadenylated RNA, localization of viral RNA and snRNP and, 135:303
Polycystic kidney disease, sea urchin sperm receptor for egg jelly is modular protein with extensive homology to human protein of, PKD1, 133:809
Polymeric immunoglobulin receptor, signal transduction by, role in regulation of receptor transcytosis, 133:997
Polymerization
of actin, effected by elongation factor 1 α, 135:1309
of actin, effects of capping protein and polyphosphoinositides on, 135:169
active transport of photoactivated tubulin molecules in growing axons revealed by new electron microscopic analysis, 133:1347
of fibronectin, β1 integrin–dependent and –independent, 132:227
novel role for integrin-binding III-10 module in fibronectin matrix assembly, 133:431
tubulin transport in neurons, 133:1355
Polymorphonuclear leukocytes
adhesion triggers disorganization of endothelial adherens junctions, 135:497
chloride ion efflux induced by tumor necrosis factor in, 135:511
deficiency of Src family kinases $p59/61^{hck}$ and $p58^{c\text{-}fgr}$ results in defective adhesion-dependent neutrophil functions, 133:895
Polypeptide D1, biochemical characterization, cryoelectron microscopy, and localization of polypeptide cytochrome b559 and, two-dimensional crystals of photosystem II, 132:823
Polyphosphoinositides, effects on capping protein of actin, role in regulating actin polymerization, 135:169
Polysialylation, contractile activity regulates isoform expression and, NCAM in cultured myotubes, involvement of Ca^{2+} and protein kinase C, 132:969
Pore, formation and kinetics of fusion, induced by baculovirus GP64, 135:1831
Position, of genes, in periphery of chromosome territories, independent of gene expression, 135:1195
Post-Golgi vesicles, processes and requirements of coat and bud assembly and vesicle scission in formation of, behavior of PKC on formation stages of, 135:355
Postsynaptic compartment, characterized by presence of postsynaptic density, in type I synapses, 134:1051
Potassium ion channel, in MDCK cells, basis for localization of polarized and clustered distribution of Kv2.1, correlation of detergent solubility with, 135:1619
Potocytosis, folate receptors clustered in caveolae necessary for, 134:1169
$pp125^{FAK}$, functions to suppress apoptosis in fibroblasts, 135:1383
Pre-mRNA
dynamic transcription-dependent organization of splicing factors of, polytene nuclei, 133:929
intron-dependent recruitment of splicing factors of, sites of transcription, 133:719
Pre-B cells, interaction with stromal cell products, 134:771
Precursor proteins, interactions with components of chloroplast envelope, 134:315
Primary hyperoxaluria type I, AGT dimerization in peroxisome-to-mitochondrion mistargeting in, 135:939
proα2(I), far-upstream enhancer in, regulates expression of reporter genes in transgenic mice, 134:1333
Procollagen, intracellular interaction of collagen-specific stress protein HSP47 with newly synthesized, 133:469
Profilin
binding of, controlled by proline-rich repeats in ActA protein of *Listeria monocytogenes*, 135:647
mediates acceleration of actin-based motility, *Shigella*, two classes of oligoproline sequences, 133:49
Programmed cell death
constitutive expression of machinery for, 133:1053
role of Ced-3/ICE-family proteases in staurosporine-induced, 133:1041
Prokaryotes, roles of molecular chaperones in protein degradation, 132:255
Proliferation
cellular, regulated by Discs large (Dlg) protein, in *Drosophila* epithelia, 134:1469
of myogenesis, stimulated by IGF-I, 135:431
Proline-rich repeats, in ActA protein, of *Listeria monocytogenes*, control actin-based motility, bacterial motility and VASP and profilin binding, 135:647
Prometaphase, kinetochore movement and response during, 135:315
Proprotein convertase, PC5, COOH-terminal variant isoforms of, biosynthesis of, sorting to different subcellular locations by, 135:1261
Protease content, of mouse mast cells during *Trichinella spiralis* infection, 135:279
Protease receptor, ligand-induced translocation of, caveolae, mechanism for regulating cell surface proteolysis of tissue factor–dependent coagulation pathway, 133:293
Proteases, role of Ced-3/ICE-family proteases in staurosporine-induced programmed cell death, 133:1041
Protein
E3/19K, activation of transcription factor NF-κB by, adenovirus, requires ER retention, 132:511
encoded by Stra8, responsive to retinoic acid, 135:469
34k-Dalton, actin-binding function in *Dictyostelium*, effects on filopodia, 135:965
M_2, ion channel activity of, influenza virus, transport through Golgi apparatus, 133:733
Protein 4.1
hDlg binding sites of, PDZ and I3 domain interactions with, ezrin interactions with, 135:1125

interactions of PDZ domain-proteins with, 135:753
Protein degradation
ER-associated, in vitro assembly is dependent on cytosol, calnexin, and ATP, 132:291
roles of molecular chaperones in, 132:255
Protein export, new pathway for, *Saccharomyces cerevisiae*, 133:1017
Protein folding, chaperonin-mediated folding of actin and tubulin, 132:1
Protein import
of DHFR hybrid protein into glycosomes in vivo not inhibited by folate-analogue aminopterin, 132:311
RAN/TC4 mutants identify common requirement for snRNP and, nucleus, 133:485
repeated cycling between cytoplasm and peroxisomes by Pex5p involved in, 135:1763
Protein kinase
detection of, specifically activated at metaphase-anaphase transition, 132:635
role of glycolipid-based domains, actin cytoskeleton and, endocytosis of GPI-anchored proteins in human lymphocytes, 133:791
Protein kinase A
role in developmental induction of myotome and sclerotome, 134:1241
subunits, in embryonic and skeletal muscle, 134:1241
Protein kinase C
actin-binding motif is unique to epsilon isoform of, participates in regulation of synaptic function, 132:77
atypical, involvement in platelet-derived growth factor stimulation of α_2 gene expression, 134:1301
involved in contractile activity that regulates isoform expression and polysialylation of NCAM in cultured myotubes, Ca^{2+}, 132:969
mitosis-specific phosphorylation of vimentin by, coupled with reorganization of intracellular membranes, 133:141
regulates localization of cytoskeleton proteins and phosphorylation of focal adhesion kinase induced by $\alpha v\beta 5$, 134:1323
Protein kinase C μ (PKCμ), located at Golgi compartment, 134:1401
Protein regulation, integrin-associated, 134:1313
Protein synthesis
constitutive expression of machinery for programmed cell death, 133:1053
mitochondrial, a thermosensitive process, 134:1375
Protein targeting, by tyrosine- and di-leucine-based signals, 135:341
Protein translocation, ER, action of *emo-1* in, 134:699
Protein transmission, of *wingless* in *Drosophila*, role of glycosaminoglycans in, 135:819
Protein transport
chloroplast protein import intermediate reveals distinct translocation machineries in two envelope membranes, 132:63
dynamic localization of nuclear import receptor and its interactions with transport factors, 133:1163
of external Lys-Asp-Glu-Leu protein from plasma membrane to endoplasmic reticulum, KDEL, cholera toxin in vero cells, 133:777
evidence that RAN/TC4 GTPase mediates essential function independent of, nuclear import, green fluorescent protein-glucocorticoid receptor chimera, 133:985
GTPase distinct from Ran involved in, nuclear import, 133:971
targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 132:277
Protein tyrosine kinase, inhibitor, blocks platelet-derived growth factor stimulation of α_2 promoter fragment/CAT construct, 134:1301
Proteins
14-3-3, associate with phosphorylated simple epithelial keratins during cell cycle progression and act as solubility cofactor, 133:345
binding properties of hyaluronic acid effected by variants of CD44, 135:1139
of PDZ domains, structural organization of, interactions of ATP and protein 4.1 with, 135:753
soluble misfolded, receptor-mediated transport to yeast vacuoles of, 135:623
Proteoglycan, GPI-anchored, heparan sulfate expression in polarized epithelial cells, apical sorting of glypican inversely related to heparan sulfate content, 132:487
Proteolysis, mechanism for regulating cell surface, tissue factor–dependent coagulation pathway, ligand-induced protease receptor translocation into caveolae, 133:293
Proteolytic cleavage, TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 132:427
Protofilament formations, of tubulin family, 135:5
Protrusions, in cells transfected with ADP ribosylation factor (ARF), 134:935
PSA-NCAM, interaction of oligodendrocyte-type 2 astrocytes with NMDA and, 135:1565
P-selectin
in PC12 cells, 134:1229
relationship between L-selectin and, 135:523
Pseudocleavage furrow, in syncytial blastoderm, *dah* involved in, 134:923
Pseudopod, formation of, role of phosphoinositide 3-kinase in signaling, 135:1249
Psoriasis, formed by overexpression of bone morphogenetic protein-6 in transgenic mice, 135:227
PTP1B-like tyrosine phosphatase, regulates N-cadherin function by dephosphorylating β-catenin, 134:801
PTS1 receptor
multiple PEX genes required for distribution and stability of, 135:1763
Pex13p as docking factor for, 135:85
Quiescent cells, nuclear proteins of, *Xenopus laevis*, inhibit DNA replication in intact and permeabilized nuclei, 133:955
Rab8, role in fibroblasts of, reorganization of actin filaments and microtubules promoted by, formation of cell protrusions caused by, polarized membrane transport promoted by, 135:153
Rab11, distribution and function of, interaction through pericentriolar recycling compartment of transferrin and, 135:913
Rabbit, nontetrameric species is major soluble form of keratin in reticulocyte lysates of, *Xenopus* oocytes, 132:153
Rabphilin-3A, function in calcium-dependent cortical granule exocytosis at fertilization in mouse eggs, 135:1741
Rac, cell type–specific roles for Cdc42, RhoL and, *Drosophila* oogenesis, 133:617
Rac1, role in site restriction during formation of wing hairs in *Drosophila*, 135:1277
Raji cells, syndecan-1 mediates cell spreading in transfected human, lymphoblastoid cells, 132:1209
Ran
function of, modification of activating protein of, 135:1457
GTPase distinct from, nuclear protein import, 133:971
Ran binding domain (RBD), isolated from Ran binding protein (RanBP1), localizes to cell nucleus, 134:1157
Ran binding protein (RanBP1)
cytoplasmic location of, important to nuclear protein import, 134:1157
nuclear export signal essential for cytosolic localization of, 134:1157
Ran proteins, RAN/TC4 mutants identify common requirement for snRNP and protein import into nucleus, 133:485
RanBP1, effect on p97 interactions in nuclear protein import, 135:559
RanGAP1, identification and localization of novel form of, transport across nuclear pore complex regulated by, 135:1457
Ran/TC4, evidence that RAN/TC4 GTPase mediates essential function independent of nuclear protein import, green fluorescent protein-glucocorticoid receptor chimera, 133:985
ras1, novel suppressor of, fission yeast, dosage-dependent inhibitor of cytokinesis, *byr4*, 133:1307
Ras/Mitogen-activated protein kinase (MAPK), signaling pathway in acetylcholine receptor–inducing activity, 134:465
Rat
kidney cells, role of midzone microtubule bundles in cytokinesis of, 135:981
neurotrophins promote survival and development of neurons in cerebellum of hypothyroid, in vivo, 133:631
rbA, and BI, isoforms of α_{1A}, in synapses and cerebellum, differential localization of, 134:511
RBL-2H3 cells, interface with large Sepharose beads, 134:1427
rDNA enhancer, acts by making more genes transcriptionally active, metazoan, 133:943
rDNA transcription, machinery is assembled during mitosis in active NORs and absent in inactive NORs, 133:235
Reactive oxygen intermediates, UV activates growth factor receptors via, 133:211
Reactive oxygen species, inhibition of IκB-α phosphorylation and degradation and subsequent NF-κB activation by glutathione peroxidase overexpression, 133:1083
Receptor tyrosine kinases, erbB, interaction with heregulin, 134:465
Receptor-like protein tyrosine phosphatase (RPTPμ), does not interact with cadherin complexes, 134:1513
Recognition feature, cytoplasmic, not required for internalization of all cell surface proteins, 134:339
Recycling, role of rab11 through pericentriolar recycling compartment, 135:913

Recycling compartment, specific domain of transferrin receptor causes reorganization of, 135:1749
Reductase, role of N-myristoylation in targeting of, 135:1501
Refolding activity, of Ssa1/2p in yeast cytosol, 135:1229
Regulated secretory organelles, targeting of P-selectin to, in PC12 cells, 134:1229
Regulated secretory pathway, in constitutive secretory cells, 133:1177
Renal failure, resulting from lack of collagen α3(IV), related to Alport syndrome, 135:1403
Replication
of DNA in *Xenopus*, chromosome structure dictates site-specific initiation of, 135:1207
of mtDNA, spatiotemporal constraints on, 135:883
Reporter genes, far-upstream expression enhancer in mouse proα2(I) collagen gene, 134:1333
Repressor activator protein 1 (Rap1), colocalization with telomeres, 134:1349
Respiratory substrates, control by supply of mitochondria with, characterization of autophagy in higher plant cells subjected to carbon deprivation, 133:1251
Reticulocyte lysates, nontetrameric species is major soluble form of keratin in, rabbit, *Xenopus* oocytes, 132:153
Retinoblastoma protein
adhesion-dependent cell cycle progression linked to phosphorylation of, expression of cyclin D1, activation of cyclin E-cdk2, 133:391
defects in MHC expression specific to lack of, 135:441
Retinoic acid
Bcl-2 inhibits apoptosis induced by, neural differentiation of embryonal stem cells, 132:955
characterization of Stra8-encoded protein responsive to, 135:469
Retinoic acid receptor-β2 (RARβ2) gene, tissue-specific expression of, in mouse embryos, regulated at translational level by short upstream open reading frames in 5′-untranslated region, 134:827
Retinoid, synthetic CD437 activates and regulates AP-1, inhibits growth and induces apoptosis in human melanoma cells, 135:1889
Retinoid-dependent pathways, crucial role in regulation of brain and heart development, and myocardial differentiation, 134:827
Retrograde transport, localization of yeast early Golgi mannosyltransferase involves, Och1p, 132:985
Rev protein, facilitation of nuclear export of HIV-1 RNA by, 135:9
RGD sequence, in matrix assembly, 134:573
RGD/synergy site, required for embryonic cell migration on fibronectin, 134:227
Rho
contractility stimulated by, drives formation of stress fibers and focal adhesions, 133:1403
involvement in regulation of CD44/ERM interaction, 135:37
role in actin cable assembly during wound healing of, 135:1097
rho family, novel member of, small GTP-binding proteins required for cytokinesis, 133:1321
RhoL, cell type–specific roles for Cdc42, Rac and, *Drosophila* oogenesis, 133:617
Rho-type GTPases, associations among PH, SH3 domain–containing proteins and, yeast, 133:879
RIα transcripts, localize to neuromuscular junction, 134:1241
Ribosomes, native 3D structure of eukaryotic 80S ribosome, morphological homology with *E. coli* 70S ribosome, 133:495
Ring canals, formation of actin filament bundles in, developing *Drosophila* follicles, 133:61
Ring formations, of tubulin family, model for microtubule nucleation, 135:5
RNA
alterations in trafficking of, produced by Nup153 NP-domain, 134:1141
of HIV-1, distribution and spatial relationships of, exported by Rev protein, 135:9
localization in specific nuclear domain of snRNP and, localization and export of spliced exons of viral, localization of polyadenylated, 135:303
RNA polymerase I
in vivo evidence that TATA-binding protein/SL1 colocalizes with UBF and, rRNA synthesis is either active or inactive, 133:225
rDNA transcription machinery is assembled during mitosis in active NORs and absent in inactive NORs, 133:235
Rna1p, dynamic localization of nuclear import receptor and its interactions with transport factors, 133.1163
RNP, heterogeneous nuclear (hnRNP) proteins, nuclear export signal–bearing, mediate mRNA export, 134:1365
Rolling, cell body, in coordination of protrusion and translocation of keratocyte, 134:1209
Rolling adhesions
effect of interactions of leukocytes through L-selectin on, 135:849
mediated by L-selectin ligands on leukocytes in shear flow, 135:837
Rotation, of cytoplasm during locomotion, within keratocyte cell body, 134:1209
rRNA synthesis, in vivo evidence that TATA-binding protein/SL1 colocalizes with UBF and RNA polymerase I during, active or inactive, 133:225
Ryanodine receptor
clusters in dysgenic microtubules, 134:375
in excitation–contraction coupling of *Caenorhabditis elegans* striated muscle, 134:885
80S monosome, native 3D structure of eukaryotic 80S ribosome, morphological homology with *E. coli* 70S ribosome, 133:495
Saccharomyces
characterization of mutants defective in Pea2p and Spa2p, polarized growth and budding requirements in, 135:725
effects of Ste6 on **a**-factor in cell fusion during mating of, 135:741
function of myosin and actin in vacuole inheritance in, 135:1535
MHP1 necessary in formation and stabilization of microtubules in, 135:1323
new clathrin-dependent endocytosis signal with NPFXD sequence in, 135:1789
receptor-mediated transport of soluble misfolded proteins to vacuoles in, 135:623
temporal and spatial characterization of Chs2p and Chs3p in, 135:597
Saccharomyces cerevisiae
actin-related protein ARP2, in actin cytoskeleton, involved in membrane growth, polarity, and endocytosis, 134:117
BED1 required for polarized growth and efficient bud emergence in, gene encoding galactosyltransferase homologue, 132:137
budding patterns of, 134:413
α-factor receptor (Ste3p) in, subject to phosphorylation and ubiquitination, 134:661
identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 132:999
identification of developmentally regulated septin and involvement of septins in spore formation in, 132:399
localization of yeast early Golgi mannosyltransferase involves retrograde transport, Och1p, 132:985
neck filament assembly and septin in, 134:1345
new pathway for protein export in, 133:1017
PAL1 gene product is peroxisomal ATP-binding cassette transporter in, 132:549
Pds1p plays critical role in APC and checkpoint pathways in, inhibitor of anaphase in budding yeast, 133:99
Pds1p required for execution of anaphase in, 133:85
role of calmodulin and Spc110p interaction in proper assembly of spindle pole body components, 133:111
signal recognition particle (SRP)–independent preprotein pathway in, 134:269
Spc42p is phosphorylated component of spindle pole body in, essential function during SPB duplication, 132:887
spindle pole body component in, phosphoprotein is modified in cell cycle-dependent manner, 132:903
spindle pole body duplication gene MPS1 is part of mitotic checkpoint in, 132:111
synthetic lethality screen identifies novel myosin I gene in, *MYO5*, myosin I proteins required for polarization of actin cytoskeleton, 133:1277
thioredoxin required for vacuole inheritance in, 132:787
TUB4 gene of, 134:443
ubiquitin-like genes involved in duplication of microtubule organizing center in, 133:1331
Vps10p cycles between late-Golgi and prevacuolar compartments in its function as sorting receptor for multiple vacuolar hydrolases in, 133:529
S-Acylation, in plasma membrane and in Golgi apparatus, mediated by distinct S-acyltransferases, 134:647
S-Acyltransferase, localizes to plasma membrane, 134:647
α-satellite, in DNA domain of centromeres, elastic behavior of, 135:545
Scaffold, centromeric topoisomerase II α in, 134:1097
Schizosaccharomyces pombe
mitotic timing determined by dephosphorylation of inhibitory sites in CDC2, 134:963
novel suppressor of *ras1* in, dosage-dependent inhibitor of cytokinesis, *byr4*, 133:1307
Schwann cells, granulocyte macrophage colony stimulating factor produced in lesioned peripheral nerves induces upregulation of cell surface expression of MAC-2 by macrophages and, 133:159

SDYQRL motif, role in targeting to TGN and in determining physical characteristics of endosomal compartments, 135:1749
Sea urchins
embryo, increases in calcium near nucleus are required to trigger entry into mitosis in, 135:191
lamin B receptor-like integral membrane protein mediates membrane targeting to chromatin in, 135:1715
sperm receptor in, egg jelly, modular protein with extensive homology to human polycystic kidney disease protein, PKD1, 133:809
Sec12p, two mechanisms of ER localization, 134:279
SEC62, *SEC63*, impair translocation of signal recognition particle–independent preproteins, 134:269
Sec63p, ER membrane protein complex required for nuclear fusion, 132:499
Secretion
biosynthetic regulated secretory pathway in constitutive secretory cells, 133:1177
of catecholamine requires translocation of p36 to subplasmalemmal region in chromaffin cells, annexin II in exocytosis, 133:1217
Secretory carrier-associated membrane protein (SCAMP), in ubiquitous secretory and endocytotic pathways, 134:731
Secretory granules, in *Tetrahymena*, role of granule lattice protein 1 to effect function and composition of, 135:1775
Secretory pathway, new pathway for protein export in *Saccharomyces cerevisiae*, 133:1017
Secretory proteins, TRAM required for translocation of, 134:25
Secretory vesicles, from TGN of endocrine cells, tyrosine phosphatase and tyrosine kinase are required for formation of, 135:1471
Semliki Forest virus, infects cells by acid-dependent membrane fusion reaction catalyzed by virus spike protein, 134:863
Septate junction, Discs large (Dlg) protein, critical component of, 134:1469
Septins
Drosophila complex with, forms filaments and exhibits GTPase activity, 133:605
essential for neck filament assembly, 134:1345
involvement in spore formation in *Saccharomyces cerevisiae*, identification of developmentally regulated septin, 132:399
Sequence NPFXD, new clathrin-dependent endocytosis signal in *Saccharomyces* with, 135:1789
Serpentine receptors, in adhesion and chemotaxis, 134:255
Serum starvation, of cathepsin B, induces expression in myoblasts, but downregulates it in myotubes, 134:837
SH3, conserved domain of Discs large (Dlg) protein, in regulation and maintenance of septate junctions, 134:1469
SH3 domain
cytoplasmic orientation of, essential for Pex13p function, 135:85
of Pex13p, role in interaction between Pex13p and Pex5p, 135:111
role in Pex13p function, interaction of Pex5p with Pex13p in, 135:97
Shc, adaptor molecule, recruitment to ligated $\alpha_6\beta_4$ partially suppressed by epidermal growth factor, 134:241
Shedding, novel mechanism in early steps of CD95-mediated apoptosis, 134:1089
She4p, endocytosis role of, identified by novel genetic screen, 135:1485
Shigella flexneri, two classes of oligoproline sequences in profilin-mediated acceleration of actin-based motility in, 133:49
SHL tripeptide, internal, not directly involved in peroxisomal targeting, 134:849
Shr3p, on endoplasmic reticulum, packaging amino acid permeases into transport vesicles by COPII and, 135:585
Signal anchor, positively charged domain of, in yeast Mas70p and NADH cytochrome b_5, functions independently as matrix-targeting signal in yeast mitochondria in vitro, 134:307
Signal recognition particle (SRP)–independent preproteins, mutant alleles of *SEC62* and *SEC63* impair translocation of, 134:269
Signal sequence
structure of, determines whether TRAM-dependent or -independent, 134:25
TRAM dependence of secretory proteins determined by, 134:25
translocation pathway use determined by, 134:269
Signal transduction
β1D integrin displaces β1A isoform in striated muscles, localization at junctional structures and signaling potential in nonmuscle cells, 132:211
deficiency of Src family kinases p59/61^{hck} and p58$^{c\text{-}fgr}$ results in defective adhesion-dependent neutrophil functions, 133:895
gene *unc-89* encodes giant modular protein composed of domains of Ig and, required for muscle M-line assembly, *Caenorhabditus elegans*, 132:835
by polymeric immunoglobulin receptor suggests role in regulation of receptor transcytosis, 133:997
Signaling
activities of *Wnt*-1 class of secreted factors of, antagonized by *Wnt*-5A class and by dominant negative cadherin in early *Xenopus* development, 133:1123
binding to cadherins antagonizes, β-catenin, axis formation in *Xenopus*, 132:1105
contact-mediated, role of receptor-like protein tyrosine phosphatase (RPTPμ) in, 134:1513
endocytosis of VAMP facilitated by synaptic vesicle targeting signal, 132:537
ion channel subunit required for mechanotransduction in *Caenorhabditis elegans*, sequence and transmembrane topology of MEC-4, 133:1071
transport of vesicular stomatitis virus G protein to cell surface is mediated by, polarized and nonpolarized cells, 133:543
Signaling pathway, involving MAP kinase, angiotensin II activation of, 135:1609
Signals, nuclear export
mediate nuclear export of shuttling heterogeneous nuclear RNP (hnRNP) proteins, 134:1365
in Ran binding protein (RanBP1), 134:1157
Silencing, telomere proximal, repressor activator protein (Rap1), Sir3p, and Sir4p essential for, 134:1349
Simian immunodeficiency virus, internalization signal in, transmembrane protein cytoplasmic domain, modulates expression of envelope glycoproteins on cell surface, 132:795
Single-base 3′ overhangs, present in DNA breaks during apoptosis, 135:1369
Sir3p
colocalization with telomeres, 134:1349
phosphorylation of, in transcriptional silencing regulation, activated by MAP kinase, 135:571
Sir4p, colocalization with telomeres, 134:1349
Skeletal muscle
differentially expressed fibroblast growth factors regulate development of, autocrine and paracrine mechanisms, 132:1151
functions of α_4 integrin in, 135:829
identification of small cytoplasmic ankyrin in kidney and, binds βIΣ* spectrin and associates with Golgi apparatus, Ank_{G119}, 133:819
insulin-stimulated transport mediated by Glut4 in transverse tubules in, 135:415
interaction of F-actin with dystrophin, 135:661
three-dimensional structure of Z band in normal mammalian, 133:571
Skeletal muscle cells, lack of retinoblastoma protein on, accumulation in S and G_2 phases of, 135:441
Skin
inactivation of integrin β6 subunit gene reveals role of epithelial integrins in regulating inflammation in lungs and, 133:921
re-epithelialization after injury to, correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 132:381
severe fragility of, changes of cytokeratin expression in epidermis, gene targeting at mouse cytokeratin 10 locus, 132:925
SKW3 cells, binding with stromal matrix, 134:1075
SM22α, expression of promoter of in transgenic mice, distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells, 132:849
Smooth muscle cells
distinct transcriptional regulatory programs in vascular and visceral, expression of *SM22α* promoter in transgenic mice, 132:849
expression of smoothelin in, 134:401
Smooth muscles, contractibility, importance of side-polar structure of crossbridges to, 134:53
Smoothelin, human, encoded by a single copy gene on chromosome 22, 134:401
SNAP receptor (SNARE) mechanism, works for membrane expansion in growth cone, 134:205
SNAREs, v-SNARE implicated in intra-Golgi transport, 133:507
snRNP
localization in specific nuclear domain of RNA and, 135:303
RAN/TC4 mutants identify common requirement for import of, protein import into nucleus, 133:485
Sodium channels
controlling density of at plasma membrane, regulates neuronal excitability, 134:499
internalization and degradation, 134:499
Soluble misfolded proteins, receptor-mediated transport to yeast vacuoles of, 135:623
Soluble *N*-ethylmaleimide-sensitive fusion attachment protein receptor, assembly of ER to Golgi SNARE complex requires Uso1p, 132:755

Somatomedin B (SMB) domain, binding site for urokinase type plasminogen activator receptor in vitronectin, 134:1563
Sorting
apical sorting of glypican (GPI-anchored proteoglycan) inversely related to heparan sulfate content, heparan sulfate expression in polarized epithelial cells, 132:487
of COOH-terminal variant isoforms of proprotein convertase to different subcellular locations, 135:1261
of newly synthesized galactosphingolipids to two surface domains of epithelial cells, 132:813
targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 132:565
of transferrin receptors, by apical and basolateral endosomes of MDCK cells, 135:139
Sox17 gene, identification of two Sox17 messenger RNA isoforms, with and without high mobility group box region, differential expression in mouse spermatogenesis, 133:667
Spa2p, in *Saccharomyces*, characterization of mutants defective in Pea2p and, role in polarized growth, mating and budding of, 135:725
SPC4, SPC6, and SPC7, bone morphogenetic proteins, coexpressed with, at distinct sites during embryogenesis, 134:181
SPC7, novel protease, expressed ubiquitously throughout embryonic development, 134:181
Spc42p, phosphorylated component of spindle pole body with essential function during SPB duplication, *Saccharomyces cerevisiae*, 132:887
Spc110p, role of interaction of calmodulin and, proper assembly of spindle pole body components, 133:111
Spectrin
dynactin associations mediated by interaction between centractin and, 135:1815
identification of small cytoplasmic ankyrin that binds βIΣ* spectrin, kidney and muscle, associates with Golgi apparatus, Ank_{G119}, 133:819
Sperm membrane potential, mouse and bovine, depolarized during contact with zona pellucida, by ZP3-dependent activation of pertussis toxin–insensitive pathway with characteristics of poorly selected cation channel, 134:637
Sperm receptor, for egg jelly is modular protein with extensive homology to human polycystic kidney disease protein, sea urchin, PKD1, 133:809
Spermatogenesis
identification of two Sox17 messenger RNA isoforms and their differential expression in, mouse, with and without high mobility group box region, 133:667
role of Stra8-encoded protein in premeiotic, 135:469
S-phase, camptothecin toxicity mediated in, 134:757
Sphingolipid galactosyltransferases, topology of, ER and Golgi, transbilayer movement of monohexosyl sphingolipids required for higher glycosphingolipid biosynthesis, 133:15
Sphingolipids, sorting of, in chlamydia-infected cells, 134:363
Spike protein, Semliki Forest virus, mediates fusion, 134:863
Spindle, aberrantly segregating centromeres activate spindle assembly checkpoint in budding yeast, 133:75
Spindle assembly, in *Drosophila* female meiosis, 134:455
Spindle elongation, DSK1 involved in, anaphase, novel kinesin-related protein from diatom *Cylindrotheca fusiformis*, 133:595
Spindle formation
mitotic, 134:429
in mitotic human cells, polo-like kinase1 required for, 135:1701
Spindle morphogenesis, implications of microtubule dynamics during interphase/mitosis transition for, 135:201
Spindle pole body
duplication gene MPS1 of, part of mitotic checkpoint, *Saccharomyces cerevisiae*, 132:111
110-kD component of, phosphoprotein is modified in cell cycle-dependent manner, *Saccharomyces cerevisiae*, 132:903
role of calmodulin and Spc110p interaction in proper assembly of components of, 133:111
in *Saccharomyces cerevisiae*, microtubule organizing functions provided by, 134:429
Spc42p is phosphorylated component of, essential function during SPB duplication, *Saccharomyces cerevisiae* 887,
Tub4p associated dynamically with, 134:443
yeast ubiquitin-like genes involved in duplication of microtubule organizing center, 133:1331
Spindle tension, influences ZW10 association with individual chromosomes, 134:1127
Spindles, molecular characterization of 50-kD subunit of dynactin reveals function for complex in chromosome alignment and organization of, mitosis, 132:617
Splicing factors
dynamic transcription-dependent organization of pre-mRNA, polytene nuclei, 133:929
intron-dependent recruitment of pre-mRNA, sites of transcription, 133:719
Spore formation, identification of developmentally regulated septin and involvement of septins in, *Saccharomyces cerevisiae*, 132:399
Spreading, intracellular pH regulation during, human neutrophils, 133:1391
src homology 4 domain, features of $p59^{fyn}$, define motif for immune-receptor tyrosine-based activation motif binding and for plasma membrane localization, 133:1007
Src homology 3 ligand domains, MDC9 is widely expressed cellular disintegrin containing, cytoplasmic, 132:717
src homology region 3 domain-binding proteins
associations among PH domain, Rho-type GTPases and, yeast, 133:879
involved in bud formation, yeast, 133:865
Src family kinases, deficiency of Src family kinases $p59/61^{hck}$ and $p58^{c\text{-}fgr}$ results in defective adhesion-dependent neutrophil functions, 133:895
Src kinase, actin cytoskeleton mediates translocation to cell periphery of, controlled by G proteins, 135:1551
Ssa1/2p, refolding activity of, in yeast cytosol, protein translocation role of, 135:1229
Ssc1, increased expression of, partially suppresses cold-sensitive growth defect of *SSH1* mutant, 134:603
Ssh1, required for normal mitochondrial DNA replication, 134:603
SSH1 gene, deletion of, results in extremely slow growth at 23° or 30°C, but nearly wild-type growth at 37°C, 134:603
stardust, zonula adherens formation, and maintenance of polarized blastoderm epithelium in *Drosophila*, 134:149
Starfish, triggering of M-phase in oocytes of, novel component of maturation-promoting factor besides cdc2 kinase, 132:125
Stathmin, is phosphoprotein essential for nerve growth factor–stimulated differentiation, 133:1383
Staurosporine
constitutive expression of machinery for programmed cell death, 133:1053
role of Ced-3/ICE-family proteases in programmed cell death induced by, 133:1041
Ste6, effects on a-factor in cell fusion during mating of *Saccharomyces*, 135:741
stmF, *Dictyostelium* streamer F mutant, produces prolonged peak of cGMP accumulation upon cAMP stimulation, 134:911
Stra8, characterization of protein responsive to retinoic acid encoded by, role in premeiotic spermatogenesis of, 135:469
Stress fibers
Rho-stimulated contractility drives formation of focal adhesions and, 133:1403
α-smooth muscle actin expression in, correlates with focal contact formation, 134:67
Striatin, member of WD-repeat family, novel calmodulin-binding protein, 134:1051
Striatum, dorsal part of, striatin conspicuous in, 134:1051
Stroma, human amnion contains novel laminin variant, laminin 7, like laminin 6 is covalently associated with laminin 5 to promote stable epithelial–stromal attachment, 132:1189
Stromal cell surface molecules, interaction with pre-B cells, 134:771
Stromal interaction molecule, 4-9, in orientation of stromal cells within tissues, 134:771
Stromelysin-1, in mammary epithelial cells of mice, effects of crossing TIMP-1 with, 135:1669
Strongylocentrotus purpuratus, sequence and localization of 115-kD accessory subunit of heterotrimeric kinesin-II complex, $KRP_{85/95}$, 132:371
Structure–function correlations, of COOH-terminal domain of dystrophin, 134:93
α subunit, of acetylcholine receptor, formation of heterodimeric binding site by, effect of tethering on, 135:809
Sulfation, HeLa protein kinase C μ (PKCμ) transfectants show enhance rate of, 134:1401
Sulfhydryl region, reactive, importance of motion in, 134:895
sup-pf-2 mutations, in *Chlamydomonas*, activity of outer dynein arms altered by, γ-dynein heavy chain modified by, 135:1865
Surface structure, of active and inactive human X interphase chromosome territories, three-dimensional analysis of, 135:1427

Sympathetic neurons, in rat, velocity and magnitude of tubulin transport in, 135: 701
Symplekin, novel tight junction plaque protein, 134:1003
Synapses, calcium-triggered exocytosis at, 134:329
Synapsin I, knock-out mice, vesicle recycling in, 134:1219
Synapsins, in regulation of neurotransmitter release, 134:1219
Synapsis initiation, telomere movements toward cluster site precede, 134:1109
Synaptic junctions, adhesion of cadherin-associated catenin proteins in, 135:767
Synaptic vesicles
actin-binding motif participates in function of, epsilon isoform of protein kinase C, 132:77
cycle of, single vesicle budding step involving clathrin and dynamin, 133:1237
endocytosis of VAMP facilitated by targeting signal of, 132:537
recycling pathways of, in active and nonactive zones, 135:797
Synaptic-like micro-vesicles, targeting signals to, 134:1229
Synaptobrevin/vesicle-associated membrane protein, intracellular distribution of, with respect to glucose transporter (GLUT-4), 134:625
Synaptonemal complex, pairing of homologous chromosomes during meiotic prophase culminates in formation of, 134:1109
Synaptophysin, homologue of, 134:731
Syndecan-1, mediates cell spreading in transfected human lymphoblastoid cells, Raji cells, 132:1209
Syndecans, stimulation of fibroblast growth factor receptor-1 occupancy and signaling by cell surface–associated glypican and, 133:405
Synergy site, PPSRN in type III-9 repeat, cell recognition of, positionally regulated in early embryo, 134:227
Syntaxin, in growth cone, acts through SNAP receptor (SNARE) mechanism, 134:205
Synthetic lethality screen, identifies novel yeast myosin I gene, *MYO5*, myosin I proteins required for polarization of actin cytoskeleton, 133:1277
T cell receptor, role of CD3γ in assembly of, 132:299
T cells, fucosyltransferase FucT-VII regulates E-selectin ligand synthesis in human, 133:911
Talin, in filopodial motility, 134:1197
Targeting
across ER membrane, catalyzed by signal recognition particle (SRP) and SRP receptor, 134:269
and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 132:277
identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 132:999
of NADH-cytochrome b_5 reductase, role of N-myristoylation in, 135:1501
Targeting signal, sorting sequence of peroxisomal integral membrane protein PMP47 is contained within short hydrophilic loop, 133:269
TATA-binding protein, in vivo evidence that TATA-binding protein/SL1 colocalizes with UBF and RNA polymerase I when rRNA synthesis is either active or inactive, 133:225
Tau, selective stabilization in axons, microtubule-associated protein 2C in cell bodies and dendrites, polarized localization of cytoskeletal proteins in mature neurons, 132:667
Taxol, G1 arrest in transformed and nontransformed cells caused by, microtubule sensitivity to, 135:689
TCR, features of $p59^{fyn}$ src homology 4 domain define motif for immune-receptor tyrosine-based activation motif binding and for plasma membrane localization, 133:1007
Telomeres
clustering and colocalization with Rap1, Sir3, and Sir4 proteins, in wild-type *Saccharomyces cerevisiae*, 134:1349
distribution patterns in mouse and human meiocytes, 134:1109
mammalian, interaction of TRF with DNA of, 135:867
size, changes in, during replicative aging and cell transformation, 134:1
Temperature, reduced, and cell adhesion, 134:549
Temperature-sensitive mutants, role of calmodulin and Spc110p interaction in proper assembly of spindle pole body components, 133:111
Tenascin-C
contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons, 132:681
dynamic expression in developing organs, 134:1499
Tenascin-R, expression restricted to central nervous system, 134:1499
Tenascin-Y, expressed and secreted by cells within muscle connective tissue, 134: 1499
Terminal COOH, truncation mutants of, of fusion protein, importance in various stages of fusion, 135:73
Tethering
effect on acetylcholine receptor α subunit in formation of heterodimeric binding site, 135:809
mediated by L-selectin ligands on leukocytes in shear flow, 135:837
Tetrahymena
conjugation of, phosphorylation of linker histone δ in, 135:1219
role of granule lattice protein 1 to effect function and composition of secretory granules in, 135:1775
TGN, role of specific domain of transferrin receptor in targeting to, 135:1749
Thermotolerance, mitochondrial, molecular chaperone Hsp78 crucial in, 134:1375
Thiolase, Peb1p is intraperoxisomal receptor for NH_2-terminal, type 2, peroxisomal targeting sequence of, Peb1p targeted to peroxisomes by NH_2-terminal peptide, Pas7p, 132:325
Thioredoxin, required for vacuole inheritance in *Saccharomyces cerevisiae*, 132: 787
Three-dimensional analysis, comparison of active and inactive human X interphase chromosome territories using, 135:1427
Thrombospondin, effects on vitronectin of melanoma cells through integrin-associated protein, 135:533
Tight junctions
antiglycolipid antibody interferes with basolateral polarization and formation of, inhibits Madin-Darby canine kidney cell adhesion to laminin, 133:695
catenins and Zonula Occludens-1 form complex during early stages in assembly of, 132:451
interspecies diversity of occludin sequence, cDNA cloning of human, mouse, dog, and rat–kangaroo homologues, 133:43
permeability of, 134:1031
symplekin in, 134:1003
TIMP-1, in mammary epithelial cells of mice, effects of crossing stromelysin-1 with, 135:1669
Tissue distribution, Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues, 132:132:167
Tissue factor, mechanism for regulating cell surface proteolysis of coagulation pathway dependent on, ligand-induced protease receptor translocation into caveolae, 133:293
Titin, association with myomesin and M-protein, 134:1441
Tonsillar stromal cells, rolling interactions of lymphocytes on, 134:1075
Topoisomerase-I, DNA, inhibitors of, induce apoptosis independent of DNA synthesis, 134:757
Topoisomerase II α, required for proper centromere/kinetochore structure, 134:1097
Touch dome mechanoreceptor units, enlarged by elevated NT-3, 134:487
Trafficking, polarity, detergent solubility and, glycosylphosphatidylinositol–anchored protein after LDL-deprivation of MDCK cells, 133:1265
Transcription
of α1(VI) collagen chain, in transgenic mice, specific regions involved in regulation of, 135:1163
dynamic organization of pre-mRNA splicing factors dependent on, polytene nuclei, 133:929
metazoan rDNA enhancer acts by making more genes active in, 133:943
micronuclei during *Tetrahymena* conjugation, associated with phosphorylation of linker histone δ, 135:1219
Transcription factor NF-κB, activated by adenovirus E3/19K protein, requires ER retention, 132:511
Transcription sites, intron-dependent recruitment of pre-mRNA splicing factors to, 133:719
Transcriptional silencing, regulation of, by phosphorylation of Sir3p activated by MAP kinase, 135:571
Transcytosis, signal transduction by polymeric immunoglobulin receptor suggests role in regulation of receptor, 133:997
Transepithelial electrical resistance, expression of both wild-type and mutant occludin induces, 134:1031
Transferrin, interaction through pericentriolar recycling compartment of rab11 and, 135:913
Transferrin receptors
polarized sorting mechanism in MDCK cells of, 135:139
specific domain of, reorganization of recycling compartment caused by, 135:1749
Transforming growth factor α, population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF, and, chemically defined (HGM) medium, 132:1133
Transforming growth factor-β, endoglin modulates cellular responses to TGF-β1, 133:1109

Transforming growth factor–β2, expression of in osteoblasts results in osteoporosis-like phenotype, 132:195
Transglutaminase, membrane-bound, keratinocyte-specific, in lamellar ichthyosis, 134:715
trans-Golgi network
biogenesis of MHC class II compartment in human I-Cell Disease B lymphoblasts, 132:769
different biosynthetic transport routes to plasma membrane in BHK and CHO cells, 133:247
of endocrine cells, tyrosine phosphatase and tyrosine kinase are required for formation of secretory vesicles from, 135:1471
mediation of transport of newly synthesized soluble lysosomal enzymes from, by two distinct mannose 6-phosphate receptors, 134:615
novel adaptor-related protein complex, 133:749
sorting of newly synthesized galactosphingolipids to two surface domains of epithelial cells, 132:813
Transit sequence, IAP86 and IAP75, components of multisubunit complex mediating energy-independent recognition of, 134:315
Translational activation, of CUG-initiated FGF-2 for cell transformation and stress response, 135:1391
Translational regulation, tissue-specific, 134:827
Translocating chain–associating membrane (TRAM), in cotranslational translocation of proteins across mammalian ER membrane, 134:25
Translocation
early-phase TRAM protein in, 134:25
role of yeast heat shock proteins Ssa1/2 in, 135:1229
of Src kinase to cell periphery, mediated by actin cytoskeleton, controlled by G proteins, 135:1551
Transmembrane domain
required for Rer1p-dependent retrieval in ER localization of Sec12p, 134:279
role of CD3γ in T cell receptor assembly, 132:299
Transmembrane ligands, mutations in *Drosophila hook* gene inhibit endocytosis of boss transmembrane ligand into multivesicular bodies, 133:1205
Transmembrane protein, internalization signal in cytoplasmic domain of, simian immunodeficiency virus, modulates expression of envelope glycoproteins on cell surface, 132:795
Transport
across nuclear pore complex, identification and localization of novel form of RanGAP1 involved with regulation of, 135:1457
forward, role of COP I vesicles in, 134:1411
of MHC class II molecules, from lysosomal structures to plasma membrane, 135:611
of neurofilaments, effects of NF-H on, 135:711
receptor-mediated, of soluble misfolded proteins to yeast vacuoles, 135:623
retrograde, role of COP I vesicles in, 134:1411
velocity and magnitude of tubulin, in rat sympathetic neurons, 135:701
Transport factors, dynamic localization of nuclear import receptor and its interactions with, 133:1163
Transport vesicles
constitutive, defined by pantophysin, 134:731
roles of COPII and Shr3p in packaging amino acid permeases into, 135:585
transport of vesicular stomatitis virus G protein to cell surface is signal mediated in polarized and nonpolarized cells, 133:543
yeast Sec12p in formation of, 134:279
TRF, interaction between mammalian telomeric DNA and, 135:867
Triad formation, in dysgenic (*mdg/mdg*) line cells, 134:375
Triton X-100, endocytosis of GPI-linked membrane folate receptor-α, 132:35
TrkA receptor, TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 132:427
Tropomyosin, mutants in *Drosophila* flight muscle, 135:673
Troponin T, mutations of in striated and nonstriated muscle cells of *Caenorhabditis elegans*, 132:1061
Truncation, mutants of COOH-terminal region of fusion protein, 135:73
Trypanosoma cruzi, invasion of mammalian cells by, occurs by actin-independent mechanism distinct from phagocytosis, 134:349
Trypanosomes
host cell invasion by, requires lysosomes and microtubule/kinesin-mediated transport, 134:349
mitochondria of, have more than one hsp70, 134:603
Trypsin, treatment of surface of intact mitochondria with, for removal of proximal components of outer membrane import machinery, 134:307
TUB4, in microtubule-organizing center function, 134:443
Tub4p, novel tubulin in *Saccharomyces cerevisiae*, role in microtubule organization, 134:429
Tub4p function, perturbation of, results in spindle and spindle body defects, 134:443
Tubulin
active transport of photoactivated molecules of, growing axons, revealed by new electron microscopic analysis, 133:1347
chaperonin-mediated folding of actin and, 132:1
α-, repression of synthesis of, mechanism of translational inhibition of, effect of colchicine on, 135:1525
transport in neurons, 133:1355
velocity and magnitude of transport of, in rat sympathetic neurons, 135:701
Tubulin family, protofilament and ring formations of all members of, 135:5
Tumor necrosis factor, chloride ion efflux induced by, 135:511
Tumor necrosis factor-α, cooperative effect of bFGF, VEGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 132:1177
Tumor suppression, by plakoglobin, augmenting effect of N-cadherin, 133:199
Tumor suppressor gene, role of p53 to induce postmitotic neuronal apoptosis, 135:1085
Tumorigenesis
and epidermal growth factor–induced changes in $\alpha_6\beta_4$ function, 134:241
in squamous epithelium, inadvertent expression of nonepithelial cadherin causes, 135:1643
Type III repeats, first, deletion of, alter rate of fibronectin matrix assembly, 134:573
Type V collagen, in chicken cornea, role in regulation of fibril diameter, 135:1415
Tyrosinase, augmentation of, by cAMP elevating agents, stimulates melanogenesis, 134:747
Tyrosine
agrin-induced acetylcholine receptor clustering in mammalian muscle requires phosphorylation of, 132:937
anchorage mediated by integrin α6β4 to laminin 5 regulates phosphorylation of, epiligrin, membrane-associated protein, 132:727
phosphatase and kinase, formation of secretory vesicles from TGN of endocrine cells require, 135:1471
stimulation of β_1 integrins on fibroblasts induces PDGF independent phosphorylation of, PDGF β-receptors, 132:741
targeting of LAMP1 to lysomes is dependent on spacing of cytoplasmic tail tyrosine sorting motif relative to membrane, 132:565
TrkA receptor ectodomain cleavage generates tyrosine-phosphorylated cell-associated fragment, 132:427
Tyrosine kinase
activity increased in nuclear fractions with translocation of fibroblast growth factor receptor to nucleus, 134:529
Lck, associated with glycolipid-enriched membrane domains, relationship to CD45 of, 135:1515
$pp125^{FAK}$, functions to suppress apoptosis in fibroblasts, 135:1383
role of Fcγ receptors and, N^+/H^+ exchange activity during phagocytosis in human neutrophils, 132:1037
TrkA, neurotrophin receptor forms complex with, gp75, 132:945
Tyrosine kinase receptors, motogenic and morphogenic activity of epithelial, 133:1095
Tyrosine phosphatases, transmembrane, LAR subfamily of, cadherin–catenin complex associated with, 134:1519
Tyrosine phosphorylation
of Cek9, 135:781
fibroblast growth factor receptor-1 substrates, 134:783
Rho-stimulated contractility drives formation of stress fibers and focal adhesions, 133:1403
Tyrosine-based signals, protein targeting mediated by, distinct saturable components of targeting mechanisms of di-leucine-based and, 135:341
U937 cells, binding of vitronectin to, through somatomedin B domain, 134:1563
Ubiquitin, yeast ubiquitin-like genes involved in duplication of microtubule organizing center, 133:1331
Ubiquitination
of α-factor receptor (Ste3p), 134:661
receptor, decrease in, correlates with impairment of receptor degradative turnover in mutant strain of α-factor receptor, 134:661
Ultraviolet radiation, activates growth factor receptors via reactive oxygen intermediates, 133:211
unc-7, mutations in, severely impair forward locomotion, 134:537

unc-68, involvement in *Caenorhabditis elegans* body-wall muscle contraction, 134:885
unc-89 gene, encodes giant modular protein composed of Ig and signal transduction domains, *Caenorhabditus elegans*, required for muscle M-line assembly, 132:835
Upstream binding factor, in vivo evidence that TATA-binding protein/SL1 colocalizes with RNA polymerase I and, rRNA synthesis is either active or inactive, 133:225
Upstream open reading frame (uORF)4, amino acid coding sequence, inhibits effect on translation of downstream major open reading frame, 134:827
Urokinase, role of, cooperative effect of TNFα, bFGF, and VEGF on formation of tubular structures of human microvascular endothelial cells in fibrin matrix, 132:1177
Uso1p, assembly of ER to Golgi SNARE complex requires, 132:755
Vacuolar protein sorting protein 10, cycles between late-Golgi and prevacuolar compartments in its function as sorting receptor for multiple yeast vacuolar hydrolases, 133:529
Vacuolating cytotoxin A, oligomeric and subunit structure of *Helicobacter pylori* vacuolating cytotoxin, 133:801
Vacuoles
  identification of cytoplasm to vacuole targeting determinant in aminopeptidase I, 132:999
  in *Saccharomyces*, function of myosin and actin in inheritance of, 135:1535
  thioredoxin required for inheritance of, *Saccharomyces cerevisiae*, 132:787
  of yeast, receptor-mediated transport of soluble misfolded proteins to, 135:623
VAMP, endocytosis of, facilitated by synaptic vesicle targeting signal, 132:537
Vascular cell adhesion molecule (VCAM)-1, effect of CC chemokines on monocyte adhesion to, 134:1063
Vascular endothelial cell growth factor, cooperative effect of TNFα, bFGF, and, formation of tubular structures of human microvascular endothelial cells in fibrin matrix, role of urokinase activity, 132:1177
Vasodilator-stimulated phosphoprotein, binding of, controlled by proline-rich repeats in ActA protein of *Listeria monocytogenes*, 135:647
V-ATPase, Golgi maintained at acidic level by, 134:1387
Ventricular zone gene-1, in neurogenic regions of cerebral cortex, role in LPA signaling, 135:1071
Vero cells, internalization of FITC-beta subunit of verotoxin (VT1B) in, 134:1387
Verotoxin, B subunit of (VT1B), from enterohemorrhagic *Escherichia coli*, responsible for attachment of holotoxin to cell surface, 134:1387
Very late antigen (VLA)-4, transient activation and deactivation of, may facilitate transendothelial diapedesis, 134:1063
Vesicle recycling, synaptic, total pool, reduced in synapsin I–deficient mice, 134:1219
Vesicle repriming, synaptic, normal in synapsin I–deficient mice, 134:1219
Vesicle scission, requirements of second stage of post-Golgi vesicle formation, role of PKC in, 135:355
Vesicle-associated membrane protein-2 (VAMP-2), segregated from recycling endosomes in insulin-sensitive cells, 134:625
Vesicle-associated membrane protein/synaptobrevin, isoforms 1 and 2 of, widely and differentially expressed in nonneuronal tissues, 132:167
Vesicles
  cycle of synaptic, single vesicle budding step involving clathrin and dynamin, 133:1237
  endosomal βCOP involved in pH-dependent formation of transport, destined for late endosomes, 133:29
  novel class of clathrin-coated, budding from endosomes, 132:21
  synaptic, recycling pathways of, in active and nonactive zones, 135:797
  targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes, 132:277
Vesicular carriers, structure and organization in endoplasmic reticulum export complexes, COPII-coated, 135:19
vesicular SNAP receptor (vSNARE), synaptophysin binding to, 134:731
Vesicular stomatitis virus G protein, transport to cell surface, signal mediated in polarized and nonpolarized cells, 133:543
Vesicular transport
  association of dynamin-like protein with Golgi apparatus in mammalian cells, 133:761
  of MHC class II molecules, from lysosomal structures to plasma membrane, 135:611
Vimentin
  disrupted glial fibrillary acidic protein network in astrocytes from vimentin knockout mice, 133:853
  filaments, decoration of, with plectin repeat 5 domain, 134:1455
  mitosis-specific phosphorylation of, protein kinase C coupled with reorganization of intracellular membranes, 133:141
Vimentin intermediate filaments, disruption of state of assembly by microinjection of 1A peptides in cultured cells, 134:971
Vinculin
  cell adhesiveness in endoplasmic reticulum controlled by calreticulin-regulated expression of, 135:1913
  in filopodial motility, 134:1197
  interaction domain on paxillin of, 135:1109
Viral genomes, transport to nuclear domain (ND)10, does not require viral gene expression, 134:815
Viral RNA, localization of polyadenylated RNA and snRNP and, 135:303
Virus, Gag precursor of Moloney murine leukemia, targeting to budding site of, 135:1841
Virus replication sites, DNA, 134:815
Visualization, new method of, of chromatin organization and DNA sequences using lac operator-repressor staining, 135:1685
Vitronectin
  bead binding, is integrin-associated protein (IAP) and integrin dependent, 134:1313
  interaction of urokinase type plasminogen activator receptor (uPAR) with, promotes cell adhesion in extracellular matrix, 134:1563
  of melanoma cells, effects of thrombospondin through integrin-associated protein on, 135:533
  spreading of β3 or β5 cells on, 134:1323
v-SNARE, implicated in intra-Golgi transport, 133:507
Wallerian degeneration, granulocyte macrophage colony stimulating factor produced in lesioned peripheral nerves induces up-regulation of cell surface expression of MAC-2 by macrophages and Schwann cells, 133:159
Weaver granule neurons, rescue mechanisms of, 134:477
Wing hairs, in *Drosophila*, roles of Cdc42 and Rac1 in formation of, 135:1277
wingless protein, in *Drosophila*, role of glycosaminoglycans in transmission of, 135:819
Wingless signaling, role of Armadillo in, 134:1283
Wnt genes, activities of *Wnt*-1 class of secreted signaling factors are antagonized by *Wnt*-5A class and by dominant negative cadherin in early *Xenopus* development, 133:1123
Wortmannin, potential sites of PI-3 kinase function in endocytic pathway revealed by, PI-3 kinase inhibitor, 132:595
Wound healing
  in chick embryo, actin purse-string assembly during, roles of cadherin, myosin and Rho during, 135:1097
  and epidermal growth factor–induced changes in $\alpha_6\beta_4$ function, 134:241
  re-epithelialization after skin injury correlates with reorganization of keratin filaments in wound edge keratinocytes, potential role for keratin 16, 132:381
Wounding, extracellular matrix alters PDGF regulation of fibroblast integrins, 132:239
wt-p53, endogenous, of C2C212 cells, transcriptional activity during myogenesis inhibited by p53 dominant-negative expression, 134:193
X chromosome, XIST RNA paints the inactive, interphase, novel RNA involved in nuclear/chromosome structure, 132:259
Xanthosine triphosphate, GTPase distinct from Ran involved in nuclear protein import, 133:971
Xenopus
  activities of *Wnt*-1 class of secreted signaling factors are antagonized by *Wnt*-5A class and by dominant negative cadherin in early development in, 133:1123
  binding to cadherins antagonizes signaling activity of β-catenin during axis formation in, 132:1105
  embryonic cell adhesion to fibronectin, 134:227
  embryos, calcium dynamics in, changes in cleavage stage of, 135:181
  metazoan rDNA enhancer acts by making more genes transcriptionally active, 133:943
  nonmuscle myosin heavy chain isoforms, differences in subcellular localizations and enzymatic activities, 134:675
  nontetrameric species is major soluble form of keratin in oocytes of, rabbit reticulocyte lysates, 132:153
  oligomerization and maturation of Na,K-ATPase, interaction of cytoplasmic NH_2 terminus of β subunit with α subunit, 133:1193
  site-specific initiation of DNA replication dictated by chromosome structure in, 135:1207

Xenopus laevis
actin-depolymerizing factor/cofilin in, a phosphorylation-regulated protein essential for development, 132:871
influence of M-phase chromatin on anisotropy of microtubule asters, 133:125
nuclear proteins of quiescent cells in, inhibit DNA replication in intact and permeabilized nuclei, 133:955
XIST gene, paints the inactive X chromosome in interphase, novel RNA involved in nuclear/chromosome structure, 132:259
Yeast
aberrantly segregating centromeres activate spindle assembly checkpoint in budding, 133:75
associations among PH and SH3 domain–containing proteins and Rho-type GTPases in, 133:879
Bud4p, level of, regulated in cell cycle–dependent manner, 134:413
endocytosis *dim* mutants identified through novel genetic screen of, 135:1485
movement of cortical actin patches in, 132:861
nucleoporin Nup188p interacts genetically and physically with core structures of nuclear pore complex in, 133:1153
receptor-mediated transport of soluble misfolded proteins to vacuoles in, 135:623
reduced levels of **a**-factor mating pheromone results in cell fusion defective mutants in, 135:1727
refolding activity of heat shock proteins Ssa1/2, protein translocation in, 135:1229
src homology region 3 domain-binding proteins in, involved in bud formation, 133:865
targeting and fusion of ER-derived transport vesicles with purified Golgi membranes of, 132:277
Yeast actin–related protein, Arp2p, in late stage of cell cycle, 134:117
Yeast peroxisomal transporter, Pmp47, in transport and folding of dihydroxyacetone synthase (DHAS), 134:37
Z band, three-dimensional structure of, normal mammalian skeletal muscle, 133:571
Zona pellucida, acrosome reactions triggered by contact with, during mammalian fertilization, 134:637
zonula adherens, early stages in formation, and *armadillo*, *bazooka*, and *stardust*, 134:149
Zonula occludens, novel type of protein in, associated with tight junctions of polar epithelia and Sertoli cells of testis but not vascular endothelia, 134:1003
Zonula Occludens-1, and catenins form complex during early stages in assembly of tight junctions, 132:451
zw10, mutations in, disrupt chromosome behavior in male meiosis I and II, 134:1127